

## **Instructions for using the workbook**

The information required for the exposure assessment needs to be entered in the worksheet "**Data entry**".

In the following worksheets formulas calculate the exposure values automatically

Worksheet "**Operator Outdoor Spray AOEM**" is to be for outdoor spray applications. PPE options can be selected in this worksheet

Worksheet "**Operator Granules**" is for granular applications. Currently the calculator does not allow operator exposure for indoor applications. PPE options can be selected in this worksheet

Worksheets "**Resident exposure**" and "**Bystander exposure**" are only relevant for outdoor applications

Worksheet "**Recreational Exposure**" is only applicable for golf course, turf, other sports lawns or amenity turf/grassland areas where members of the public are likely to have access

The combined results of the exposure assessment are presented in worksheet "**Summary**"

This calculator should be used in conjunction with the **Guidance on the Assessment of Exposure for Operators, Workers, Residents and Bystanders in Risk Assessment for Plant Protection Products**

Latest version: 23 Oct 2014 - Version produced to support guidance document published 23/10/2014

Note: Some drop-down menus depend on others. To avoid errors, please fill-in from top to bottom

<b>Substance name</b>	trifloxystrobin
<b>Product name</b>	Luna Sensation
<b>Reference value non acutely toxic active substance (RVNAS)</b>	0,06 mg/kg bw/day
<b>Reference value acutely toxic active substance (RVAAS)</b>	0,06 mg/kg bw/day
<b>Crop type</b>	Ornamentals
<b>Substance properties</b>	
Formulation type	Soluble concentrates, emulsifiable concentrate, etc.
Minimum volume water for application (liquids)	150 L/ha
Maximum application rate of active substance	0,075 kg a.s. /ha
50% Dissipation Time DT50	30 days
Initial Dislodgeable Foliar Residue	3 µg/cm <sup>2</sup> of foliage/kg a.s. applied/ha
Dermal absorption of product	1,60%
Dermal absorption of in-use dilution	15,00%
Oral absorption of active substance	60,00%
Inhalation absorption of active substance	100,00%
Vapour pressure of active substance	moderately volatile substances with a vapour pressure between 5*10 <sup>-3</sup> Pa and 10 <sup>-2</sup> Pa
<b>Scenario</b>	
Indoor or Outdoor application	Outdoor
Application method	Downward spraying
Application equipment	Vehicle-mounted
Buffer strip	2-3 m
Number of applications	5
Interval between multiple applications	7 days
Season (upward spraying orchards only)	not relevant

**Exposure assessment**

Substance	trifloxystrobin	Formulation = Soluble concentrates, emulsifiable concentrate, etc.	Application rate-0,075 kg a.s. /ha	Spray dilution = 0,5 g a.s./l	Vapour pressure = moderately volatile substances with a vapour pressure between 5*10-3Pa and 10-2Pa
Scenario	Ornamentals / Outdoor / Downward spraying / Vehicle-mounted			Buffer = 2-3	Number applications = 5, Application interval = 7 days
Percentage Absorption	Dermal for product = 1,6	Dermal for in use dilution = 15	Oral = 60	Inhalation = 100	
RVNAS	0,06 mg/kg bw/day		RVAAS	0,06 mg/kg bw/day	
DFR	3 µg a.s./cm2 per kg a.s./ha		DT50	30 days	

<b>Operator Model</b>	Mixing, loading and application AOEM				
Potential exposure	Longer term systemic exposure mg/kg bw/day		0,0092	% of RVNAS	15,32%
	Acute systemic exposure mg/kg bw/day		0,0367	% of RVAAS	61,13%
Mixing and Loading	Gloves = No		Clothing = Work wear - arms, body and legs covered	RPE = None	Soluble bags = No
Application	Gloves = No		Clothing = Work wear - arms, body and legs covered	RPE = None	Closed cabin = No
Exposure (including PPE options above)	Longer term systemic exposure mg/kg bw/day		0,0043	% of RVNAS	7,17%
	Acute systemic exposure mg/kg bw/day		0,0138	% of RVAAS	23,03%

<b>Worker - Cutting, sorting, bundling, carrying</b>	Potential exposure mg/kg bw/day	0,2340	% of RVNAS	389,92%
	Working clothing mg/kg bw/day	0,0836	% of RVNAS	139,26%
	Working clothing and gloves mg/kg bw/day	0,0234	% of RVNAS	38,99%

<b>Resident - child</b>	Spray drift (75th percentile) mg/kg bw/day	0,0020	% of RVNAS	3,37%
	Vapour (75th percentile) mg/kg bw/day	0,0161	% of RVNAS	26,75%
	Surface deposits (75th percentile) mg/kg bw/day	0,0007	% of RVNAS	1,24%
	Entry into treated crops (75th percentile) mg/kg bw/day	0,0070	% of RVNAS	11,75%
	All pathways (mean) mg/kg bw/day	0,0233	% of RVNAS	38,87%
<b>Resident - adult</b>	Spray drift (75th percentile) mg/kg bw/day	0,0005	% of RVNAS	0,80%
	Vapour (75th percentile) mg/kg bw/day	0,0035	% of RVNAS	5,75%
	Surface deposits (75th percentile) mg/kg bw/day	0,0003	% of RVNAS	0,47%
	Entry into treated crops (75th percentile) mg/kg bw/day	0,0039	% of RVNAS	6,53%
	All pathways (mean) mg/kg bw/day	0,0070	% of RVNAS	11,68%

<b>Bystander - child</b>	Spray drift (95th percentile) mg/kg bw/day	0,0046	% of RVAAS	7,68%
	Vapour (95th percentile) mg/kg bw/day	0,0161	% of RVAAS	26,75%
	Surface deposits (95th percentile) mg/kg bw/day	0,0022	% of RVAAS	3,67%
	Entry into treated crops (95th percentile) mg/kg bw/day	0,0070	% of RVAAS	11,75%
<b>Bystander - adult</b>	Spray drift (95th percentile) mg/kg bw/day	0,0012	% of RVAAS	2,07%

### Exposure assessment

Vapour (95th percentile) mg/kg bw/day	0,0035	% of RVAAS	5,75%
Surface deposits (95th percentile) mg/kg bw/day	0,0009	% of RVAAS	1,43%
Entry into treated crops (95th percentile) mg/kg bw/day	0,0039	% of RVAAS	6,53%

<b>Recreational Exposure</b>	Child % of RVNAS	Adult % of RVNAS
------------------------------	------------------	------------------

Latest version: 23 Oct 2014 - Version produced to support guidance document published 23/10/2014

**Operator exposure for Luna Sensation outdoor spray applications**

Application rate of active substance	0,075 kg a.s./ha	<i>i_AppRate</i>
Assumed area treated	10 ha/day	<i>d_AreaTreated</i>
Amount of active substance applied	0,75 kg a.s./day	<i>i_AmountAS</i>
Dermal absorption of the product	1,60%	<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	15,00%	<i>i_AbsorInuse</i>
Formulation type	Soluble concentrates, emulsifiable concentrate, etc.	
Indoor or Outdoor application	Outdoor	
Application method	Downward spraying	
Application equipment	Vehicle-mounted	
Season	not relevant	

	Exposure values	µg exposure/day mixed and loaded		Reference	Comment
		75 <sup>th</sup> centile	95 <sup>th</sup> centile		
Mixing and loading	Hands	3892	14225	AOEM	
	Body	2914	66249	AOEM	
	Head	39	3768	AOEM	
	Protected hands (gloves)	29	149	AOEM	
	Protected body (workwear or protective garment and sturdy footwear)	18	110	AOEM	
	Protected head (hood and face shield)	1	213	AOEM	
	Inhalation	3	29	AOEM	
	<b>Protective Equipment</b>	Select for inclusion		Penetration factor	Inhalation Protection factor
	Gloves	No			
	Clothing	Work wear - arms, body and legs covered		Incl. in AOEM model	
Head and respiratory PPE	None		1	1	
Water soluble bag	No		1		

	Exposure values	µg exposure/day applied		Reference	Comment
		75 <sup>th</sup> centile	95 <sup>th</sup> centile		
Application	Hands	1215	3036	AOEM	This scenario assumes that small area equipment is used
	Body	1667	2113	AOEM	
	Head	10	117	AOEM	
	Protected hands (gloves)	19	21	AOEM	
	Protected body (workwear or protective garment and sturdy footwear)	21	25	AOEM	
	Inhalation	5	34	AOEM	
	<b>Protective Equipment</b>	Select for inclusion		Penetration factor	Inhalation Protection factor
	Gloves	No			
	Clothing	Work wear - arms, body and legs covered		Incl. in AOEM model	
	Head and respiratory PPE	None		1	1
Closed cab	No		vehicle mounted upward spraying only		

**1. Total**

	Without RPE/PPE	With RPE/PPE
<b>Longer term</b>		
Total systemic exposure from mixing, loading and application (mg a.s./day)	0,5514089	0,2581023
Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	0,0091901	0,0043017
% of RVNAS	15,32%	7,17%
<b>Acute</b>		
Total systemic exposure from mixing, loading and application (mg a.s./day)	2,2005525	0,8290992

Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	0,0366759	0,0138183
% of RVAAS	61,13%	23,03%

## 2. Longer term exposure

### 2.1 Mixing and loading

	Systemic exposure [ $\mu\text{g a.s. /day}$ ]	Systemic exposure [ $\mu\text{g a.s./kg}$ ]	Formula
<b>Without RPE/PPE</b>			
Hands	62,2728002	1,0378800	D15*i_AbsorpProduct
Body	46,6258537	0,7770976	D16*i_AbsorpProduct
Head	0,6226031	0,0103767	D17*i_AbsorpProduct
Inhalation	3,3979598	0,0566327	D21*i_AbsorpInhalation
Sum	112,9192167	1,8819869	
<b>With RPE/PPE (as selected above)</b>			
Hands	62,2728002	1,0378800	D18*i_AbsorpProduct
Body	0,2946918	0,0049115	D19*i_AbsorpProduct or D15*i_AbsorpProduct*F24
Head	0,6226031	0,0103767	D20*i_AbsorpProduct or D17*i_AbsorpProduct*F25
Inhalation	3,3979598	0,0566327	D21*i_AbsorpInhalation*G25
Sum	66,5880548	1,1098009	
Water soluble bag	66,5880548	1,1098009	C70*F26

### 2.2 Application

	Systemic exposure [ $\mu\text{g a.s. /day}$ ]	Systemic exposure [ $\mu\text{g a.s./kg}$ ]	Formula
<b>Without RPE/PPE</b>			
Hands	182,2867540	3,0381126	D30*i_AbsorpInuse
Body	250,1031691	4,1683862	D31*i_AbsorpInuse
Head	1,4974241	0,0249571	D32*i_AbsorpInuse
Inhalation	4,6023745	0,0767062	D35*i_AbsorpInhalation
Sum	438,4897217	7,3081620	
<b>With RPE/PPE (as selected above)</b>			
Hands	182,2867540	3,0381126	D33*i_AbsorpInuse
Body	3,1277112	0,0521285	D34*i_AbsorpInuse or D31*i_AbsorpInuse*F38
Head	1,4974241	0,0249571	D32*i_AbsorpInuse*F39
Inhalation	4,6023745	0,0767062	D35*i_AbsorpInuse*G39
Sum	191,5142637	3,1919044	

## 3. Acute exposure

### 3.1 Mixing and loading

	Systemic exposure [ $\mu\text{g a.s. /day}$ ]	Systemic exposure [ $\mu\text{g a.s./kg}$ ]	Formula
<b>Without RPE/PPE</b>			
Hands	227,5940626	3,7932344	E15*i_AbsorpProduct
Body	1059,9898161	17,6664969	E16*i_AbsorpProduct
Head	60,2880513	1,0048009	E17*i_AbsorpProduct
Inhalation	28,5186078	0,4753101	E21*i_AbsorpInhalation
Sum	1376,3905378	22,9398423	
<b>With RPE/PPE (as selected above)</b>			
Hands	227,5940626	3,7932344	E18*i_AbsorpProduct
Body	1,7550167	0,0292503	E19*i_AbsorpProduct or E16*i_AbsorpProduct*F24
Head	60,2880513	1,0048009	E20*i_AbsorpProduct or E17*i_AbsorpProduct*F25
Inhalation	28,5186078	0,4753101	E21*i_AbsorpInhalation*G25
Sum	318,1557384	5,3025956	
Water soluble bag	318,1557384	5,3025956	C104*F26

### 2.2 Application

	Systemic exposure [ $\mu\text{g a.s. /day}$ ]	Systemic exposure [ $\mu\text{g a.s./kg}$ ]	Formula
<b>Without RPE/PPE</b>			
Hands	455,4030671	7,5900511	E30*i_AbsorpInuse
Body	316,9146815	5,2819114	E31*i_AbsorpInuse
Head	17,5780279	0,2929671	E32*i_AbsorpInuse
Inhalation	34,2661494	0,5711025	E35*i_AbsorpInhalation
Sum	824,1619259	13,7360321	
<b>With RPE/PPE (as selected above)</b>			
Hands	455,4030671	7,5900511	E33*i_AbsorpInuse
Body	3,6961685	0,0616028	E34*i_AbsorpInuse or E31*i_AbsorpInuse*F38
Head	17,5780279	0,2929671	E32*i_AbsorpInuse*F39

Inhalation	34,2661494	0,5711025	E35*1_Absorpinhalation*G39
Sum	510,9434129	8,5157235	

**Operator exposure for Luna Sensation granular applications**

Application rate of active substance	0,075 kg a.s./ha	<i>i_AppRate</i>
Assumed area treated	10 ha/day	<i>d_AreaTreated</i>
Amount of active substance applied	0,75 kg a.s./day	<i>i_AmountAS</i>
Dermal absorption of the product	1,60%	<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	15,00%	<i>i_AbsorInuse</i>
Formulation type	Soluble concentrates, emulsifiable concentrate, etc.	
Indoor or Outdoor application	Outdoor <b>This sheet is only to be used for granular applications</b>	
Application method	Downward spraying	
Application equipment	Vehicle-mounted	

	Exposure values	mg exposure/kg a.s. mixed and loaded		Reference	Comment
		75 <sup>th</sup> centile	95 <sup>th</sup> centile		
Mixing and loading	Hands	#N/A	#N/A	#N/A	#N/A
	Body	#N/A	#N/A	#N/A	#N/A
	Inhalation	#N/A	#N/A	#N/A	#N/A
	<b>Protective Equipment</b>	Choose item		Penetration factor	
	Gloves	Chemical resistant gloves			Protection for granules exposure is based on measured values
	Body PPE	Certified protective coverall			
	RPE	None		1	

	Exposure values	mg exposure/kg a.s. applied		Reference	Comment
		75 <sup>th</sup> centile	95 <sup>th</sup> centile		
Application	Hands	#N/A	#N/A	#N/A	#N/A
	Body	#N/A	#N/A	#N/A	#N/A
	Inhalation	#N/A	#N/A	#N/A	#N/A
	<b>Protective Equipment</b>	Choose item		Penetration factor	
	Gloves	Chemical resistant gloves			Protection for granules exposure is based on measured values
	Body PPE	Certified protective coverall			
	RPE	FP1, P1 and similar		0,25	

**1. Total**

	Without RPE/PPE	With RPE/PPE
<b>Longer term</b>		
Total systemic exposure from mixing, loading and application (mg a.s./day)	#N/A	#N/A
Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	#N/A	#N/A
% of RVNAS	#N/A	#N/A
<b>Acute</b>		
Total systemic exposure from mixing, loading and application (mg a.s./day)	#N/A	#N/A



Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	#N/A	#N/A	
% of RVAAS	#N/A	#N/A	

## 2. Longer term exposure

### 2.1 Mixing and loading

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
<b>Without RPE/PPE</b>			
Hands	#N/A	#N/A	$D14*100*i\_AmoutAS*i\_AbsorpProduct$
Body	#N/A	#N/A	$D15*100*i\_AmoutAS*i\_AbsorpProduct$
Inhalation	#N/A	#N/A	$D16*i\_AmoutAS*i\_AbsorpInhalation$
Sum	#N/A	#N/A	
<b>With RPE/PPE (as selected above)</b>			
Hands	#N/A	#N/A	$D14*i\_AmoutAS*i\_AbsorpProduct$
Body	#N/A	#N/A	$D15*i\_AmoutAS*i\_AbsorpProduct$
Inhalation	#N/A	#N/A	$D16*i\_AmoutAS*i\_AbsorpInhalation*F20$
Sum	#N/A	#N/A	

### 2.2 Application

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
<b>Without RPE/PPE</b>			
Hands	#N/A	#N/A	$D25*100*i\_AmoutAS*i\_AbsorpInuse$
Body	#N/A	#N/A	$D26*100*i\_AmoutAS*i\_AbsorpInuse$
Inhalation	#N/A	#N/A	$D27*i\_AmoutAS*i\_AbsorpInhalation$
Sum	#N/A	#N/A	
<b>With RPE/PPE (as selected above)</b>			
Hands	#N/A	#N/A	$D25*i\_AmoutAS*i\_AbsorpInuse$
Body	#N/A	#N/A	$D26*i\_AmoutAS*i\_AbsorpInuse$
Inhalation	#N/A	#N/A	$D27*i\_AmoutAS*i\_AbsorpInhalation*F31$
Sum	#N/A	#N/A	

## 3. Acute exposure

### 3.1 Mixing and loading

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
<b>Without RPE/PPE</b>			
Hands	#N/A	#N/A	$E14*100*i\_AmoutAS*i\_AbsorpProduct$
Body	#N/A	#N/A	$E15*100*i\_AmoutAS*i\_AbsorpProduct$
Inhalation	#N/A	#N/A	$E16*i\_AmoutAS*i\_AbsorpInhalation$
Sum	#N/A	#N/A	
<b>With RPE/PPE (as selected above)</b>			
Hands	#N/A	#N/A	$E14*100*i\_AmoutAS*i\_AbsorpProduct$
Body	#N/A	#N/A	$E15*100*i\_AmoutAS*i\_AbsorpProduct$
Inhalation	#N/A	#N/A	$E16*i\_AmoutAS*i\_AbsorpInhalation*F20$
Sum	#N/A	#N/A	

### 3.2 Application

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
<b>Without RPE/PPE</b>			
Hands	#N/A	#N/A	$E25*100*i\_AmoutAS*i\_AbsorpInuse$
Body	#N/A	#N/A	$E25*100*i\_AmoutAS*i\_AbsorpInuse$
Inhalation	#N/A	#N/A	$E26*i\_AmoutAS*i\_AbsorpInhalation$
Sum	#N/A	#N/A	
<b>With RPE/PPE (as selected above)</b>			

Hands	#N/A	#N/A	$E25*100*i\_AmountAS*i\_Absorpnuse$
Body	#N/A	#N/A	$E26*100*i\_AmountAS*i\_Absorpnuse$
Inhalation	#N/A	#N/A	$E27*i\_AmountAS*i\_Absorpinhalation*F31$
Sum	#N/A	#N/A	

## Worker exposure from residues on foliage for Luna Sensation

Crop type	Ornamentals	
Indoor or outdoor	Outdoor	
Application method	Downward spraying	
Application equipment	Vehicle-mounted	
Worker's task	Cutting, sorting, bundling, carrying	
Main body parts in contact with foliage	Hand and body	
Application rate of active substance	0,075 kg a.s./ha	<i>i_AppRate</i>
Number of applications	5	<i>i_AppNo</i>
Interval between multiple applications	7 days	<i>i_AppInt</i>
Half-life of active substance	30 days	<i>d_HalfLifeAS</i>
Multiple application factor	3,7	<i>d_MAF</i>
Dermal absorption of the product	1,60%	<i>i_AbsorpProduct</i>
Dermal absorption of the in-use dilution	15,00%	<i>i_AbsorpInuse</i>
Dislodgeable foliar residue ( $i\_AppRate * i\_DFR$ )	0,225 µg a.s./cm <sup>2</sup>	<i>d_DFR</i>
Working hours	8 hr	<i>d_WorkHr</i>
Dermal transfer coefficient - Total potential exposure	14000 cm <sup>2</sup> /hr	<i>d_DermTcUCV</i>
Dermal transfer coefficient - arms, body and legs covered	5000 cm <sup>2</sup> /hr	<i>d_DermTcCV1</i>
Dermal transfer coefficient - hands, arms, body and legs covered	1400 cm <sup>2</sup> /hr	<i>d_DermTcCV2</i>
Inhalation transfer coefficient for automated applications	NA ha/hr*10 <sup>-3</sup>	<i>d_InhalTcAut</i>
Inhalation transfer coefficient for cutting ornamentals	NA ha/hr*10 <sup>-3</sup>	<i>d_InhalTcCut</i>
Inhalation transfer coefficient for sorting / bundling ornamentals	NA ha/hr*10 <sup>-3</sup>	<i>d_InhalTcSort</i>

### 1. Total

	Potential exposure	Work wear - arms, body and legs covered	Working wear and gloves	Comments
Total systemic exposure (mg a.s./day)	14,0371096	5,0132534	1,4037110	
Total systemic exposure per kg body weight (mg/kg bw/day)	0,2339518	0,0835542	0,0233952	
% of RVNAS	389,92%	139,26%	38,99%	

### 2. Details

	Systemic exposure		Formula	Comments
	[mg a.s. /day]	[mg a.s./kg bw/day]		
Dermal - Potential	14,0371096	0,2339518	$d\_DermTcUCV * d\_WorkHr * i\_DFR * i\_MAF / 1000 * i\_AbsorpInuse$	
Dermal - Work wear - arms, body and legs covered	5,0132534	0,0835542	$d\_DermTcCV1 * d\_WorkHr * d\_DFR * d\_MAF / 1000 * i\_AbsorpInuse$	
Dermal - Working wear and gloves	1,4037110	0,0233952	$d\_DermTcCV2 * d\_WorkHr * d\_DFR * d\_MAF / 1000 * i\_AbsorpInuse$	
Inhalation				Na for outdoor activities

## Resident exposure for Luna Sensation

Croptype	Ornamentals	
Application method	Downward spraying	
Application equipment	Vehicle-mounted	<i>i_AppEquip</i>
Formulation type	Soluble concentrates, emulsifiable concentrate, etc.	<i>i_FormVal</i>
Buffer strip	2-3 m	<i>i_Buffer</i>
Application rate of the product	0,075 kg a.s./ha	<i>i_AppRate</i>
Concentration of active substance (in-use dilution for liquid applications)	0,5 g a.s./l	<i>d_ConcAS</i>
Dermal absorption of product	1,60%	<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	15,00%	<i>i_AbsorpInuse</i>
Oral absorption	60,00%	<i>i_AbsorpOrallnuse</i>
Dislodgeable foliar residue ( $i\_AppRate * i\_DFR$ )	0,225 µg a.s./cm <sup>2</sup>	<i>d_DFR</i>
Vapour pressure of in-use dilution	moderately volatile substances with a vapour pressure between 5*10-3Pa and 10-2Pa	<i>i_Volat</i>
Concentration in air	0,015 mg/m <sup>3</sup>	<i>d_AirCon</i>
Resident dermal spray drift exposure 75th percentile - adult	0,47 ml spray dilution/person	
Resident dermal spray drift exposure 75th percentile - child	0,327 ml spray dilution/person	
Resident inhal. spray drift exposure 75th percentile - adult	0,00010 ml spray dilution/person	
Resident inhal. spray drift exposure 75th percentile - child	0,00022 ml spray dilution/person	
Resident dermal spray drift exposure mean - adult	0,22318 ml spray dilution/person	
Resident dermal spray drift exposure mean - child	0,18 ml spray dilution/person	
Resident inhal. spray drift exposure mean - adult	0,00009 ml spray dilution/person	
Resident inhal. spray drift exposure mean - child	0,00017 ml spray dilution/person	
Exposure duration dermal	2 hours	<i>d_ReExpDur</i>
Exposure duration inhalation	24 hours	<i>d_ReExpDurInhal</i>
Exposure duration entry into treated crops	0,25 hours	<i>d_ExpDurTreatCrop</i>
Light clothing adjustment factor	18,0%	<i>d_ClothAF</i>
Breathing rate adult	0,23 m <sup>3</sup> /day/kg	<i>d_BreathRAAd</i>
Breathing rate child (1-3 year old)	1,07 m <sup>3</sup> /day/kg	<i>d_BreathRCh</i>
Drift percentage on surface (75th percentile)	5,60%	
Drift percentage on surface (mean)	4,10%	
Turf transferable residues percentage	5,00%	<i>d_Turf</i>
Transfer coeff. of surface deposits-adult	7300 cm <sup>2</sup> /hour	<i>d_ReTCAd</i>
Transfer coeff. of surface deposits-child (1-3 year old)	2600 cm <sup>2</sup> /hour	<i>d_ReTCCh</i>
Saliva extraction percentage	50,00%	<i>d_SalExt</i>
Surface area of hands mouthed	20 cm <sup>2</sup>	<i>d_AreaHM</i>
Frequency of hand to mouth activity	9,5 events/hour	<i>d_ReFreqHM</i>
Ingestion rate for mouthing of grass per day	25 cm <sup>2</sup>	<i>d_MouthGrass</i>
Dislodgeable residues percentage transferability for object to mouth	20,00%	<i>d_DRP</i>
Transfer coefficient for entry into treated crops (75th percentile) - adult	7500 cm <sup>2</sup> /h	<i>d_TcEntryAd</i>
Transfer coefficient for entry into treated crops (75th percentile) - child	2250 cm <sup>2</sup> /h	<i>d_TcEntryCh</i>
Transfer coefficient for entry into treated crops (mean) - adult	5980 cm <sup>2</sup> /h	<i>d_TcEntryAd</i>
Transfer coefficient for entry into treated crops (mean) - child	1794 cm <sup>2</sup> /h	<i>d_TcEntryCh</i>

1. Total

1.1 1-3 year old child

	Spray drift (75th percentile)	Vapour (75th percentile)	Surface deposits (75th percentile)	Entry into treated crops (75th percentile)	All pathways (mean)
Total systemic exposure (mg a.s./day)	0,0202205	0,1605000	0,0074397	0,0704989	0,2332280
Total systemic exposure per kg body weight (mg/kg bw/day)	0,0020221	0,0160500	0,0007440	0,0070499	0,0233228
% of RVNAS	3,37%	26,75%	1,24%	11,75%	38,87%

1.2 Adult

	Spray drift	Vapour	Surface deposits	Entry into treated crops	All pathways (mean)
Total systemic exposure (mg a.s./day)	0,0289550	0,2070000	0,0170785	0,2349963	0,4206448
Total systemic exposure per kg body weight (mg/kg bw/day)	0,0004826	0,0034500	0,0002846	0,0039166	0,0070107
% of RVNAS	0,80%	5,75%	0,47%	6,53%	11,68%

2. Resident exposure 75th Percentile

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
<b>1-3 year old child</b>				
Spray drift	0,0202205	0,0020221	$((C16 * i\_AbsorpInuse * (1 - d\_ClothAF)) + C18) * d\_ConcAS$	
Vapour	0,1605000	0,0160500	$d\_AirCon * d\_BreathRCh * d\_BwChild$	
Surface deposits				
Dermal	0,0060827	0,0006083	$(i\_AppRate / 100) * C29 * d\_Turf * d\_ReTCCh * d\_ReExpDur * MAX(i\_AbsorpProduct, i\_AbsorpInuse) * d\_MAF * IF(i\_AppEquip = "Vehicle-mounted-Drift Reduction", 0.5, 1)$	
Hand to mouth	0,0008890	0,0000889	$(i\_AppRate / 100) * C29 * d\_Turf * d\_SalExt * d\_AreaHM * d\_ReFreqHM * d\_ReExpDur * i\_AbsorpOrallnuse * d\_MAF$	
Object to mouth	0,0004679	0,0000468	$(i\_AppRate / 100) * C29 * d\_DRP * d\_MouthGrass * i\_AbsorpOrallnuse * d\_MAF$	

Entry into treated crops				
Dermal	0,0704989	0,0070499	$(d\_TcEntryCh*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_Absorplnuse)$	
Hand to mouth			$(i\_AppRate/100)*d\_Turf*d\_MAF*d\_SalExt*d\_AreaHM*d\_ReFreqHM*d\_ReExpDur*i\_AbsorpOrallnuse$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Object to mouth			$(i\_AppRate/100)*d\_DRP*d\_MouthGrass*i\_AbsorpOrallnuse*d\_MAF$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
<b>Adult</b>				
Spray drift	0,0289550	0,0004826	$(C15*i\_Absorplnuse*(1-d\_ClothAF))+C17)*d\_ConcAS$	
Vapour	0,2070000	0,0034500	$d\_AirCon*d\_BreathRAD*d\_BwAdult$	
Surface deposits (dermal)	0,0170785	0,0002846	$(i\_AppRate/100)*C30*d\_Turf*d\_ReTCAd*d\_ReExpDur*i\_Absorplnuse$	
Entry into treated crops (dermal)	0,2349963	0,0039166	$(d\_TcEntryAd*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_Absorplnuse)$	

### 3. Summing of exposure pathways mean

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
<b>1-3 year old child</b>				
Spray drift	0,0110700	0,0011070	$((C20*i\_Absorplnuse*(1-d\_ClothAF))+C22)*d\_ConcAS$	
Vapour	0,1605000	0,0160500	$d\_AirCon*d\_BreathRCh*d\_BwChild$	
Surface deposits				
Dermal	0,0044534	0,0004453	$(i\_AppRate/100)*C30*d\_Turf*d\_ReTCCh*d\_ReExpDur*MAX(i\_AbsorpProduct,i\_Absorplnuse)*d\_MAF*IF(i\_AppEquip = "Vehicle-mounted-Drift Reduction",0.5,1))$	
Hand to mouth	0,0006509	0,0000651	$(i\_AppRate/100)*C30*d\_Turf*d\_SalExt*d\_AreaHM*d\_ReFreqHM*d\_ReExpDur*i\_AbsorpOrallnuse*d\_MAF$	
Object to mouth	0,0003426	0,0000343	$(i\_AppRate/100)*C30*d\_DRP*d\_MouthGrass*i\_AbsorpOrallnuse*d\_MAF$	
Entry into treated crops				
Dermal	0,0562111	0,0056211	$(d\_TcEntryMeanCh*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_Absorplnuse))$	
Hand to mouth			$(i\_AppRate/100)*1*d\_Turf*d\_MAF*d\_SalExt*d\_AreaHM*d\_ReFreqHM*d\_ReExpDur*i\_AbsorpOrallnuse$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Object to mouth			$(i\_AppRate/100)*1*d\_DRP*d\_MouthGrass*i\_AbsorpOrallnuse*d\_MAF$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
<b>Adult</b>				
Spray drift	0,0137706	0,0002295	$"(C19*i\_Absorplnuse*(1-d\_ClothAF))+C21)*d\_ConcAS"$	
Vapour	0,2070000	0,0034500	$d\_AirCon*d\_BreathRAD*d\_BwAdult$	
Surface deposits (dermal)	0,0125039	0,0002084	$(i\_AppRate/100)*C30*d\_Turf*d\_ReTCAd*d\_ReExpDur*MAX(i\_AbsorpProduct,i\_Absorplnuse)*d\_MAF*IF(i\_AppEquip = "Vehicle-mounted-Drift Reduction",0.5,1)$	
Entry into treated crops (dermal)	0,1873703	0,0031228	$(d\_TcEntryMeanAd*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_Absorplnuse)$	

**Bystander exposure for Luna Sensation**

Croptype	<b>Ornamentals</b>		
Application method	<b>Downward spraying</b>		
Application equipment	<b>Vehicle-mounted</b>		<i>i_AppEquip</i>
Formulation type	<b>Soluble concentrates, emulsifiable concentrate, etc.</b>		
Application rate of the product	<b>0,075</b> kg a.s./ha		<i>i_AppRate</i>
Buffer strip	<b>2-3</b> m		<i>i_Buffer</i>
Concentration of active substance (in-use dilution for liquid applications)	<b>0,5</b> g a.s./l		<i>d_ConcAS</i>
Dermal absorption of product	<b>1,60%</b>		<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	<b>15,00%</b>		<i>i_AbsorpInuse</i>
Oral absorption	<b>60,00%</b>		<i>i_AbsorpOrallnuse</i>
Dislodgeable foliar residue (i_AppRate*i_DFR)	<b>0,225</b> µg a.s./cm <sup>2</sup>		<i>d_DFR</i>
Vapour pressure of in-use dilution	<b>moderately volatile substances with a vapour pressure between 5*10-3Pa and 10-2Pa</b> Pa		<i>i_Volat</i>
Concentration in air	<b>0,015</b> mg/m <sup>3</sup>		<i>d_AirCon</i>
Bystander dermal spray drift exposure - adult	<b>1,21</b> ml spray dilution/person		
Bystander dermal spray drift exposure - child	<b>0,74</b> ml spray dilution/person		
Bystander inhal. spray drift exposure - adult	<b>0,00050</b> ml spray dilution/person		
Bystander inhal. spray drift exposure - child	<b>0,00112</b> ml spray dilution/person		
Exposure duration	<b>2</b> hours		<i>d_ByExpDur</i>
Exposure duration entry into treated crops	<b>0,25</b> hours		<i>d_ExpDurTreatCrop</i>
Light clothing adjustment factor	<b>18,0%</b>		<i>d_ClothAF</i>
Breathing rate adult	<b>0,23</b> m <sup>3</sup> /hours/kg		<i>d_BreathRAd</i>
Breathing rate child (1-3 year old)	<b>1,07</b> m <sup>3</sup> /hours/kg		<i>d_BreathRCh</i>
Drift percentage on surface (90th percentile)	<b>8,50%</b>		
Turf transferable residues percentage	<b>5,00%</b>		<i>d_Turf</i>
Transfer coeff. of surface deposits-adult	<b>14500</b> cm <sup>2</sup> /hour		<i>d_ByTCAd</i>
Transfer coeff. of surface deposits-child (1-3 year old)	<b>5200</b> cm <sup>2</sup> /hour		<i>d_ByTCCh</i>
Saliva extraction percentage	<b>50,00%</b>		<i>d_SalExt</i>
Surface area of hands mouthed	<b>20</b> cm <sup>2</sup>		<i>d_AreaHM</i>
Frequency of hand to mouth activity	<b>20</b> events/hour		<i>d_ByFreqHM</i>
Ingestion rate for mouthing of grass per day	<b>25</b> cm <sup>2</sup>		<i>d_MouthGrass</i>
Dislodgeable residues percentage transferability for object to mouth	<b>20,00%</b>		<i>d_DRP</i>
Transfer coefficient for entry into treated crops - adult	<b>7500</b> cm <sup>2</sup> /h		<i>d_TcEntryAd</i>
Transfer coefficient for entry into treated crops - child	<b>2250</b> cm <sup>2</sup> /h		<i>d_TcEntryCh</i>

**1. Total**

**1.1 1-3 year old child**

	<b>Spray drift</b>	<b>Vapour</b>	<b>Surface deposits</b>	<b>Entry into treated crops</b>
Total systemic exposure (mg a.s./day)	<b>0,0460700</b>	<b>0,1605000</b>	<b>0,0220165</b>	<b>0,0704989</b>
Total systemic exposure per kg body weight (mg/kg bw/day)	<b>0,0046070</b>	<b>0,0160500</b>	<b>0,0022017</b>	<b>0,0070499</b>
% of RVAAS	<b>7,68%</b>	<b>26,75%</b>	<b>3,67%</b>	<b>11,75%</b>

**1.2 Adult**

	<b>Spray drift</b>	<b>Vapour</b>	<b>Surface deposits</b>	<b>Entry into treated crops</b>
Total systemic exposure (mg a.s./day)	<b>0,0746650</b>	<b>0,2070000</b>	<b>0,0514903</b>	<b>0,2349963</b>
Total systemic exposure per kg body weight (mg/kg bw/day)	<b>0,0012444</b>	<b>0,0034500</b>	<b>0,0008582</b>	<b>0,0039166</b>
% of RVAAS	<b>2,07%</b>	<b>5,75%</b>	<b>1,43%</b>	<b>6,53%</b>

2. Details

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
<b>1-3 year old child</b>				
Spray drift	0,0460700	0,0046070	$((C16*i\_Absorplnuse*(1-d\_ClothAF))+C18)*d\_ConcAS''$	
Vapour	0,1605000	0,0160500	$d\_AirCon*d\_BreathRCh*d\_BwChild$	
Surface deposits				
Dermal	0,0184655	0,0018465	$(i\_AppRate/100)*C24*d\_Turf*d\_ByTCCh*d\_ByExpDur*MAX(i\_AbsorpProduct,i\_Absorplnuse)*d\_MAF*IF(i\_AppEquip="Vehicle-mounted-Drift Reduction",0.5,1)$	
Hand to mouth	0,0028408	0,0002841	$(i\_AppRate/100)*C25*d\_Turf*d\_SalExt*d\_AreaHM*d\_ByFreqHM*d\_ByExpDur*i\_AbsorpOralnuse*d\_MAF$	
Object to mouth	0,0007102	0,0000710	$(i\_AppRate/100)*C25*d\_DRP*d\_MouthGrass*i\_AbsorpOralnuse*d\_MAF$	



Entry into treated crops				
Dermal	<b>0,0704989</b>	<b>0,0070499</b>	$(d\_TcEntryCh*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_Absorplnuse)$	
Hand to mouth			$(i\_AppRate/100)*d\_MAF*d\_Turf*d\_SalExt*d\_AreaHM*d\_ByFreqHM*d\_ByExpDur*i\_AbsorpOrallnuse$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Object to mouth			$(i\_AppRate/100)*d\_DRP*d\_MouthGrass*i\_AbsorpOrallnuse*d\_MAF$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
<b>Adult</b>				
Spray drift	<b>0,0746650</b>	<b>0,0012444</b>	$((C15*i\_Absorplnuse*(1-d\_ClothAF))+C17)*d\_ConcAS$	
Vapour	<b>0,2070000</b>	<b>0,0034500</b>	$d\_AirCon*d\_BreathRAD*d\_BwAdult$	
Surface deposits (dermal)	<b>0,0514903</b>	<b>0,0008582</b>	$(i\_AppRate/100)*C24*d\_Turf*d\_ByTCAd*d\_ByExpDur*MAX(i\_AbsorpProduct,i\_Absorplnuse)*d\_MAF*IF(i\_AppEquip="Vehicle-mounted-Drift Reduction",0.5,1)$	
Entry into treated crops (dermal)	<b>0,2349963</b>	<b>0,0039166</b>	$(d\_TcEntryAd*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_Absorplnuse)$	

## Recreational exposure for Luna Sensation

Croptype	Golf course, turf or other sports lawns	<b>This sheet is only to be used for treatment of grassland used for recreational purposes</b>	
Application method	Downward spraying		
Application equipment	Vehicle-mounted		<i>i_AppEquip</i>
Formulation type	Soluble concentrates, emulsifiable concentrate, etc.		<i>i_FormVal</i>
Application rate of the product	0,075 kg a.s./ha		<i>i_AppRate</i>
Dermal absorption of product	1,60%		<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	15,00%		<i>i_AbsorpInuse</i>
Oral absorption	60,00%		<i>i_AbsorpOrallnuse</i>
Dislodgeable foliar residue ( $i\_AppRate \cdot i\_DFR$ )	0,225 µg a.s./cm <sup>2</sup>		<i>d_DFR</i>
Exposure duration dermal	2 hours		<i>d_ReExpDur</i>
Light clothing adjustment factor Adult resident	18,0%		<i>d_ClothAF</i>
Drift percentage on surface	100,00%		
Turf transferable residues percentage	5,00%		<i>d_Turf</i>
Transfer coeff. of surface deposits-adult	7300 cm <sup>2</sup> /hour		<i>d_ReTCAd</i>
Transfer coeff. of surface deposits-child (1-3 year old)	2600 cm <sup>2</sup> /hour		<i>d_ReTCCh</i>
Saliva extraction percentage	50,00%		<i>d_SalExt</i>
Surface area of hands mouthed	20 cm <sup>2</sup>		<i>d_AreaHM</i>
Frequency of hand to mouth activity	9,5 events/hour		<i>d_ReFreqHM</i>
Ingestion rate for mouthing of grass per day	25 cm <sup>2</sup>		<i>d_MouthGrass</i>

### 2. Details

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
<b>1-3 year old child</b>				
Surface deposits				
Dermal	0,1086205	0,0108620	$(i\_AppRate/100) \cdot C13 \cdot d\_Turf \cdot d\_ReTCCh \cdot d\_ReExpDur \cdot \text{MAX}(i\_AbsorpProduct, i\_AbsorpInuse) \cdot d\_MAF$	
Hand to mouth	0,0158753	0,0015875	$(i\_AppRate/100) \cdot C13 \cdot d\_Turf \cdot d\_SalExt \cdot d\_AreaHM \cdot d\_ReFreqHM \cdot d\_ReExpDur \cdot i\_AbsorpOrallnuse \cdot d\_MAF$	
Object to mouth	0,0083554	0,0008355	$(i\_AppRate/100) \cdot C13 \cdot d\_DRP \cdot d\_MouthGrass \cdot i\_AbsorpOrallnuse \cdot d\_MAF$	
Total systemic exposure	0,1328512	0,0132851		
% of RVNAS				
<b>Adult</b>				
Surface deposits (dermal)	0,3049729	0,0050829	$(i\_AppRate/100) \cdot C13 \cdot d\_Turf \cdot d\_ReTCAd \cdot d\_ReExpDur \cdot \text{MAX}(i\_AbsorpProduct, i\_AbsorpInuse) \cdot d\_MAF$	
% of RVNAS				

<i>d_AirConVol</i>	Concentration in air of moderately volatile substances	<b>0,015</b> mg/m <sup>3</sup>
<i>d_AirConNonVol</i>	Concentration in air of low volatile substances	<b>0,001</b> mg/m <sup>3</sup>
<i>d_AreaHM</i>	Surface area of hands mouthed	<b>20</b> cm <sup>2</sup>
<i>d_AreaTreated</i>	Area treated (defined by crop type)	<b>10</b> ha
<i>d_BreathRAd</i>	Breathing rate adult residents	<b>0,23</b> m <sup>3</sup> /day/kg
<i>d_BreathRCh</i>	Breathing rate child (1-3 year old) residents	<b>1,07</b> m <sup>3</sup> /day/kg
<i>d_BwAdult</i>	Adult body weight	<b>60</b> kg
<i>d_BwChild</i>	Child body weight (1 to < 3 year olds)	<b>10</b> kg
<i>d_ByBreathRAd</i>	Breathing rate adult bystander	<b>0,04</b> m <sup>3</sup> /hours/kg
<i>d_ByBreathRCh</i>	Breathing rate child (1-3 year old) bystander	<b>0,19</b> m <sup>3</sup> /hours/kg
<i>d_ByExpDur</i>	Exposure duration intense activity breathing rates	<b>2</b> hours
<i>d_ByFreqHM</i>	Frequency of hand to mouth activity	<b>20</b> events/hour
<i>d_ByTCAd</i>	Transfer coeff. of surface deposits-adult	<b>14500</b> cm <sup>2</sup> /hour
<i>d_ByTCCh</i>	Transfer coeff. of surface deposits-child (1-3 year old)	<b>5200</b> cm <sup>2</sup> /hour
<i>d_ClothAF</i>	Light clothing adjustment factor resident and bystanders	<b>18,0%</b>
<i>d_ConcAs</i>	Concentration of active substance (in-use dilution for liquid applications)	<b>0,5</b> g a.s./l
<i>d_DFR</i>	Dislodgeable foliar residue (i_AppRate*i_DFR)	<b>0,225</b> µg a.s./cm <sup>2</sup>
<i>d_DRP</i>	Dislodgeable residues percentage transferability for object to mouth	<b>20,0%</b>
<i>d_HalfLifeAS</i>	Half-life of active substance (DT50)	<b>30</b> days
<i>d_InhalTcAut</i>	Inhalation transfer coefficient for automated applications	<b>NA</b> ha/hr*10 <sup>^(-3)</sup>
<i>d_InhalTcCut</i>	Inhalation transfer coefficient for cutting ornamentals	<b>NA</b> ha/hr*10 <sup>^(-3)</sup>
<i>d_InhalTcSort</i>	Inhalation transfer coefficient for sorting / bundling ornamentals	<b>NA</b> ha/hr*10 <sup>^(-3)</sup>
<i>d_MAF</i>	Multiple application factor	<b>3,71</b>
<i>d_MouthGrass</i>	Ingestion rate for mouthing of grass per day	<b>25</b> cm <sup>2</sup> grass/day
<i>d_ReExpDur</i>	Exposure duration resident dermal	<b>2</b> hours
<i>d_ReExpDurInhal</i>	Exposure duration resident inhalation	<b>24</b> hours
<i>d_ExpDurTreatCrop</i>	Exposure duration for resident and bystander entry into treated crops	<b>0,25</b> hours
<i>d_ReFreqHM</i>	Frequency of hand to mouth activity	<b>9,5</b> events/hour
<i>d_ReTCAd</i>	Transfer coeff. of surface deposits-adult	<b>7300</b> cm <sup>2</sup> /hour
<i>d_ReTCCh</i>	Transfer coeff. of surface deposits-child (1-3 year old)	<b>2600</b> cm <sup>2</sup> /hour
<i>d_SalExt</i>	Saliva extraction percentage	<b>50,0%</b>
<i>d_TcEntryAd</i>	Transfer coefficient for entry into treated crops 75th percentile - adult	<b>7500</b> cm <sup>2</sup> /h
<i>d_TcEntryCh</i>	Transfer coefficient for entry into treated crops 75th percentile - child	<b>2250</b> cm <sup>2</sup> /h
<i>d_TcEntryMeanAd</i>	Transfer coefficient for entry into treated crops mean - adult	<b>5980</b> cm <sup>2</sup> /h
<i>d_TcEntryMeanCh</i>	Transfer coefficient for entry into treated crops mean - child	<b>1794</b> cm <sup>2</sup> /h
<i>d_Turf</i>	Turf transferable residues percentage	<b>5,0%</b>
<i>d_PctExtrapolation</i>	For exposure value 75 percentiles above this amount linear extrapolation is performed	<b>1,5</b> kg
<i>d_head75ProtectionFactor</i>	Coefficient to estimate head protection factor 75 th Percentile	<b>1,79422</b>
<i>d_head95ProtectionFactor</i>	Coefficient to estimate head protection factor 95 Percentile	<b>1,24705</b>

*sys\_KeyOperator* Variables for operator exposure lookup key  
*sys\_OperatorModel* Operator model

**i\_IndoorOutdoor&i\_FormVal&i\_AppMeth&i\_AppEquip&**

**1**

RPE reduction factor	
key_MixRPE, ay_MixRPE	
None	1
FP1, P1 and similar	0,25
FP2, P2 and similar	0,1

PPE reduction factor	
key_MixPPEBody, ay_MixPPEBody	
Potential exposure	1
Work wear - arms, body and legs covered	0,1
Certified protective coverall	0,05

PPE reduction factor	
key_MixPPEHead, ay_MixPPEHead	
None	1
Hood	0,5
Hood and visor	0,05
FP1, P1 and similar	0,8
FP2, P2 and similar	0,8

Application: Gloves PPE reduction factor (depending on formulation type)	
key_AppPPEHands, ay_AppPPEHands	
Wettable powder, soluble powder	0,05
Chemical resistant gloves	0,05
Granules, fine granules	0,05
Chemical resistant gloves	0,05
Wettable granules, soluble granules	0,1
Chemical resistant gloves	0,1
Soluble concentrates, emulsifiable concentrate, etc.	1
Chemical resistant gloves	1
Wettable powder, soluble powder	1
Granules, fine granules	1
None	1
Wettable granules, soluble granules	1
None	1
Soluble concentrates, emulsifiable concentrate, etc.	1
None	1

Crop dependent exposure parameters										
key_CropType, ay_CropType	Transfer coefficients	Transfer coefficients	1400	Transfer coefficients	Area Treated					
Crop type	Arm, body and legs covered	Total potential exposure	Activity	hours per day	Body parts involved	Hands, arm, body and legs covered	Type of crop for Resident Bystander	Vehicle Mounted	Applications	
Bare soil	NA	NA	NA	NA	NA	NA	Field crops			50
Low berries and other small fruits		3000	5800 Reaching, picking		8 Hand and forearm		750 Field crops			50
Brassica vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops			50
Bulb vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops			50
Cane fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Field crops			10
Cereals		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops			50
Citrus fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops			10
Fruiting vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops			50
Grapes		10100	30000 Hand harvesting		8 Hand and body	no TC available for this assessment	Grapes			10
Grassland and lawns		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops			50
Golf course, turf or other sports lawns		2500	5800 Maintenance		8 Hand and body		580 Field crops			50
Hops		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Hops			10
Leaf vegetables and fresh herbs		2500	5800 Reaching, picking		8 Hand and body		580 Field crops			50
Legume vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops			50
Oilfruits		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops			10
Oilseeds		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops			50
Ornamentals		5000	14000 Cutting, sorting, bundling, carrying		8 Hand and body		1400 Field crops			10
Pome fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops			10
Root and tuber vegetables		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops			50
Stone fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops			10
Tree nuts		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops			10

Resident Spray Drift					
These values are the 75th Percentiles for Residents (assuming average breathing rates for inhalation exposures)					
key_ResidSpray, ay_ResidSpray	Adults Dermal	Children Dermal	Adults Inhalation	Children Inhalation	
Downward spraying2-3		0,47	0,327	0,0001	0,00022
Downward spraying5		0,24	0,22	0,00009	0,00017
Downward spraying10		0,20	0,18	0,00009	0,00013
Upward spraying2-3	NA	NA	NA	NA	
Upward spraying5		5,63	1,689	0,0021	0,00164
Upward spraying10		5,63	1,689	0,0021	0,00164

Bystander Spray Drift					
These values are the 95th Percentiles for Bystanders (assuming high breathing rates for inhalation exposures)					
key_BySpray, ay_BySpray	Adults Dermal	Children Dermal	Adults Inhalation	Children Inhalation	
Downward spraying2-3		1,21	0,74	0,0005	0,0011
Downward spraying5		0,57	0,48	0,00048	0,0008
Downward spraying10		0,48	0,39	0,00051	0,00076
Upward spraying2-3	NA	NA	NA	NA	
Upward spraying5		12,9	3,87	0,0044	0,0035
Upward spraying10		12,9	3,87	0,0044	0,0035

Mean Spray Drift					
These values are the mean values (assuming average breathing rates for inhalation exposures)					
key_AvgSpray, ay_AvgSpray	Adults Dermal	Children Dermal	Adults Inhalation	Children Inhalation	
Downward spraying2-3		0,22	0,18	0,0001	0,0002
Downward spraying5		0,12	0,12	0,0001	0,0001
Downward spraying10		0,11	0,1	0,0001	0,0001
Upward spraying2-3	NA	NA	NA	NA	
Upward spraying5		3,68	1,11	0,0017	0,0013
Upward spraying10		3,68	1,11	0,0017	0,0013

Resident and bystander Surface Deposits Drift percentage			
Ground sediments in % of the application rate calculated on the basis of percentile values (drift data acc. Rautmann)			
key_ByCropType, ay_ByCropType	Bystander surface deposit (90th Percentile)	Resident surface deposit (77th Percentile)	mean
Field cropsnot relevant2-3		0,085	0,056
Field cropsnot relevant5		0,035	0,023
Field cropsnot relevant10		0,019	0,013
Fruit cropsnot relevant2-3		0,292	0,240
Fruit cropsnot relevant5		0,199	0,158
Fruit cropsnot relevant10		0,118	0,090
Fruit cropsearly (without leaves)2-3		0,292	0,240
Fruit cropsearly (without leaves)5		0,199	0,158

Fruit cropsearly (without leaves)10	0,118	0,090	0,061
Fruit cropslate (dense foliage)2-3	0,157	0,110	0,070
Fruit cropslate (dense foliage)5	0,084	0,060	0,037
Fruit cropslate (dense foliage)10	0,036	0,027	0,016
Grapesnot relevant2-3	0,080	0,069	0,053
Grapesnot relevant5	0,036	0,031	0,023
Grapesnot relevant10	0,012	0,010	0,008
Hopsnot relevant2-3	0,193	0,159	0,100
Hopsnot relevant5	0,116	0,086	0,059
Hopsnot relevant10	0,058	0,037	0,029

1	2	3	4	5	6	7	8	9	10	11	12	13	14
<b>Mitch Mit-oid</b>	<b>Outdoor/Indoor</b>	<b>Formulation type</b>	<b>Application method</b>	<b>Application equipment</b>	<b>Type of exposure</b>	<b>Mixing &amp; Loading 75th percentile</b>	<b>Mixing &amp; Loading 95th percentile</b>	<b>Mixing &amp; Loading Comments</b>	<b>Mixing &amp; Loading Model</b>	<b>Application 75th percentile</b>	<b>Application 95th percentile</b>	<b>Application Comments</b>	<b>Application Model</b>
IndoorGranules, fine granulesApplication of granulesManualBody	Indoor	Granules, fine granules	Application of granules	Manual	Body			Value for application is for combination of mixing&loading and application	PHED	68,8708	253,4438	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
IndoorGranules, fine granulesApplication of granulesManualHands	Indoor	Granules, fine granules	Application of granules	Manual	Hands			Value for application is for combination of mixing&loading and application	PHED	26,5320	94,3636	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
IndoorGranules, fine granulesApplication of granulesManualInhalation	Indoor	Granules, fine granules	Application of granules	Manual	Inhalation			Value for application is for combination of mixing&loading and application	PHED	0,4677	1,5251	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesBroadcast application of granulesVehicle-mountedBody	Outdoor	Granules, fine granules	Broadcast application of granules	Vehicle-mounted	Body	0,0162	0,0427	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0,0047	0,0151	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesBroadcast application of granulesVehicle-mountedHands	Outdoor	Granules, fine granules	Broadcast application of granules	Vehicle-mounted	Hands	0,0015	0,0069	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0,0004	0,0018	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesBroadcast application of granulesVehicle-mountedInhalation	Outdoor	Granules, fine granules	Broadcast application of granules	Vehicle-mounted	Inhalation	0,0208	0,0784		PHED	0,0012	0,0045		PHED
OutdoorGranules, fine granulesIn furrow application of granulesVehicle-mountedBody	Outdoor	Granules, fine granules	In furrow application of granules	Vehicle-mounted	Body	0,0162	0,0427	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0,0047	0,0151	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesIn furrow application of granulesVehicle-mountedHands	Outdoor	Granules, fine granules	In furrow application of granules	Vehicle-mounted	Hands	0,0015	0,0069	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0,0004	0,0018	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesIn furrow application of granulesVehicle-mountedInhalation	Outdoor	Granules, fine granules	In furrow application of granules	Vehicle-mounted	Inhalation	0,0208	0,0784		PHED	0,0012	0,0045		PHED
OutdoorGranules, fine granulesManual application of granulesManualBody	Outdoor	Granules, fine granules	Manual application of granules	Manual	Body			Value for application is for combination of mixing&loading and application	PHED	68,8708	253,4438	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesManual application of granulesManualHands	Outdoor	Granules, fine granules	Manual application of granules	Manual	Hands			Value for application is for combination of mixing&loading and application	PHED	26,5320	94,3636	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesManual application of granulesManualInhalation	Outdoor	Granules, fine granules	Manual application of granules	Manual	Inhalation			Value for application is for combination of mixing&loading and application	PHED	0,4677	1,5251	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED





