

## **Instructions for using the workbook**

The information required for the exposure assessment needs to be entered in the worksheet "**Data entry**".

In the following worksheets formulas calculate the exposure values automatically

Worksheet "**Operator Outdoor Spray AOEM**" is to be for outdoor spray applications. PPE options can be selected in this worksheet

Worksheet "**Operator Granules**" is for granular applications. Currently the calculator does not allow operator exposure for indoor applications. PPE options can be selected in this worksheet

Worksheets "**Resident exposure**" and "**Bystander exposure**" are only relevant for outdoor applications

Worksheet "**Recreational Exposure**" is only applicable for golf course, turf, other sports lawns or amenity turf/grassland areas where members of the public are likely to have access

The combined results of the exposure assessment are presented in worksheet "**Summary**"

This calculator should be used in conjunction with the **Guidance on the Assessment of Exposure for Operators, Workers, Residents and Bystanders in Risk Assessment for Plant Protection Products**

Latest version: 23 Oct 2014 - Version produced to support guidance document published 23/10/2014

Note: Some drop-down menus depend on others. To avoid errors, please fill-in from top to bottom

<b>Substance name</b>	Kresoxim-methyl
<b>Product name</b>	Collis
<b>Reference value non acutely toxic active substance (RVNAS)</b>	0,9 mg/kg bw/day
<b>Reference value acutely toxic active substance (RVAAS)</b>	0,9 mg/kg bw/day
<b>Crop type</b>	Grapes
<b>Substance properties</b>	
Formulation type	Soluble concentrates, emulsifiable concentrate, etc.
Minimum volume water for application (liquids)	200 L/ha
Maximum application rate of active substance	0,064 kg a.s. /ha
50% Dissipation Time DT50	30 days
Initial Dislodgeable Foliar Residue	3 µg/cm <sup>2</sup> of foliage/kg a.s. applied/ha
Dermal absorption of product	100,00%
Dermal absorption of in-use dilution	100,00%
Oral absorption of active substance	63,00%
Inhalation absorption of active substance	100,00%
Vapour pressure of active substance	moderately volatile substances with a vapour pressure between 5*10 <sup>-3</sup> Pa and 10 <sup>-2</sup> Pa
<b>Scenario</b>	
Indoor or Outdoor application	Outdoor
Application method	Upward spraying
Application equipment	Manual-Hand held
Buffer strip	5 m
Number of applications	3
Interval between multiple applications	7 days
Season (upward spraying orchards only)	not relevant

**Exposure assessment**

Substance	Kresoxim-methyl	Formulation = Soluble concentrates, emulsifiable concentrate, etc.	Application rate-0,064 kg a.s. /ha	Spray dilution = 0,32 g a.s./l	Vapour pressure = moderately volatile substances with a vapour pressure between 5*10-3Pa and 10-2Pa
Scenario	Grapes / Outdoor / Upward spraying / Manual-Hand held			Buffer = 5	Number applications = 3, Application interval = 7 days
Percentage Absorption	Dermal for product = 100	Dermal for in use dilution = 100	Oral = 63	Inhalation = 100	
RVNAS	0,9 mg/kg bw/day		RVAAS	0,9 mg/kg bw/day	
DFR	3 µg a.s./cm2 per kg a.s./ha		DT50	30 days	

<b>Operator Model</b>	Mixing, loading and application AOEM				
Potential exposure	Longer term systemic exposure mg/kg bw/day		0,8956	% of RVNAS	99,52%
	Acute systemic exposure mg/kg bw/day		3,9155	% of RVAAS	435,06%
Mixing and Loading	Gloves = No		Clothing = Work wear - arms, body and legs covered	RPE = None	Soluble bags = No
Application	Gloves = No		Clothing = Work wear - arms, body and legs covered	RPE = None	Closed cabin = No
Exposure (including PPE options above)	Longer term systemic exposure mg/kg bw/day		0,0625	% of RVNAS	6,95%
	Acute systemic exposure mg/kg bw/day		0,1991	% of RVAAS	22,12%

<b>Worker - Hand harvesting</b>	Potential exposure mg/kg bw/day		1,9771	% of RVNAS	219,67%
	Working clothing mg/kg bw/day		0,6656	% of RVNAS	73,96%
	Working clothing and gloves mg/kg bw/day			% of RVNAS	

<b>Resident - child</b>	Spray drift (75th percentile) mg/kg bw/day		#NA	% of RVNAS	#NA
	Vapour (75th percentile) mg/kg bw/day		#NA	% of RVNAS	#NA
	Surface deposits (75th percentile) mg/kg bw/day		#NA	% of RVNAS	#NA
	Entry into treated crops (75th percentile) mg/kg bw/day		#NA	% of RVNAS	#NA
	All pathways (mean) mg/kg bw/day		#NA	% of RVNAS	#NA

<b>Resident - adult</b>	Spray drift (75th percentile) mg/kg bw/day		#NA	% of RVNAS	#NA
	Vapour (75th percentile) mg/kg bw/day		#NA	% of RVNAS	#NA
	Surface deposits (75th percentile) mg/kg bw/day		#NA	% of RVNAS	#NA
	Entry into treated crops (75th percentile) mg/kg bw/day		#NA	% of RVNAS	#NA
	All pathways (mean) mg/kg bw/day		#NA	% of RVNAS	#NA

<b>Bystander - child</b>	Spray drift (95th percentile) mg/kg bw/day		#NA	% of RVAAS	#NA
	Vapour (95th percentile) mg/kg bw/day		#NA	% of RVAAS	#NA
	Surface deposits (95th percentile) mg/kg bw/day		#NA	% of RVAAS	#NA
	Entry into treated crops (95th percentile) mg/kg bw/day		#NA	% of RVAAS	#NA
<b>Bystander - adult</b>	Spray drift (95th percentile) mg/kg bw/day		#NA	% of RVAAS	#NA

### Exposure assessment

Vapour (95th percentile) mg/kg bw/day	#NA	% of RVAAS	#NA
Surface deposits (95th percentile) mg/kg bw/day	#NA	% of RVAAS	#NA
Entry into treated crops (95th percentile) mg/kg bw/day	#NA	% of RVAAS	#NA

<b>Recreational Exposure</b>	Child % of RVNAS	Adult % of RVNAS
------------------------------	------------------	------------------

Latest version: 23 Oct 2014 - Version produced to support guidance document published 23/10/2014

**Operator exposure for Collis outdoor spray applications**

Application rate of active substance	0,064 kg a.s./ha	<i>i_AppRate</i>
Assumed area treated	4 ha/day	<i>d_AreaTreated</i>
Amount of active substance applied	0,256 kg a.s./day	<i>i_AmountAS</i>
Dermal absorption of the product	100,00%	<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	100,00%	<i>i_AbsorInuse</i>
Formulation type	Soluble concentrates, emulsifiable concentrate, etc.	
Indoor or Outdoor application	Outdoor	
Application method	Upward spraying	
Application equipment	Manual-Hand held	
Season	not relevant	

	Exposure values	µg exposure/day mixed and loaded		Reference	Comment
		75 <sup>th</sup> centile	95 <sup>th</sup> centile		
Mixing and loading	Hands	1701	6160	AOEM	
	Body	1369	48480	AOEM	
	Head	13	73	AOEM	
	Protected hands (gloves)	14	51	AOEM	
	Protected body (workwear or protective garment and sturdy footwear)	7	37	AOEM	
	Protected head (hood and face shield)	0	4	AOEM	
	Inhalation	2	28	AOEM	
	<b>Protective Equipment</b>	Select for inclusion		Penetration factor	Inhalation Protection factor
	Gloves	No			
	Clothing	Work wear - arms, body and legs covered		Incl. in AOEM model	
Head and respiratory PPE	None		1	1	
Water soluble bag	No		1		
	Exposure values	µg exposure/day applied		Reference	Comment
		75 <sup>th</sup> centile	95 <sup>th</sup> centile		
Application	Hands	871	3103	AOEM	No data available for a drift reduction scenario
	Body	49658	176482	AOEM	
	Head	98	526	AOEM	
	Protected hands (gloves)	6	32	AOEM	
	Protected body (workwear or protective garment and sturdy footwear)	1033	1938	AOEM	
	Inhalation	26	81	AOEM	
	<b>Protective Equipment</b>	Select for inclusion		Penetration factor	Inhalation Protection factor
	Gloves	No			
	Clothing	Work wear - arms, body and legs covered		Incl. in AOEM model	
	Head and respiratory PPE	None		1	1
Closed cab	No		vehicle mounted upward spraying only		

1. Total			
	Without RPE/PPE	With RPE/PPE	
Longer term			
Total systemic exposure from mixing, loading and application (mg a.s./day)	53,7387828	3,7522529	
Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	0,8956464	0,0625375	
% of RVNAS	99,52%	6,95%	
Acute			

Total systemic exposure from mixing, loading and application (mg a.s./day)	234,9327357	11,9461763	
--	-------------	------------	--

Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	3,9155456	0,1991029	
% of RVAAS	435,06%	22,12%	

## 2. Longer term exposure

### 2.1 Mixing and loading

	Systemic exposure [ $\mu\text{g a.s. /day}$ ]	Systemic exposure [ $\mu\text{g a.s./kg bw/day}$ ]	Formula
<b>Without RPE/PPE</b>			
Hands	1701,3958631	28,3565977	$D15^*i\_AbsorpProduct$
Body	1368,8760302	22,8146005	$D16^*i\_AbsorpProduct$
Head	13,2821985	0,2213700	$D17^*i\_AbsorpProduct$
Inhalation	2,4676992	0,0411283	$D21^*i\_AbsorpInhalation$
Sum	3086,0217909	51,4336965	
<b>With RPE/PPE (as selected above)</b>			
Hands	1701,3958631	28,3565977	$D18^*i\_AbsorpProduct$
Body	7,1034881	0,1183915	$D19^*i\_AbsorpProduct$ or $D15^*i\_AbsorpProduct^*F24$
Head	13,2821985	0,2213700	$D20^*i\_AbsorpProduct$ or $D17^*i\_AbsorpProduct^*F25$
Inhalation	2,4676992	0,0411283	$D21^*i\_AbsorpInhalation^*G25$
Sum	1724,2492488	28,7374875	
Water soluble bag	1724,2492488	28,7374875	$C70^*F26$

### 2.2 Application

	Systemic exposure [ $\mu\text{g a.s. /day}$ ]	Systemic exposure [ $\mu\text{g a.s./kg bw/day}$ ]	Formula
<b>Without RPE/PPE</b>			
Hands	870,6203933	14,5103399	$D30^*i\_AbsorpInuse$
Body	49657,8279411	827,6304657	$D31^*i\_AbsorpInuse$
Head	98,3108799	1,6385147	$D32^*i\_AbsorpInuse$
Inhalation	26,0018308	0,4333638	$D35^*i\_AbsorpInhalation$
Sum	50652,7610452	844,2126841	
<b>With RPE/PPE (as selected above)</b>			
Hands	870,6203933	14,5103399	$D33^*i\_AbsorpInuse$
Body	1033,0705948	17,2178432	$D34^*i\_AbsorpInuse$ or $D31^*i\_AbsorpInuse^*F38$
Head	98,3108799	1,6385147	$D32^*i\_AbsorpInuse^*F39$
Inhalation	26,0018308	0,4333638	$D35^*i\_AbsorpInuse^*G39$
Sum	2028,0036988	33,8000616	

## 3. Acute exposure

### 3.1 Mixing and loading

	Systemic exposure [ $\mu\text{g a.s. /day}$ ]	Systemic exposure [ $\mu\text{g a.s./kg bw/day}$ ]	Formula
<b>Without RPE/PPE</b>			
Hands	6159,5662599	102,6594377	$E15^*i\_AbsorpProduct$
Body	48479,9177922	807,9986299	$E16^*i\_AbsorpProduct$
Head	72,8467138	1,2141119	$E17^*i\_AbsorpProduct$
Inhalation	27,7980464	0,4633008	$E21^*i\_AbsorpInhalation$
Sum	54740,1288124	912,3354802	
<b>With RPE/PPE (as selected above)</b>			
Hands	6159,5662599	102,6594377	$E18^*i\_AbsorpProduct$
Body	37,4403556	0,6240059	$E19^*i\_AbsorpProduct$ or $E16^*i\_AbsorpProduct^*F24$
Head	72,8467138	1,2141119	$E20^*i\_AbsorpProduct$ or $E17^*i\_AbsorpProduct^*F25$
Inhalation	27,7980464	0,4633008	$E21^*i\_AbsorpInhalation^*G25$
Sum	6297,6513758	104,9608563	
Water soluble bag	6297,6513758	104,9608563	$C104^*F26$

### 2.2 Application

	Systemic exposure [ $\mu\text{g a.s. /day}$ ]	Systemic exposure [ $\mu\text{g a.s./kg bw/day}$ ]	Formula
<b>Without RPE/PPE</b>			
Hands	3102,9282850	51,7154714	$E30^*i\_AbsorpInuse$
Body	176482,2882208	2941,3714703	$E31^*i\_AbsorpInuse$
Head	526,4062331	8,7734372	$E32^*i\_AbsorpInuse$
Inhalation	80,9841462	1,3497358	$E35^*i\_AbsorpInhalation$

Sum	180192,6068851	3003,2101148	
With RPE/PPE (as selected above)			
Hands	3102,9282850	51,7154714	<i>E33*i_Absorpnuse</i>
Body	1938,2062961	32,3034383	<i>E34*i_Absorpnuse or E31*i_Absorpnuse*F38</i>
Head	526,4062331	8,7734372	<i>E32*i_Absorpnuse*F39</i>
Inhalation	80,9841462	1,3497358	<i>E35*i_Absorpnhalation*G39</i>
Sum	5648,5249604	94,1420827	



### Operator exposure for Collis granular applications

Application rate of active substance	0,064 kg a.s./ha	<i>i_AppRate</i>
Assumed area treated	4 ha/day	<i>d_AreaTreated</i>
Amount of active substance applied	0,256 kg a.s./day	<i>i_AmountAS</i>
Dermal absorption of the product	100,00%	<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	100,00%	<i>i_AbsorInuse</i>
Formulation type	Soluble concentrates, emulsifiable concentrate, etc.	
Indoor or Outdoor application	Outdoor <b>This sheet is only to be used for granular applications</b>	
Application method	Upward spraying	
Application equipment	Manual-Hand held	

	Exposure values	mg exposure/kg a.s. mixed and loaded		Reference	Comment
		75 <sup>th</sup> centile	95 <sup>th</sup> centile		
Mixing and loading	Hands	#N/A	#N/A	#N/A	#N/A
	Body	#N/A	#N/A	#N/A	#N/A
	Inhalation	#N/A	#N/A	#N/A	#N/A
	<b>Protective Equipment</b>	Choose item		Penetration factor	
	Gloves	Chemical resistant gloves			Protection for granules exposure is based on measured values
	Body PPE	Certified protective coverall			
	RPE	None		1	

	Exposure values	mg exposure/kg a.s. applied		Reference	Comment
		75 <sup>th</sup> centile	95 <sup>th</sup> centile		
Application	Hands	#N/A	#N/A	#N/A	#N/A
	Body	#N/A	#N/A	#N/A	#N/A
	Inhalation	#N/A	#N/A	#N/A	#N/A
	<b>Protective Equipment</b>	Choose item		Penetration factor	
	Gloves	Chemical resistant gloves			Protection for granules exposure is based on measured values
	Body PPE	Certified protective coverall			
	RPE	FP1, P1 and similar		0,25	

#### 1. Total

	Without RPE/PPE	With RPE/PPE
Longer term		
Total systemic exposure from mixing, loading and application (mg a.s./day)	#N/A	#N/A
Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	#N/A	#N/A
% of RVNAS	#N/A	#N/A

Acute			
Total systemic exposure from mixing, loading and application (mg a.s./day)	#N/A	#N/A	

Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	#N/A	#N/A
% of RVAAS	#N/A	#N/A

## 2. Longer term exposure

### 2.1 Mixing and loading

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
<b>Without RPE/PPE</b>			
Hands	#N/A	#N/A	$D14*100*i\_AmountAS*i\_AbsorpProduct$
Body	#N/A	#N/A	$D15*100*i\_AmountAS*i\_AbsorpProduct$
Inhalation	#N/A	#N/A	$D16*i\_AmountAS*i\_AbsorpInhalation$
Sum	#N/A	#N/A	
<b>With RPE/PPE (as selected above)</b>			
Hands	#N/A	#N/A	$D14*i\_AmountAS*i\_AbsorpProduct$
Body	#N/A	#N/A	$D15*i\_AmountAS*i\_AbsorpProduct$
Inhalation	#N/A	#N/A	$D16*i\_AmountAS*i\_AbsorpInhalation*F20$
Sum	#N/A	#N/A	

### 2.2 Application

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
<b>Without RPE/PPE</b>			
Hands	#N/A	#N/A	$D25*100*i\_AmountAS*i\_AbsorpInuse$
Body	#N/A	#N/A	$D26*100*i\_AmountAS*i\_AbsorpInuse$
Inhalation	#N/A	#N/A	$D27*i\_AmountAS*i\_AbsorpInhalation$
Sum	#N/A	#N/A	
<b>With RPE/PPE (as selected above)</b>			
Hands	#N/A	#N/A	$D25*i\_AmountAS*i\_AbsorpInuse$
Body	#N/A	#N/A	$D26*i\_AmountAS*i\_AbsorpInuse$
Inhalation	#N/A	#N/A	$D27*i\_AmountAS*i\_AbsorpInhalation*F31$
Sum	#N/A	#N/A	

## 3. Acute exposure

### 3.1 Mixing and loading

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
<b>Without RPE/PPE</b>			
Hands	#N/A	#N/A	$E14*100*i\_AmountAS*i\_AbsorpProduct$
Body	#N/A	#N/A	$E15*100*i\_AmountAS*i\_AbsorpProduct$
Inhalation	#N/A	#N/A	$E16*i\_AmountAS*i\_AbsorpInhalation$
Sum	#N/A	#N/A	
<b>With RPE/PPE (as selected above)</b>			
Hands	#N/A	#N/A	$E14*100*i\_AmountAS*i\_AbsorpProduct$
Body	#N/A	#N/A	$E15*100*i\_AmountAS*i\_AbsorpProduct$
Inhalation	#N/A	#N/A	$E16*i\_AmountAS*i\_AbsorpInhalation*F20$
Sum	#N/A	#N/A	

### 3.2 Application

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
<b>Without RPE/PPE</b>			
Hands	#N/A	#N/A	$E25*100*i\_AmountAS*i\_AbsorpInuse$
Body	#N/A	#N/A	$E25*100*i\_AmountAS*i\_AbsorpInuse$

Inhalation	#N/A	#N/A	$E26 * i\_AmountAS * i\_Absorplnhalation$
Sum	#N/A	#N/A	
<b>With RPE/PPE (as selected above)</b>			
Hands	#N/A	#N/A	$E25 * 100 * i\_AmountAS * i\_Absorplnuse$
Body	#N/A	#N/A	$E26 * 100 * i\_AmountAS * i\_Absorplnuse$
Inhalation	#N/A	#N/A	$E27 * i\_AmountAS * i\_Absorplnhalation * F31$
Sum	#N/A	#N/A	

## Worker exposure from residues on foliage for Collis

Crop type	Grapes	
Indoor or outdoor	Outdoor	
Application method	Upward spraying	
Application equipment	Manual-Hand held	
Worker's task	Hand harvesting	
Main body parts in contact with foliage	Hand and body	
Application rate of active substance	0,064 kg a.s./ha	<i>i_AppRate</i>
Number of applications	3	<i>i_AppNo</i>
Interval between multiple applications	7 days	<i>i_AppInt</i>
Half-life of active substance	30 days	<i>d_HalfLifeAS</i>
Multiple application factor	2,6	<i>d_MAF</i>
Dermal absorption of the product	100,00%	<i>i_AbsorpProduct</i>
Dermal absorption of the in-use dilution	100,00%	<i>i_Absorplnuse</i>
Dislodgeable foliar residue ( <i>i_AppRate</i> * <i>i_DFR</i> )	0,192 µg a.s./cm <sup>2</sup>	<i>d_DFR</i>
Working hours	8 hr	<i>d_WorkHr</i>
Dermal transfer coefficient - Total potential exposure	30000 cm <sup>2</sup> /hr	<i>d_DermTcUCV</i>
Dermal transfer coefficient - arms, body and legs covered	10100 cm <sup>2</sup> /hr	<i>d_DermTcCV1</i>
Dermal transfer coefficient - hands, arms, body and legs covered	no TC available for this assessment cm <sup>2</sup> /hr	<i>d_DermTcCV2</i>
Inhalation transfer coefficient for automated applications	NA ha/hr*10 <sup>-3</sup>	<i>d_InhalTcAut</i>
Inhalation transfer coefficient for cutting ornamentals	NA ha/hr*10 <sup>-3</sup>	<i>d_InhalTcCut</i>
Inhalation transfer coefficient for sorting / bundling ornamentals	NA ha/hr*10 <sup>-3</sup>	<i>d_InhalTcSort</i>

### 1. Total

	Potential exposure	Work wear - arms, body and legs covered	Working wear and gloves	Comments
Total systemic exposure (mg a.s./day)	118,6238260	39,9366881	no TC available for this assessment	
Total systemic exposure per kg body weight (mg/kg bw/day)	1,9770638	0,6656115		
% of RVNAS	219,67%	73,96%		

### 2. Details

	Systemic exposure		Formula	Comments
	[mg a.s. /day]	[mg a.s./kg bw/day]		
Dermal - Potential	118,6238260	1,9770638	$d\_DermTcUCV * d\_WorkHr * i\_DFR * i\_MAF / 1000 * i\_Absorplnuse$	
Dermal - Work wear - arms, body and legs covered	39,9366881	0,6656115	$d\_DermTcCV1 * d\_WorkHr * d\_DFR * d\_MAF / 1000 * i\_Absorplnuse$	
Dermal - Working wear and gloves	no TC available for this assessment		$d\_DermTcCV2 * d\_WorkHr * d\_DFR * d\_MAF / 1000 * i\_Absorplnuse$	
Inhalation				Na for outdoor activities

## Resident exposure for Collis

Croptype	Grapes	
Application method	Upward spraying	
Application equipment	Manual-Hand held	<i>i_AppEquip</i>
Formulation type	Soluble concentrates, emulsifiable concentrate, etc.	<i>i_FormVal</i>
Buffer strip	5 m	<i>i_Buffer</i>
Application rate of the product	0,064 kg a.s./ha	<i>i_AppRate</i>
Concentration of active substance (in-use dilution for liquid applications)	0,32 g a.s./l	<i>d_ConcAS</i>
Dermal absorption of product	100,00%	<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	100,00%	<i>i_AbsorpInuse</i>
Oral absorption	63,00%	<i>i_AbsorpOrallnuse</i>
Dislodgeable foliar residue ( $i\_AppRate * i\_DFR$ )	0,192 µg a.s./cm <sup>2</sup>	<i>d_DFR</i>
Vapour pressure of in-use dilution	moderately volatile substances with a vapour pressure between 5*10-3Pa and 10-2Pa	<i>i_Volat</i>
Concentration in air	0,015 mg/m <sup>3</sup>	<i>d_AirCon</i>
Resident dermal spray drift exposure 75th percentile - adult	5,63 ml spray dilution/person	
Resident dermal spray drift exposure 75th percentile - child	1,689 ml spray dilution/person	
Resident inhal. spray drift exposure 75th percentile - adult	0,00210 ml spray dilution/person	
Resident inhal. spray drift exposure 75th percentile - child	0,00164 ml spray dilution/person	
Resident dermal spray drift exposure mean - adult	3,68 ml spray dilution/person	
Resident dermal spray drift exposure mean - child	1,11 ml spray dilution/person	
Resident inhal. spray drift exposure mean - adult	0,00170 ml spray dilution/person	
Resident inhal. spray drift exposure mean - child	0,00133 ml spray dilution/person	
Exposure duration dermal	2 hours	<i>d_ReExpDur</i>
Exposure duration inhalation	24 hours	<i>d_ReExpDurInhal</i>
Exposure duration entry into treated crops	0,25 hours	<i>d_ExpDurTreatCrop</i>
Light clothing adjustment factor	18,0%	<i>d_ClothAF</i>
Breathing rate adult	0,23 m <sup>3</sup> /day/kg	<i>d_BreathRAAd</i>
Breathing rate child (1-3 year old)	1,07 m <sup>3</sup> /day/kg	<i>d_BreathRCh</i>
Drift percentage on surface (75th percentile)	3,07%	
Drift percentage on surface (mean)	2,32%	
Turf transferable residues percentage	5,00%	<i>d_Turf</i>
Transfer coeff. of surface deposits-adult	7300 cm <sup>2</sup> /hour	<i>d_ReTCAd</i>
Transfer coeff. of surface deposits-child (1-3 year old)	2600 cm <sup>2</sup> /hour	<i>d_ReTCCh</i>
Saliva extraction percentage	50,00%	<i>d_SalExt</i>
Surface area of hands mouthed	20 cm <sup>2</sup>	<i>d_AreaHM</i>
Frequency of hand to mouth activity	9,5 events/hour	<i>d_ReFreqHM</i>
Ingestion rate for mouthing of grass per day	25 cm <sup>2</sup>	<i>d_MouthGrass</i>
Dislodgeable residues percentage transferability for object to mouth	20,00%	<i>d_DRP</i>
Transfer coefficient for entry into treated crops (75th percentile) - adult	7500 cm <sup>2</sup> /h	<i>d_TcEntryAd</i>
Transfer coefficient for entry into treated crops (75th percentile) - child	2250 cm <sup>2</sup> /h	<i>d_TcEntryCh</i>
Transfer coefficient for entry into treated crops (mean) - adult	5980 cm <sup>2</sup> /h	<i>d_TcEntryAd</i>
Transfer coefficient for entry into treated crops (mean) - child	1794 cm <sup>2</sup> /h	<i>d_TcEntryCh</i>

1. Total

1.1 1-3 year old child

	Spray drift (75th percentile)	Vapour (75th percentile)	Surface deposits (75th percentile)	Entry into treated crops (75th percentile)	All pathways (mean)
Total systemic exposure (mg a.s./day)	0,4437195	0,1605000	0,0136128	0,2780246	0,6837295
Total systemic exposure per kg body weight (mg/kg bw/day)	0,0443720	0,0160500	0,0013613	0,0278025	0,0683729
% of RVNAS	4,93%	1,78%	0,15%	3,09%	7,60%

1.2 Adult

	Spray drift	Vapour	Surface deposits	Entry into treated crops	All pathways (mean)
Total systemic exposure (mg a.s./day)	1,4779840	0,2070000	0,0369233	0,9267486	1,9400065
Total systemic exposure per kg body weight (mg/kg bw/day)	0,0246331	0,0034500	0,0006154	0,0154458	0,0323334
% of RVNAS	2,74%	0,38%	0,07%	1,72%	3,59%

2. Resident exposure 75th Percentile

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
<b>1-3 year old child</b>				
Spray drift	0,4437195	0,0443720	$((C16 * i\_Absorpnuse * (1 - d\_ClothAF)) + C18) * d\_ConcAS$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	0,1605000	0,0160500	$d\_AirCon * d\_BreathRCh * d\_BwChild$	
Surface deposits				
Dermal	0,0131508	0,0013151	$(i\_AppRate / 100) * C29 * d\_Turf * d\_ReTCCh * d\_ReExpDur * MAX(i\_AbsorpProduct, i\_Absorpnuse) * d\_MAF * IF(i\_AppEquip = "Vehicle-mounted-Drift Reduction", 0.5, 1)$	
Hand to mouth	0,0003027	0,0000303	$(i\_AppRate / 100) * C29 * d\_Turf * d\_SalExt * d\_AreaHM * d\_ReFreqHM * d\_ReExpDur * i\_AbsorpOralnuse * d\_MAF$	
Object to mouth	0,0001593	0,0000159	$(i\_AppRate / 100) * C29 * d\_DRP * d\_MouthGrass * i\_AbsorpOralnuse * d\_MAF$	

Entry into treated crops				
Dermal	0,2780246	0,0278025	$(d\_TcEntryCh*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_Absorplnuse)$	
Hand to mouth			$(i\_AppRate/100)*d\_Turf*d\_MAF*d\_SalExt*d\_AreaHM*d\_ReFreqHM*d\_ReExpDur*i\_AbsorpOrallnuse$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Object to mouth			$(i\_AppRate/100)*d\_DRP*d\_MouthGrass*i\_AbsorpOrallnuse*d\_MAF$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
<b>Adult</b>				
Spray drift	1,4779840	0,0246331	$(C15*i\_Absorplnuse*(1-d\_ClothAF))+C17)*d\_ConcAS$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	0,2070000	0,0034500	$d\_AirCon*d\_BreathRAD*d\_BwAdult$	
Surface deposits (dermal)	0,0369233	0,0006154	$(i\_AppRate/100)*C30*d\_Turf*d\_ReTCA*d\_ReExpDur*i\_AbsorpProduct*d\_MAF$	
Entry into treated crops (dermal)	0,9267486	0,0154458	$(d\_TcEntryAd*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_Absorplnuse)$	

### 3. Summing of exposure pathways mean

	Systemic exposure [mg a.s./day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
<b>1-3 year old child</b>				
Spray drift	0,2912640	0,0291264	$((C20*i\_Absorplnuse*(1-d\_ClothAF))+C22)*d\_ConcAS$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	0,1605000	0,0160500	$d\_AirCon*d\_BreathRCh*d\_BwChild$	
Surface deposits				
Dermal	0,0099380	0,0009938	$(i\_AppRate/100)*C30*d\_Turf*d\_ReTCh*d\_ReExpDur*MAX(i\_AbsorpProduct,i\_Absorplnuse)*d\_MAF*IF(i\_AppEquip = "Vehicle-mounted-Drift Reduction",0.5,1)$	
Hand to mouth	0,0002288	0,0000229	$(i\_AppRate/100)*C30*d\_Turf*d\_SalExt*d\_AreaHM*d\_ReFreqHM*d\_ReExpDur*i\_AbsorpOrallnuse*d\_MAF$	
Object to mouth	0,0001204	0,0000120	$(i\_AppRate/100)*C30*d\_DRP*d\_MouthGrass*i\_AbsorpOrallnuse*d\_MAF$	
Entry into treated crops				
Dermal	0,2216783	0,0221678	$(d\_TcEntryMeanCh*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_Absorplnuse)$	
Hand to mouth			$(i\_AppRate/100)*1*d\_Turf*d\_MAF*d\_SalExt*d\_AreaHM*d\_ReFreqHM*d\_ReExpDur*i\_AbsorpOrallnuse$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Object to mouth			$(i\_AppRate/100)*1*d\_DRP*d\_MouthGrass*i\_AbsorpOrallnuse*d\_MAF$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
<b>Adult</b>				
Spray drift	0,9661760	0,0161029	$"(C19*i\_Absorplnuse*(1-d\_ClothAF))+C21)*d\_ConcAS"$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	0,2070000	0,0034500	$d\_AirCon*d\_BreathRAD*d\_BwAdult$	
Surface deposits (dermal)	0,0279030	0,0004650	$(i\_AppRate/100)*C30*d\_Turf*d\_ReTCA*d\_ReExpDur*MAX(i\_AbsorpProduct,i\_Absorplnuse)*d\_MAF*IF(i\_AppEquip = "Vehicle-mounted-Drift Reduction",0.5,1)$	
Entry into treated crops (dermal)	0,7389276	0,0123155	$(d\_TcEntryMeanAd*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_Absorplnuse)$	



## Bystander exposure for Collis

Croptype	Grapes	
Application method	Upward spraying	
Application equipment	Manual-Hand held	<i>i_AppEquip</i>
Formulation type	Soluble concentrates, emulsifiable concentrate, etc.	
Application rate of the product	0,064 kg a.s./ha	<i>i_AppRate</i>
Buffer strip	5 m	<i>i_Buffer</i>
Concentration of active substance (in-use dilution for liquid applications)	0,32 g a.s./l	<i>d_ConcAS</i>
Dermal absorption of product	100,00%	<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	100,00%	<i>i_AbsorpInuse</i>
Oral absorption	63,00%	<i>i_AbsorpOrallnuse</i>
Dislodgeable foliar residue ( <i>i_AppRate</i> * <i>i_DFR</i> )	0,192 µg a.s./cm <sup>2</sup>	<i>d_DFR</i>
Vapour pressure of in-use dilution	moderately volatile substances with a vapour pressure between 5*10-3Pa and 10-2Pa Pa	<i>i_Volat</i>
Concentration in air	0,015 mg/m <sup>3</sup>	<i>d_AirCon</i>
Bystander dermal spray drift exposure - adult	12,9 ml spray dilution/person	
Bystander dermal spray drift exposure - child	3,87 ml spray dilution/person	
Bystander inhal. spray drift exposure - adult	0,00440 ml spray dilution/person	
Bystander inhal. spray drift exposure - child	0,00348 ml spray dilution/person	
Exposure duration	2 hours	<i>d_ByExpDur</i>
Exposure duration entry into treated crops	0,25 hours	<i>d_ExpDurTreatCrop</i>
Light clothing adjustment factor	18,0%	<i>d_ClothAF</i>
Breathing rate adult	0,23 m <sup>3</sup> /hours/kg	<i>d_BreathRAd</i>
Breathing rate child (1-3 year old)	1,07 m <sup>3</sup> /hours/kg	<i>d_BreathRCh</i>
Drift percentage on surface (90th percentile)	3,62%	
Turf transferable residues percentage	5,00%	<i>d_Turf</i>
Transfer coeff. of surface deposits-adult	14500 cm <sup>2</sup> /hour	<i>d_ByTCAd</i>
Transfer coeff. of surface deposits-child (1-3 year old)	5200 cm <sup>2</sup> /hour	<i>d_ByTCCh</i>
Saliva extraction percentage	50,00%	<i>d_SalExt</i>
Surface area of hands mouthed	20 cm <sup>2</sup>	<i>d_AreaHM</i>
Frequency of hand to mouth activity	20 events/hour	<i>d_ByFreqHM</i>
Ingestion rate for mouthing of grass per day	25 cm <sup>2</sup>	<i>d_MouthGrass</i>
Dislodgeable residues percentage transferability for object to mouth	20,00%	<i>d_DRP</i>
Transfer coefficient for entry into treated crops - adult	7500 cm <sup>2</sup> /h	<i>d_TcEntryAd</i>
Transfer coefficient for entry into treated crops - child	2250 cm <sup>2</sup> /h	<i>d_TcEntryCh</i>

### 1. Total

#### 1.1 1-3 year old child

	Spray drift	Vapour	Surface deposits	Entry into treated crops
Total systemic exposure (mg a.s./day)	1,0166027	0,1605000	0,0319529	0,2780246
Total systemic exposure per kg body weight (mg/kg bw/day)	0,1016603	0,0160500	0,0031953	0,0278025
% of RVAAS	11,30%	1,78%	0,36%	3,09%

#### 1.2 Adult

	Spray drift	Vapour	Surface deposits	Entry into treated crops
Total systemic exposure (mg a.s./day)	3,3863680	0,2070000	0,0864801	0,9267486
Total systemic exposure per kg body weight (mg/kg bw/day)	0,0564395	0,0034500	0,0014413	0,0154458
% of RVAAS	6,27%	0,38%	0,16%	1,72%

2. Details

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
<b>1-3 year old child</b>				
Spray drift	<b>1,0166027</b>	<b>0,1016603</b>	$((C16*i\_Absorplnuse*(1-d\_ClothAF))+C18)*d\_ConcAS$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	<b>0,1605000</b>	<b>0,0160500</b>	$d\_AirCon*d\_BreathRCh*d\_BwChild$	
Surface deposits				
Dermal	<b>0,0310135</b>	<b>0,0031014</b>	$(i\_AppRate/100)*C24*d\_Turf*d\_ByTCCh*d\_ByExpDur*MAX(i\_AbsorpProduct,i\_Absorplnuse)*d\_MAF*IF(i\_AppEquip="Vehicle-mounted-Drift Reduction",0.5,1)$	
Hand to mouth	<b>0,0007515</b>	<b>0,0000751</b>	$(i\_AppRate/100)*C25*d\_Turf*d\_SalExt*d\_AreaHM*d\_ByFreqHM*d\_ByExpDur*i\_AbsorpOrallnuse*d\_MAF$	
Object to mouth	<b>0,0001879</b>	<b>0,0000188</b>	$(i\_AppRate/100)*C25*d\_DRP*d\_MouthGrass*i\_AbsorpOrallnuse*d\_MAF$	

Entry into treated crops				
Dermal	<b>0,2780246</b>	<b>0,0278025</b>	$(d\_TcEntryCh*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_Absorplnuse)$	
Hand to mouth			$(i\_AppRate/100)*d\_MAF*d\_Turf*d\_SalExt*d\_AreaHM*d\_ByFreqHM*d\_ByExpDur*i\_AbsorpOrallnuse$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Object to mouth			$(i\_AppRate/100)*d\_DRP*d\_MouthGrass*i\_AbsorpOrallnuse*d\_MAF$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
<b>Adult</b>				
Spray drift	<b>3,3863680</b>	<b>0,0564395</b>	$((C15*i\_Absorplnuse*(1-d\_ClothAF))+C17)*d\_ConcAS$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	<b>0,2070000</b>	<b>0,0034500</b>	$d\_AirCon*d\_BreathRAD*d\_BwAdult$	
Surface deposits (dermal)	<b>0,0864801</b>	<b>0,0014413</b>	$(i\_AppRate/100)*C24*d\_Turf*d\_ByTCAd*d\_ByExpDur*MAX(i\_AbsorpProduct,i\_Absorplnuse)*d\_MAF*IF(i\_AppEquip="Vehicle-mounted-Drift Reduction",0.5,1)$	
Entry into treated crops (dermal)	<b>0,9267486</b>	<b>0,0154458</b>	$(d\_TcEntryAd*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_Absorplnuse)$	

## Recreational exposure for Collis

Golf course, turf or other sports lawns		This sheet is only to be used for treatment of grassland used for recreational purposes	
Croptype			
Application method	Upward spraying		
Application equipment	Manual-Hand held		<i>i_AppEquip</i>
Formulation type	Soluble concentrates, emulsifiable concentrate, etc.		<i>i_FormVal</i>
Application rate of the product	0,064 kg a.s./ha		<i>i_AppRate</i>
Dermal absorption of product	100,00%		<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	100,00%		<i>i_Absorplnuse</i>
Oral absorption	63,00%		<i>i_AbsorpOrallnuse</i>
Dislodgeable foliar residue ( $i\_AppRate \cdot i\_DFR$ )	0,192 $\mu\text{g a.s./cm}^2$		<i>d_DFR</i>
Exposure duration dermal	2 hours		<i>d_ReExpDur</i>
Light clothing adjustment factor Adult resident	18,0%		<i>d_ClothAF</i>
Drift percentage on surface	100,00%		
Turf transferable residues percentage	5,00%		<i>d_Turf</i>
Transfer coeff. of surface deposits-adult	7300 $\text{cm}^2/\text{hour}$		<i>d_ReTCAd</i>
Transfer coeff. of surface deposits-child (1-3 year old)	2600 $\text{cm}^2/\text{hour}$		<i>d_ReTCCh</i>
Saliva extraction percentage	50,00%		<i>d_SalExt</i>
Surface area of hands mouthed	20 $\text{cm}^2$		<i>d_AreaHM</i>
Frequency of hand to mouth activity	9,5 events/hour		<i>d_ReFreqHM</i>
Ingestion rate for mouthing of grass per day	25 $\text{cm}^2$		<i>d_MouthGrass</i>

### 2. Details

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
<b>1-3 year old child</b>				
Surface deposits				
Dermal	0,4283638	0,0428364	$(i\_AppRate/100) \cdot C13 \cdot d\_Turf \cdot d\_ReTCCh \cdot d\_ReExpDur \cdot \text{MAX}(i\_AbsorpProduct, i\_Absorplnuse) \cdot d\_MAF$	
Hand to mouth	0,0098606	0,0009861	$(i\_AppRate/100) \cdot C13 \cdot d\_Turf \cdot d\_SalExt \cdot d\_AreaHM \cdot d\_ReFreqHM \cdot d\_ReExpDur \cdot i\_AbsorpOrallnuse \cdot d\_MAF$	
Object to mouth	0,0051898	0,0005190	$(i\_AppRate/100) \cdot C13 \cdot d\_DRP \cdot d\_MouthGrass \cdot i\_AbsorpOrallnuse \cdot d\_MAF$	
Total systemic exposure	0,4434142	0,0443414		
% of RVNAS				
<b>Adult</b>				
Surface deposits (dermal)	1,2027138	0,0200452	$(i\_AppRate/100) \cdot C13 \cdot d\_Turf \cdot d\_ReTCAd \cdot d\_ReExpDur \cdot \text{MAX}(i\_AbsorpProduct, i\_Absorplnuse) \cdot d\_MAF$	
% of RVNAS				

<i>d_AirConVol</i>	Concentration in air of moderately volatile substances	<b>0,015</b> mg/m <sup>3</sup>
<i>d_AirConNonVol</i>	Concentration in air of low volatile substances	<b>0,001</b> mg/m <sup>3</sup>
<i>d_AreaHM</i>	Surface area of hands mouthed	<b>20</b> cm <sup>2</sup>
<i>d_AreaTreated</i>	Area treated (defined by crop type)	<b>4</b> ha
<i>d_BreathRAd</i>	Breathing rate adult residents	<b>0,23</b> m <sup>3</sup> /day/kg
<i>d_BreathRCh</i>	Breathing rate child (1-3 year old) residents	<b>1,07</b> m <sup>3</sup> /day/kg
<i>d_BwAdult</i>	Adult body weight	<b>60</b> kg
<i>d_BwChild</i>	Child body weight (1 to < 3 year olds)	<b>10</b> kg
<i>d_ByBreathRAd</i>	Breathing rate adult bystander	<b>0,04</b> m <sup>3</sup> /hours/kg
<i>d_ByBreathRCh</i>	Breathing rate child (1-3 year old) bystander	<b>0,19</b> m <sup>3</sup> /hours/kg
<i>d_ByExpDur</i>	Exposure duration intense activity breathing rates	<b>2</b> hours
<i>d_ByFreqHM</i>	Frequency of hand to mouth activity	<b>20</b> events/hour
<i>d_ByTCAd</i>	Transfer coeff. of surface deposits-adult	<b>14500</b> cm <sup>2</sup> /hour
<i>d_ByTCCh</i>	Transfer coeff. of surface deposits-child (1-3 year old)	<b>5200</b> cm <sup>2</sup> /hour
<i>d_ClothAF</i>	Light clothing adjustment factor resident and bystanders	<b>18,0%</b>
<i>d_ConcAs</i>	Concentration of active substance (in-use dilution for liquid applications)	<b>0,32</b> g a.s./l
<i>d_DFR</i>	Dislodgeable foliar residue (i_AppRate*i_DFR)	<b>0,192</b> µg a.s./cm <sup>2</sup>
<i>d_DRP</i>	Dislodgeable residues percentage transferability for object to mouth	<b>20,0%</b>
<i>d_HalfLifeAS</i>	Half-life of active substance (DT50)	<b>30</b> days
<i>d_InhalTcAut</i>	Inhalation transfer coefficient for automated applications	<b>NA</b> ha/hr*10 <sup>^(-3)</sup>
<i>d_InhalTcCut</i>	Inhalation transfer coefficient for cutting ornamentals	<b>NA</b> ha/hr*10 <sup>^(-3)</sup>
<i>d_InhalTcSort</i>	Inhalation transfer coefficient for sorting / bundling ornamentals	<b>NA</b> ha/hr*10 <sup>^(-3)</sup>
<i>d_MAF</i>	Multiple application factor	<b>2,57</b>
<i>d_MouthGrass</i>	Ingestion rate for mouthing of grass per day	<b>25</b> cm <sup>2</sup> grass/day
<i>d_ReExpDur</i>	Exposure duration resident dermal	<b>2</b> hours
<i>d_ReExpDurInhal</i>	Exposure duration resident inhalation	<b>24</b> hours
<i>d_ExpDurTreatCrop</i>	Exposure duration for resident and bystander entry into treated crops	<b>0,25</b> hours
<i>d_ReFreqHM</i>	Frequency of hand to mouth activity	<b>9,5</b> events/hour
<i>d_ReTCAd</i>	Transfer coeff. of surface deposits-adult	<b>7300</b> cm <sup>2</sup> /hour
<i>d_ReTCCh</i>	Transfer coeff. of surface deposits-child (1-3 year old)	<b>2600</b> cm <sup>2</sup> /hour
<i>d_SalExt</i>	Saliva extraction percentage	<b>50,0%</b>
<i>d_TcEntryAd</i>	Transfer coefficient for entry into treated crops 75th percentile - adult	<b>7500</b> cm <sup>2</sup> /h
<i>d_TcEntryCh</i>	Transfer coefficient for entry into treated crops 75th percentile - child	<b>2250</b> cm <sup>2</sup> /h
<i>d_TcEntryMeanAd</i>	Transfer coefficient for entry into treated crops mean - adult	<b>5980</b> cm <sup>2</sup> /h
<i>d_TcEntryMeanCh</i>	Transfer coefficient for entry into treated crops mean - child	<b>1794</b> cm <sup>2</sup> /h
<i>d_Turf</i>	Turf transferable residues percentage	<b>5,0%</b>
<i>d_PctExtrapolation</i>	For exposure value 75 percentiles above this amount linear extrapolation is performed	<b>1,5</b> kg
<i>d_head75ProtectionFactor</i>	Coefficient to estimate head protection factor 75 th Percentile	<b>1,79422</b>
<i>d_head95ProtectionFactor</i>	Coefficient to estimate head protection factor 95 Percentile	<b>1,24705</b>

*sys\_KeyOperator* Variables for operator exposure lookup key  
*sys\_OperatorModel* Operator model

**i\_IndoorOutdoor&i\_FormVal&i\_AppMeth&i\_AppEquip&**

**1**

RPE reduction factor	
key_MixRPE, ay_MixRPE	
None	1
FP1, P1 and similar	0,25
FP2, P2 and similar	0,1

PPE reduction factor	
key_MixPPEBody, ay_MixPPEBody	
Potential exposure	1
Work wear - arms, body and legs covered	0,1
Certified protective coverall	0,05

PPE reduction factor	
key_MixPPEHead, ay_MixPPEHead	
None	1
Hood	0,5
Hood and visor	0,05
FP1, P1 and similar	0,8
FP2, P2 and similar	0,8

Application: Gloves PPE reduction factor (depending on formulation type)		
key_AppPPEHands, ay_AppPPEHands		
Wettable powder, soluble powder	Chemical resistant gloves	0,05
Granules, fine granules	Chemical resistant gloves	0,05
Wettable granules, soluble granules	Chemical resistant gloves	0,05
Soluble concentrates, emulsifiable concentrate, etc.	Chemical resistant gloves	0,1
Wettable powder, soluble powder	None	1
Granules, fine granules	None	1
Wettable granules, soluble granules	None	1
Soluble concentrates, emulsifiable concentrate, etc.	None	1

Crop dependent exposure parameters									
key_CropType, ay_CropType	Transfer coefficients	Transfer coefficients	no TC available for this assessment	Transfer coefficients	Area Treated				
Crop type	Arm, body and legs covered	Total potential exposure	Activity	hours per day	Body parts involved	Hands, arm, body and legs covered	Type of crop for Resident Bystander	Vehicle Mounted Applications	
Bare soil	NA	NA	NA	NA	NA	NA	Field crops		50
Low berries and other small fruits		3000	5800 Reaching, picking		8 Hand and forearm		750 Field crops		50
Brassica vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops		50
Bulb vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops		50
Cane fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Field crops		10
Cereals		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops		50
Citrus fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops		10
Fruiting vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops		50
Grapes		10100	30000 Hand harvesting		8 Hand and body	no TC available for this assessment	Grapes		10
Grassland and lawns		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops		50
Golf course, turf or other sports lawns		2500	5800 Maintenance		8 Hand and body		580 Field crops		50
Hops		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Hops		50
Leaf vegetables and fresh herbs		2500	5800 Reaching, picking		8 Hand and body		580 Field crops		10
Legume vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops		50
Oilfruits		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops		10
Oilseeds		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops		50
Ornamentals		5000	14000 Cutting, sorting, bundling, carrying		8 Hand and body		1400 Field crops		10
Pome fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops		10
Root and tuber vegetables		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops		50
Stone fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops		10
Tree nuts		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops		10

Resident Spray Drift					
These values are the 75th Percentiles for Residents (assuming average breathing rates for inhalation exposures)					
key_ResidSpray, ay_ResidSpray	Adults Dermal	Children Dermal	Adults Inhalation	Children Inhalation	
Downward spraying2-3		0,47	0,327	0,0001	0,00022
Downward spraying5		0,24	0,22	0,00009	0,00017
Downward spraying10		0,20	0,18	0,00009	0,00013
Upward spraying2-3	NA	NA	NA	NA	
Upward spraying5		5,63	1,689	0,0021	0,00164
Upward spraying10		5,63	1,689	0,0021	0,00164

Bystander Spray Drift					
These values are the 95th Percentiles for Bystanders (assuming high breathing rates for inhalation exposures)					
key_BySpray, ay_BySpray	Adults Dermal	Children Dermal	Adults Inhalation	Children Inhalation	
Downward spraying2-3		1,21	0,74	0,0005	0,0011
Downward spraying5		0,57	0,48	0,00048	0,0008
Downward spraying10		0,48	0,39	0,00051	0,00076
Upward spraying2-3	NA	NA	NA	NA	
Upward spraying5		12,9	3,87	0,0044	0,0035
Upward spraying10		12,9	3,87	0,0044	0,0035

Mean Spray Drift					
These values are the mean values (assuming average breathing rates for inhalation exposures)					
key_AvgSpray, ay_AvgSpray	Adults Dermal	Children Dermal	Adults Inhalation	Children Inhalation	
Downward spraying2-3		0,22	0,18	0,0001	0,0002
Downward spraying5		0,12	0,12	0,0001	0,0001
Downward spraying10		0,11	0,1	0,0001	0,0001
Upward spraying2-3	NA	NA	NA	NA	
Upward spraying5		3,68	1,11	0,0017	0,0013
Upward spraying10		3,68	1,11	0,0017	0,0013

Resident and bystander Surface Deposits Drift percentage			
Ground sediments in % of the application rate calculated on the basis of percentile values (drift data acc. Rautmann)			
key_ByCropType, ay_ByCropType	Bystander surface deposit (90th Percentile)	Resident surface deposit (77th Percentile)	mean
Field cropsnot relevant2-3		0,085	0,056
Field cropsnot relevant5		0,035	0,023
Field cropsnot relevant10		0,019	0,013
Fruit cropsnot relevant2-3		0,292	0,240
Fruit cropsnot relevant5		0,199	0,158
Fruit cropsnot relevant10		0,118	0,090
Fruit cropsearly (without leaves)2-3		0,292	0,240
Fruit cropsearly (without leaves)5		0,199	0,158

Fruit cropsearly (without leaves)10	0,118	0,090	0,061
Fruit cropslate (dense foliage)2-3	0,157	0,110	0,070
Fruit cropslate (dense foliage)5	0,084	0,060	0,037
Fruit cropslate (dense foliage)10	0,036	0,027	0,016
Grapenot relevant2-3	0,080	0,069	0,053
Grapenot relevant5	0,036	0,031	0,023
Grapenot relevant10	0,012	0,010	0,008
Hopsnot relevant2-3	0,193	0,159	0,100
Hopsnot relevant5	0,116	0,086	0,059
Hopsnot relevant10	0,058	0,037	0,029



1	2	3	4	5	6	7	8	9	10	11	12	13	14
<b>Mitch Mit-oid</b>	<b>Outdoor/Indoor</b>	<b>Formulation type</b>	<b>Application method</b>	<b>Application equipment</b>	<b>Type of exposure</b>	<b>Mixing &amp; Loading 75th percentile</b>	<b>Mixing &amp; Loading 95th percentile</b>	<b>Mixing &amp; Loading Comments</b>	<b>Mixing &amp; Loading Model</b>	<b>Application 75th percentile</b>	<b>Application 95th percentile</b>	<b>Application Comments</b>	<b>Application Model</b>
IndoorGranules, fine granulesApplication of granulesManualBody	Indoor	Granules, fine granules	Application of granules	Manual	Body			Value for application is for combination of mixing&loading and application	PHED	68,8708	253,4438	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
IndoorGranules, fine granulesApplication of granulesManualHands	Indoor	Granules, fine granules	Application of granules	Manual	Hands			Value for application is for combination of mixing&loading and application	PHED	26,5320	94,3636	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
IndoorGranules, fine granulesApplication of granulesManualInhalation	Indoor	Granules, fine granules	Application of granules	Manual	Inhalation			Value for application is for combination of mixing&loading and application	PHED	0,4677	1,5251	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesBroadcast application of granulesVehicle-mountedBody	Outdoor	Granules, fine granules	Broadcast application of granules	Vehicle-mounted	Body	0,0162	0,0427	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0,0047	0,0151	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesBroadcast application of granulesVehicle-mountedHands	Outdoor	Granules, fine granules	Broadcast application of granules	Vehicle-mounted	Hands	0,0015	0,0069	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0,0004	0,0018	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesBroadcast application of granulesVehicle-mountedInhalation	Outdoor	Granules, fine granules	Broadcast application of granules	Vehicle-mounted	Inhalation	0,0208	0,0784		PHED	0,0012	0,0045		PHED
OutdoorGranules, fine granulesIn furrow application of granulesVehicle-mountedBody	Outdoor	Granules, fine granules	In furrow application of granules	Vehicle-mounted	Body	0,0162	0,0427	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0,0047	0,0151	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesIn furrow application of granulesVehicle-mountedHands	Outdoor	Granules, fine granules	In furrow application of granules	Vehicle-mounted	Hands	0,0015	0,0069	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0,0004	0,0018	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesIn furrow application of granulesVehicle-mountedInhalation	Outdoor	Granules, fine granules	In furrow application of granules	Vehicle-mounted	Inhalation	0,0208	0,0784		PHED	0,0012	0,0045		PHED
OutdoorGranules, fine granulesManual application of granulesManualBody	Outdoor	Granules, fine granules	Manual application of granules	Manual	Body			Value for application is for combination of mixing&loading and application	PHED	68,8708	253,4438	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesManual application of granulesManualHands	Outdoor	Granules, fine granules	Manual application of granules	Manual	Hands			Value for application is for combination of mixing&loading and application	PHED	26,5320	94,3636	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesManual application of granulesManualInhalation	Outdoor	Granules, fine granules	Manual application of granules	Manual	Inhalation			Value for application is for combination of mixing&loading and application	PHED	0,4677	1,5251	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED



