

Instructions for using the workbook

The information required for the exposure assessment needs to be entered in the worksheet "**Data entry**".

In the following worksheets formulas calculate the exposure values automatically

Worksheet "**Operator Outdoor Spray AOEM**" is to be for outdoor spray applications. PPE options can be selected in this worksheet

Worksheet "**Operator Granules**" is for granular applications. Currently the calculator does not allow operator exposure for indoor applications. PPE options can be selected in this worksheet

Worksheets "**Resident exposure**" and "**Bystander exposure**" are only relevant for outdoor applications

Worksheet "**Recreational Exposure**" is only applicable for golf course, turf, other sports lawns or amenity turf/grassland areas where members of the public are likely to have access

The combined results of the exposure assessment are presented in worksheet "**Summary**"

This calculator should be used in conjunction with the **Guidance on the Assessment of Exposure for Operators, Workers, Residents and Bystanders in Risk Assessment for Plant Protection Products**

Latest version: 23 Oct 2014 - Version produced to support guidance document published 23/10/2014

Note: Some drop-down menus depend on others. To avoid errors, please fill-in from top to bottom

Substance name	Captan
Product name	Malvin WG
Reference value non acutely toxic active substance (RVNAS)	0,1 mg/kg bw/day
Reference value acutely toxic active substance (RVAAS)	0,1 mg/kg bw/day
Crop type	Pome fruit
Substance properties	
Formulation type	Wettable granules, soluble granules
Minimum volume water for application (liquids)	500 L/ha
Maximum application rate of active substance	0,75 kg a.s. /ha
50% Dissipation Time DT50	30 days
Initial Dislodgeable Foliar Residue	3 µg/cm ² of foliage/kg a.s. applied/ha
Dermal absorption of product	1,00%
Dermal absorption of in-use dilution	10,00%
Oral absorption of active substance	81,00%
Inhalation absorption of active substance	100,00%
Vapour pressure of active substance	moderately volatile substances with a vapour pressure between 5*10 ⁻³ Pa and 10 ⁻² Pa
Scenario	
Indoor or Outdoor application	Outdoor
Application method	Upward spraying
Application equipment	Vehicle-mounted
Buffer strip	5 m
Number of applications	5
Interval between multiple applications	7 days
Season (upward spraying orchards only)	late (dense foliage)

Exposure assessment

Substance	Captan	Formulation = Wettable granules, soluble granules	Application rate-0,75 kg a.s. /ha	Spray dilution = 1,5 g a.s./l	Vapour pressure = moderately volatile substances with a vapour pressure between 5*10 ⁻³ Pa and 10 ⁻² Pa
Scenario	Pome fruit late (dense foliage) / Outdoor / Upward spraying / Vehicle-mounted			Buffer = 5	Number applications = 5, Application interval = 7 days
Percentage Absorption	Derma for product = 1	Derma for in use dilution = 10	Oral = 81	Inhalation = 100	
RVNAS	0,1 mg/kg bw/day		RVAAS	0,1 mg/kg bw/day	
DFR	3 µg a.s./cm ² per kg a.s./ha		DT50	30 days	

Operator Model	Mixing, loading and application AOEM			
Potential exposure	Longer term systemic exposure mg/kg bw/day	0,1561	% of RVNAS	156,07%
	Acute systemic exposure mg/kg bw/day	0,8363	% of RVAAS	836,26%
Mixing and Loading	Gloves = No	Clothing = Work wear - arms, body and legs covered	RPE = None	Soluble bags = No
Application	Gloves = No	Clothing = Work wear - arms, body and legs covered	RPE = None	Closed cabin = No
Exposure (including PPE options above)	Longer term systemic exposure mg/kg bw/day	0,0465	% of RVNAS	46,53%
	Acute systemic exposure mg/kg bw/day	0,1916	% of RVAAS	191,64%

Worker - Searching, reaching, picking	Potential exposure mg/kg bw/day	2,5066	% of RVNAS	2506,63%
	Working clothing mg/kg bw/day	0,5013	% of RVNAS	501,33%
	Working clothing and gloves mg/kg bw/day	0,2507	% of RVNAS	250,66%

Resident - child	Spray drift (75th percentile) mg/kg bw/day	0,0210	% of RVNAS	21,02%
	Vapour (75th percentile) mg/kg bw/day	0,0161	% of RVNAS	16,05%
	Surface deposits (75th percentile) mg/kg bw/day	0,0063	% of RVNAS	6,35%
	Entry into treated crops (75th percentile) mg/kg bw/day	0,0470	% of RVNAS	47,00%
	All pathways (mean) mg/kg bw/day	0,0711	% of RVNAS	71,10%
Resident - adult	Spray drift (75th percentile) mg/kg bw/day	0,0116	% of RVNAS	11,59%
	Vapour (75th percentile) mg/kg bw/day	0,0035	% of RVNAS	3,45%
	Surface deposits (75th percentile) mg/kg bw/day	0,0020	% of RVNAS	2,05%
	Entry into treated crops (75th percentile) mg/kg bw/day	0,0261	% of RVNAS	26,11%
	All pathways (mean) mg/kg bw/day	0,0331	% of RVNAS	33,12%

Bystander - child	Spray drift (95th percentile) mg/kg bw/day	0,0481	% of RVAAS	48,12%
	Vapour (95th percentile) mg/kg bw/day	0,0161	% of RVAAS	16,05%
	Surface deposits (95th percentile) mg/kg bw/day	0,0169	% of RVAAS	16,92%
	Entry into treated crops (95th percentile) mg/kg bw/day	0,0470	% of RVAAS	47,00%
Bystander - adult	Spray drift (95th percentile) mg/kg bw/day	0,0266	% of RVAAS	26,56%

Exposure assessment

Vapour (95th percentile) mg/kg bw/day	0,0035	% of RVAAS	3,45%
Surface deposits (95th percentile) mg/kg bw/day	0,0057	% of RVAAS	5,66%
Entry into treated crops (95th percentile) mg/kg bw/day	0,0261	% of RVAAS	26,11%

Recreational Exposure	Child % of RVNAS	Adult % of RVNAS
------------------------------	------------------	------------------

Latest version: 23 Oct 2014 - Version produced to support guidance document published 23/10/2014

Operator exposure for Malvin WG outdoor spray applications

Application rate of active substance	0,75 kg a.s./ha	<i>i_AppRate</i>
Assumed area treated	10 ha/day	<i>d_AreaTreated</i>
Amount of active substance applied	7,5 kg a.s./day	<i>i_AmountAS</i>
Dermal absorption of the product	1,00%	<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	10,00%	<i>i_AbsorInuse</i>
Formulation type	Wettable granules, soluble granules	
Indoor or Outdoor application	Outdoor	
Application method	Upward spraying	
Application equipment	Vehicle-mounted	
Season	late (dense foliage)	

Mixing and loading	Exposure values	µg exposure/day mixed and loaded		Reference	Comment
		75 th centile	95 th centile		
		Hands	6183		
Body	5091	28857	AOEM		
Head	49	12448	AOEM		
Protected hands (gloves)	61	236	AOEM		
Protected body (workwear or protective garment and sturdy footwear)	111	467	AOEM		
Protected head (hood and face shield)	1	705	AOEM		
Inhalation	68	272	AOEM		
Protective Equipment	Select for inclusion		Penetration factor	Inhalation Protection factor	
Gloves	No				
Clothing	Work wear - arms, body and legs covered		Incl. in AOEM model		
Head and respiratory PPE	None		1	1	
Water soluble bag	No		1		

Application	Exposure values	µg exposure/day applied		Reference	Comment
		75 th centile	95 th centile		
		Hands	15065		
Body	66088	385622	AOEM		
Head	8685	53304	AOEM		
Protected hands (gloves)	264	6896	AOEM		
Protected body (workwear or protective garment and sturdy footwear)	862	1685	AOEM		
Inhalation	199	620	AOEM		
Protective Equipment	Select for inclusion		Penetration factor	Inhalation Protection factor	
Gloves	No				
Clothing	Work wear - arms, body and legs covered		Incl. in AOEM model		
Head and respiratory PPE	None		1	1	
Closed cab	No		vehicle mounted upward spraying only		

1. Total

	Without RPE/PPE	With RPE/PPE
Longer term		
Total systemic exposure from mixing, loading and application (mg a.s./day)	9,3642412	2,7918863
Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	0,1560707	0,0465314
% of RVNAS	156,07%	46,53%

Acute			
Total systemic exposure from mixing, loading and application (mg a.s./day)	50,1757146	11,4981791	

Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	0,8362619	0,1916363	
% of RVAAS	836,26%	191,64%	

2. Longer term exposure

2.1 Mixing and loading

	Systemic exposure [$\mu\text{g a.s. /day}$]	Systemic exposure [$\mu\text{g a.s./kg bw/day}$]	Formula
Without RPE/PPE			
Hands	61,8341525	1,0305692	$D15^*i_AbsorpProduct$
Body	50,9092635	0,8484877	$D16^*i_AbsorpProduct$
Head	0,4870026	0,0081167	$D17^*i_AbsorpProduct$
Inhalation	68,0155528	1,1335925	$D21^*i_AbsorpInhalation$
Sum	181,2459713	3,0207662	
With RPE/PPE (as selected above)			
Hands	61,8341525	1,0305692	$D18^*i_AbsorpProduct$
Body	1,1092720	0,0184879	$D19^*i_AbsorpProduct$ or $D15^*i_AbsorpProduct^*F24$
Head	0,4870026	0,0081167	$D20^*i_AbsorpProduct$ or $D17^*i_AbsorpProduct^*F25$
Inhalation	68,0155528	1,1335925	$D21^*i_AbsorpInhalation^*G25$
Sum	131,4459798	2,1907663	
Water soluble bag	131,4459798	2,1907663	$C70^*F26$

2.2 Application

	Systemic exposure [$\mu\text{g a.s. /day}$]	Systemic exposure [$\mu\text{g a.s./kg bw/day}$]	Formula
Without RPE/PPE			
Hands	1506,4616369	25,1076939	$D30^*i_AbsorpInuse$
Body	6608,7795211	110,1463254	$D31^*i_AbsorpInuse$
Head	868,5028809	14,4750480	$D32^*i_AbsorpInuse$
Inhalation	199,2511667	3,3208528	$D35^*i_AbsorpInhalation$
Sum	9182,9952056	153,0499201	
With RPE/PPE (as selected above)			
Hands	1506,4616369	25,1076939	$D33^*i_AbsorpInuse$
Body	86,2246149	1,4370769	$D34^*i_AbsorpInuse$ or $D31^*i_AbsorpInuse^*F38$
Head	868,5028809	14,4750480	$D32^*i_AbsorpInuse^*F39$
Inhalation	199,2511667	3,3208528	$D35^*i_AbsorpInuse^*G39$
Sum	2660,4402994	44,3406717	

3. Acute exposure

3.1 Mixing and loading

	Systemic exposure [$\mu\text{g a.s. /day}$]	Systemic exposure [$\mu\text{g a.s./kg bw/day}$]	Formula
Without RPE/PPE			
Hands	300,1794534	5,0029909	$E15^*i_AbsorpProduct$
Body	288,5733548	4,8095559	$E16^*i_AbsorpProduct$
Head	124,4833489	2,0747225	$E17^*i_AbsorpProduct$
Inhalation	272,2857278	4,5380955	$E21^*i_AbsorpInhalation$
Sum	985,5218850	16,4253647	
With RPE/PPE (as selected above)			
Hands	300,1794534	5,0029909	$E18^*i_AbsorpProduct$
Body	4,6661776	0,0777696	$E19^*i_AbsorpProduct$ or $E16^*i_AbsorpProduct^*F24$
Head	124,4833489	2,0747225	$E20^*i_AbsorpProduct$ or $E17^*i_AbsorpProduct^*F25$
Inhalation	272,2857278	4,5380955	$E21^*i_AbsorpInhalation^*G25$
Sum	701,6147077	11,6935785	
Water soluble bag	701,6147077	11,6935785	$C104^*F26$

2.2 Application

	Systemic exposure [$\mu\text{g a.s. /day}$]	Systemic exposure [$\mu\text{g a.s./kg bw/day}$]	Formula
Without RPE/PPE			
Hands	4677,5804252	77,9596738	$E30^*i_AbsorpInuse$

Body	38562,1521102	642,7025352	E31*i_Absorplnuse
Head	5330,4179581	88,8402993	E32*i_Absorplnuse
Inhalation	620,0422212	10,3340370	E35*i_Absorplnhalation
Sum	49190,1927146	819,8365452	
With RPE/PPE (as selected above)			
Hands	4677,5804252	77,9596738	E33*i_Absorplnuse
Body	168,5238078	2,8087301	E34*i_Absorplnuse or E31*i_Absorplnuse*F38
Head	5330,4179581	88,8402993	E32*i_Absorplnuse*F39
Inhalation	620,0422212	10,3340370	E35*i_Absorplnhalation*G39
Sum	10796,5644122	179,9427402	

Operator exposure for Malvin WG granular applications

Application rate of active substance	0,75 kg a.s./ha	<i>i_AppRate</i>
Assumed area treated	10 ha/day	<i>d_AreaTreated</i>
Amount of active substance applied	7,5 kg a.s./day	<i>i_AmountAS</i>
Dermal absorption of the product	1,00%	<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	10,00%	<i>i_AbsorInuse</i>
Formulation type	Wettable granules, soluble granules	
Indoor or Outdoor application	Outdoor This sheet is only to be used for granular applications	
Application method	Upward spraying	
Application equipment	Vehicle-mounted	

	Exposure values	mg exposure/kg a.s. mixed and loaded		Reference	Comment
		75 th centile	95 th centile		
Mixing and loading	Hands	#N/A	#N/A	#N/A	#N/A
	Body	#N/A	#N/A	#N/A	#N/A
	Inhalation	#N/A	#N/A	#N/A	#N/A
	Protective Equipment	Choose item		Penetration factor	
	Gloves	Chemical resistant gloves			Protection for granules exposure is based on measured values
	Body PPE	Certified protective coverall			
	RPE	None		1	

	Exposure values	mg exposure/kg a.s. applied		Reference	Comment
		75 th centile	95 th centile		
Application	Hands	#N/A	#N/A	#N/A	#N/A
	Body	#N/A	#N/A	#N/A	#N/A
	Inhalation	#N/A	#N/A	#N/A	#N/A
	Protective Equipment	Choose item		Penetration factor	
	Gloves	Chemical resistant gloves			Protection for granules exposure is based on measured values
	Body PPE	Certified protective coverall			
	RPE	FP1, P1 and similar		0,25	

1. Total

	Without RPE/PPE	With RPE/PPE
Longer term		
Total systemic exposure from mixing, loading and application (mg a.s./day)	#N/A	#N/A
Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	#N/A	#N/A
% of RVNAS	#N/A	#N/A
Acute		

Total systemic exposure from mixing, loading and application (mg a.s./day)	#N/A	#N/A	
--	------	------	--

Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	#N/A	#N/A
% of RVAAS	#N/A	#N/A

2. Longer term exposure

2.1 Mixing and loading

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
Without RPE/PPE			
Hands	#N/A	#N/A	$D14*100*i_AmountAS*i_AbsorpProduct$
Body	#N/A	#N/A	$D15*100*i_AmountAS*i_AbsorpProduct$
Inhalation	#N/A	#N/A	$D16*i_AmountAS*i_AbsorpInhalation$
Sum	#N/A	#N/A	
With RPE/PPE (as selected above)			
Hands	#N/A	#N/A	$D14*i_AmountAS*i_AbsorpProduct$
Body	#N/A	#N/A	$D15*i_AmountAS*i_AbsorpProduct$
Inhalation	#N/A	#N/A	$D16*i_AmountAS*i_AbsorpInhalation*F20$
Sum	#N/A	#N/A	

2.2 Application

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
Without RPE/PPE			
Hands	#N/A	#N/A	$D25*100*i_AmountAS*i_AbsorpInuse$
Body	#N/A	#N/A	$D26*100*i_AmountAS*i_AbsorpInuse$
Inhalation	#N/A	#N/A	$D27*i_AmountAS*i_AbsorpInhalation$
Sum	#N/A	#N/A	
With RPE/PPE (as selected above)			
Hands	#N/A	#N/A	$D25*i_AmountAS*i_AbsorpInuse$
Body	#N/A	#N/A	$D26*i_AmountAS*i_AbsorpInuse$
Inhalation	#N/A	#N/A	$D27*i_AmountAS*i_AbsorpInhalation*F31$
Sum	#N/A	#N/A	

3. Acute exposure

3.1 Mixing and loading

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
Without RPE/PPE			
Hands	#N/A	#N/A	$E14*100*i_AmountAS*i_AbsorpProduct$
Body	#N/A	#N/A	$E15*100*i_AmountAS*i_AbsorpProduct$
Inhalation	#N/A	#N/A	$E16*i_AmountAS*i_AbsorpInhalation$
Sum	#N/A	#N/A	
With RPE/PPE (as selected above)			
Hands	#N/A	#N/A	$E14*100*i_AmountAS*i_AbsorpProduct$
Body	#N/A	#N/A	$E15*100*i_AmountAS*i_AbsorpProduct$
Inhalation	#N/A	#N/A	$E16*i_AmountAS*i_AbsorpInhalation*F20$
Sum	#N/A	#N/A	

3.2 Application

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
Without RPE/PPE			
Hands	#N/A	#N/A	$E25*100*i_AmountAS*i_AbsorpInuse$
Body	#N/A	#N/A	$E25*100*i_AmountAS*i_AbsorpInuse$
Inhalation	#N/A	#N/A	$E26*i_AmountAS*i_AbsorpInhalation$

Sum	#N/A	#N/A	
With RPE/PPE (as selected above)			
Hands	#N/A	#N/A	$E25 * 100 * i_AmoutAS * i_Absorpnuse$
Body	#N/A	#N/A	$E26 * 100 * i_AmoutAS * i_Absorpnuse$
Inhalation	#N/A	#N/A	$E27 * i_AmoutAS * i_AbsorInhalation * F31$
Sum	#N/A	#N/A	

Worker exposure from residues on foliage for Malvin WG

Crop type	Pome fruit	
Indoor or outdoor	Outdoor	
Application method	Upward spraying	
Application equipment	Vehicle-mounted	
Worker's task	Searching, reaching, picking	
Main body parts in contact with foliage	Hand and body	
Application rate of active substance	0,75 kg a.s./ha	<i>i_AppRate</i>
Number of applications	5	<i>i_AppNo</i>
Interval between multiple applications	7 days	<i>i_AppInt</i>
Half-life of active substance	30 days	<i>d_HalfLifeAS</i>
Multiple application factor	3,7	<i>d_MAF</i>
Dermal absorption of the product	1,00%	<i>i_AbsorpProduct</i>
Dermal absorption of the in-use dilution	10,00%	<i>i_AbsorpInuse</i>
Dislodgeable foliar residue ($i_AppRate * i_DFR$)	2,25 µg a.s./cm ²	<i>d_DFR</i>
Working hours	8 hr	<i>d_WorkHr</i>
Dermal transfer coefficient - Total potential exposure	22500 cm ² /hr	<i>d_DermTcUCV</i>
Dermal transfer coefficient - arms, body and legs covered	4500 cm ² /hr	<i>d_DermTcCV1</i>
Dermal transfer coefficient - hands, arms, body and legs covered	2250 cm ² /hr	<i>d_DermTcCV2</i>
Inhalation transfer coefficient for automated applications	NA ha/hr*10 ⁻³	<i>d_InhalTcAut</i>
Inhalation transfer coefficient for cutting ornamentals	NA ha/hr*10 ⁻³	<i>d_InhalTcCut</i>
Inhalation transfer coefficient for sorting / bundling ornamentals	NA ha/hr*10 ⁻³	<i>d_InhalTcSort</i>

1. Total

	Potential exposure	Work wear - arms, body and legs covered	Working wear and gloves	Comments
Total systemic exposure (mg a.s./day)	150,3976024	30,0795205	15,0397602	
Total systemic exposure per kg body weight (mg/kg bw/day)	2,5066267	0,5013253	0,2506627	
% of RVNAS	2506,63%	501,33%	250,66%	

2. Details

	Systemic exposure		Formula	Comments
	[mg a.s. /day]	[mg a.s./kg bw/day]		
Dermal - Potential	150,3976024	2,5066267	$d_DermTcUCV * d_WorkHr * i_DFR * i_MAF / 1000 * i_AbsorpInuse$	
Dermal - Work wear - arms, body and legs covered	30,0795205	0,5013253	$d_DermTcCV1 * d_WorkHr * d_DFR * d_MAF / 1000 * i_AbsorpInuse$	
Dermal - Working wear and gloves	15,0397602	0,2506627	$d_DermTcCV2 * d_WorkHr * d_DFR * d_MAF / 1000 * i_AbsorpInuse$	
Inhalation				Na for outdoor activities

Resident exposure for Malvin WG

Croptype	Pome fruit	
Application method	Upward spraying	
Application equipment	Vehicle-mounted	<i>i_AppEquip</i>
Formulation type	Wettable granules, soluble granules	<i>i_FormVal</i>
Buffer strip	5 m	<i>i_Buffer</i>
Application rate of the product	0,75 kg a.s./ha	<i>i_AppRate</i>
Concentration of active substance (in-use dilution for liquid applications)	1,5 g a.s./l	<i>d_ConcAS</i>
Dermal absorption of product	1,00%	<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	10,00%	<i>i_AbsorpInuse</i>
Oral absorption	100,00%	<i>i_AbsorpOrallnuse</i>
Dislodgeable foliar residue ($i_AppRate * i_DFR$)	2,25 µg a.s./cm ²	<i>d_DFR</i>
Vapour pressure of in-use dilution	moderately volatile substances with a vapour pressure between 5*10-3Pa and 10-2Pa	<i>i_Volat</i>
Concentration in air	0,015 mg/m ³	<i>d_AirCon</i>
Resident dermal spray drift exposure 75th percentile - adult	5,63 ml spray dilution/person	
Resident dermal spray drift exposure 75th percentile - child	1,689 ml spray dilution/person	
Resident inhal. spray drift exposure 75th percentile - adult	0,00210 ml spray dilution/person	
Resident inhal. spray drift exposure 75th percentile - child	0,00164 ml spray dilution/person	
Resident dermal spray drift exposure mean - adult	3,68 ml spray dilution/person	
Resident dermal spray drift exposure mean - child	1,11 ml spray dilution/person	
Resident inhal. spray drift exposure mean - adult	0,00170 ml spray dilution/person	
Resident inhal. spray drift exposure mean - child	0,00133 ml spray dilution/person	
Exposure duration dermal	2 hours	<i>d_ReExpDur</i>
Exposure duration inhalation	24 hours	<i>d_ReExpDurInhal</i>
Exposure duration entry into treated crops	0,25 hours	<i>d_ExpDurTreatCrop</i>
Light clothing adjustment factor	18,0%	<i>d_ClothAF</i>
Breathing rate adult	0,23 m ³ /day/kg	<i>d_BreathRAAd</i>
Breathing rate child (1-3 year old)	1,07 m ³ /day/kg	<i>d_BreathRCh</i>
Drift percentage on surface (75th percentile)	6,04%	
Drift percentage on surface (mean)	3,73%	
Turf transferable residues percentage	5,00%	<i>d_Turf</i>
Transfer coeff. of surface deposits-adult	7300 cm ² /hour	<i>d_ReTCAd</i>
Transfer coeff. of surface deposits-child (1-3 year old)	2600 cm ² /hour	<i>d_ReTCCh</i>
Saliva extraction percentage	50,00%	<i>d_SalExt</i>
Surface area of hands mouthed	20 cm ²	<i>d_AreaHM</i>
Frequency of hand to mouth activity	9,5 events/hour	<i>d_ReFreqHM</i>
Ingestion rate for mouthing of grass per day	25 cm ²	<i>d_MouthGrass</i>
Dislodgeable residues percentage transferability for object to mouth	20,00%	<i>d_DRP</i>
Transfer coefficient for entry into treated crops (75th percentile) - adult	7500 cm ² /h	<i>d_TcEntryAd</i>
Transfer coefficient for entry into treated crops (75th percentile) - child	2250 cm ² /h	<i>d_TcEntryCh</i>
Transfer coefficient for entry into treated crops (mean) - adult	5980 cm ² /h	<i>d_TcEntryAd</i>
Transfer coefficient for entry into treated crops (mean) - child	1794 cm ² /h	<i>d_TcEntryCh</i>

1. Total

1.1 1-3 year old child

	Spray drift (75th percentile)	Vapour (75th percentile)	Surface deposits (75th percentile)	Entry into treated crops (75th percentile)	All pathways (mean)
Total systemic exposure (mg a.s./day)	0,2102122	0,1605000	0,0634956	0,4699925	0,7109824
Total systemic exposure per kg body weight (mg/kg bw/day)	0,0210212	0,0160500	0,0063496	0,0469993	0,0710982
% of RVNAS	21,02%	16,05%	6,35%	47,00%	71,10%

1.2 Adult

	Spray drift	Vapour	Surface deposits	Entry into treated crops	All pathways (mean)
Total systemic exposure (mg a.s./day)	0,6956400	0,2070000	0,1228024	1,5666417	1,9871622
Total systemic exposure per kg body weight (mg/kg bw/day)	0,0115940	0,0034500	0,0020467	0,0261107	0,0331194
% of RVNAS	11,59%	3,45%	2,05%	26,11%	33,12%

2. Resident exposure 75th Percentile

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
1-3 year old child				
Spray drift	0,2102122	0,0210212	$((C16 * i_Absorpnuse * (1 - d_ClothAF)) + C18) * d_ConcAS$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	0,1605000	0,0160500	$d_AirCon * d_BreathRCh * d_BwChild$	
Surface deposits				
Dermal	0,0437379	0,0043738	$(i_AppRate / 100) * C29 * d_Turf * d_ReTCCh * d_ReExpDur * MAX(i_AbsorpProduct, i_Absorpnuse) * d_MAF * IF(i_AppEquip = "Vehicle-mounted-Drift Reduction", 0.5, 1)$	
Hand to mouth	0,0129447	0,0012945	$(i_AppRate / 100) * C29 * d_Turf * d_SalExt * d_AreaHM * d_ReFreqHM * d_ReExpDur * i_AbsorpOralnuse * d_MAF$	
Object to mouth	0,0068130	0,0006813	$(i_AppRate / 100) * C29 * d_DRP * d_MouthGrass * i_AbsorpOralnuse * d_MAF$	

Entry into treated crops				
Dermal	0,4699925	0,0469993	$(d_TcEntryCh*0.25*d_DFR*d_MAF)/1000*MAX(i_AbsorpProduct,i_Absorplnuse)$	
Hand to mouth			$(i_AppRate/100)*d_Turf*d_MAF*d_SalExt*d_AreaHM*d_ReFreqHM*d_ReExpDur*i_AbsorpOrallnuse$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Object to mouth			$(i_AppRate/100)*d_DRP*d_MouthGrass*i_AbsorpOrallnuse*d_MAF$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Adult				
Spray drift	0,6956400	0,0115940	$(C15*i_Absorplnuse*(1-d_ClothAF))+C17)*d_ConcAS$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	0,2070000	0,0034500	$d_AirCon*d_BreathRAD*d_BwAdult$	
Surface deposits (dermal)	0,1228024	0,0020467	$(i_AppRate/100)*C30*d_Turf*d_ReTCAd*d_ReExpDur*i_Absorplnuse$	
Entry into treated crops (dermal)	1,5666417	0,0261107	$(d_TcEntryAd*0.25*d_DFR*d_MAF)/1000*MAX(i_AbsorpProduct,i_Absorplnuse)$	

3. Summing of exposure pathways mean

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
1-3 year old child				
Spray drift	0,1365300	0,0136530	$((C20*i_Absorplnuse*(1-d_ClothAF))+C22)*d_ConcAS$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	0,1605000	0,0160500	$d_AirCon*d_BreathRCh*d_BwChild$	
Surface deposits				
Dermal	0,0270103	0,0027010	$(i_AppRate/100)*C30*d_Turf*d_ReTCCh*d_ReExpDur*MAX(i_AbsorpProduct,i_Absorplnuse)*d_MAF*IF(i_AppEquip = "Vehicle-mounted-Drift Reduction",0.5,1)$	
Hand to mouth	0,0079940	0,0007994	$(i_AppRate/100)*C30*d_Turf*d_SalExt*d_AreaHM*d_ReFreqHM*d_ReExpDur*i_AbsorpOrallnuse*d_MAF$	
Object to mouth	0,0042074	0,0004207	$(i_AppRate/100)*C30*d_DRP*d_MouthGrass*i_AbsorpOrallnuse*d_MAF$	
Entry into treated crops				
Dermal	0,3747407	0,0374741	$(d_TcEntryMeanCh*0.25*d_DFR*d_MAF)/1000*MAX(i_AbsorpProduct,i_Absorplnuse)$	
Hand to mouth			$(i_AppRate/100)*1*d_Turf*d_MAF*d_SalExt*d_AreaHM*d_ReFreqHM*d_ReExpDur*i_AbsorpOrallnuse$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Object to mouth			$(i_AppRate/100)*1*d_DRP*d_MouthGrass*i_AbsorpOrallnuse*d_MAF$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Adult				
Spray drift	0,4551900	0,0075865	$"(C19*i_Absorplnuse*(1-d_ClothAF))+C21)*d_ConcAS"$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	0,2070000	0,0034500	$d_AirCon*d_BreathRAD*d_BwAdult$	
Surface deposits (dermal)	0,0758366	0,0012639	$(i_AppRate/100)*C30*d_Turf*d_ReTCAd*d_ReExpDur*MAX(i_AbsorpProduct,i_Absorplnuse)*d_MAF*IF(i_AppEquip = "Vehicle-mounted-Drift Reduction",0.5,1)$	
Entry into treated crops (dermal)	1,2491356	0,0208189	$(d_TcEntryMeanAd*0.25*d_DFR*d_MAF)/1000*MAX(i_AbsorpProduct,i_Absorplnuse)$	

Bystander exposure for Malvin WG

Croptype	Pome fruit	
Application method	Upward spraying	
Application equipment	Vehicle-mounted	<i>i_AppEquip</i>
Formulation type	Wettable granules, soluble granules	
Application rate of the product	0,75 kg a.s./ha	<i>i_AppRate</i>
Buffer strip	5 m	<i>i_Buffer</i>
Concentration of active substance (in-use dilution for liquid applications)	1,5 g a.s./l	<i>d_ConcAS</i>
Dermal absorption of product	1,00%	<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	10,00%	<i>i_AbsorpInuse</i>
Oral absorption	100,00%	<i>i_AbsorpOrallnuse</i>
Dislodgeable foliar residue (i_AppRate*i_DFR)	2,25 µg a.s./cm²	<i>d_DFR</i>
Vapour pressure of in-use dilution	moderately volatile substances with a vapour pressure between 5*10-3Pa and 10-2Pa	<i>i_Volat</i>
Concentration in air	0,015 mg/m³	<i>d_AirCon</i>
Bystander dermal spray drift exposure - adult	12,9 ml spray dilution/person	
Bystander dermal spray drift exposure - child	3,87 ml spray dilution/person	
Bystander inhal. spray drift exposure - adult	0,00440 ml spray dilution/person	
Bystander inhal. spray drift exposure - child	0,00348 ml spray dilution/person	
Exposure duration	2 hours	<i>d_ByExpDur</i>
Exposure duration entry into treated crops	0,25 hours	<i>d_ExpDurTreatCrop</i>
Light clothing adjustment factor	18,0%	<i>d_ClothAF</i>
Breathing rate adult	0,23 m³/hours/kg	<i>d_BreathRAd</i>
Breathing rate child (1-3 year old)	1,07 m³/hours/kg	<i>d_BreathRCh</i>
Drift percentage on surface (90th percentile)	8,41%	
Turf transferable residues percentage	5,00%	<i>d_Turf</i>
Transfer coeff. of surface deposits-adult	14500 cm²/hour	<i>d_ByTCAd</i>
Transfer coeff. of surface deposits-child (1-3 year old)	5200 cm²/hour	<i>d_ByTCCh</i>
Saliva extraction percentage	50,00%	<i>d_SalExt</i>
Surface area of hands mouthed	20 cm²	<i>d_AreaHM</i>
Frequency of hand to mouth activity	20 events/hour	<i>d_ByFreqHM</i>
Ingestion rate for mouthing of grass per day	25 cm²	<i>d_MouthGrass</i>
Dislodgeable residues percentage transferability for object to mouth	20,00%	<i>d_DRP</i>
Transfer coefficient for entry into treated crops - adult	7500 cm²/h	<i>d_TcEntryAd</i>
Transfer coefficient for entry into treated crops - child	2250 cm²/h	<i>d_TcEntryCh</i>

1. Total

1.1 1-3 year old child

	Spray drift	Vapour	Surface deposits	Entry into treated crops
Total systemic exposure (mg a.s./day)	0,4812350	0,1605000	0,1692314	0,4699925
Total systemic exposure per kg body weight (mg/kg bw/day)	0,0481235	0,0160500	0,0169231	0,0469993
% of RVAAS	48,12%	16,05%	16,92%	47,00%

1.2 Adult

	Spray drift	Vapour	Surface deposits	Entry into treated crops
Total systemic exposure (mg a.s./day)	1,5933000	0,2070000	0,3396340	1,5666417
Total systemic exposure per kg body weight (mg/kg bw/day)	0,0265550	0,0034500	0,0056606	0,0261107
% of RVAAS	26,56%	3,45%	5,66%	26,11%

2. Details

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
1-3 year old child				
Spray drift	0,4812350	0,0481235	$((C16*i_Absorpnuse*(1-d_ClothAF))+C18)*d_ConcAS$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	0,1605000	0,0160500	$d_AirCon*d_BreathRCh*d_BwChild$	
Surface deposits				
Dermal	0,1217998	0,0121800	$(i_AppRate/100)*C24*d_Turf*d_ByTCCh*d_ByExpDur*MAX(i_AbsorpProduct,i_Absorpnuse)*d_MAF*IF(i_AppEquip="Vehicle-mounted-Drift Reduction",0.5,1)$	
Hand to mouth	0,0379453	0,0037945	$(i_AppRate/100)*C25*d_Turf*d_SalExt*d_AreaHM*d_ByFreqHM*d_ByExpDur*i_AbsorpOralnuse*d_MAF$	
Object to mouth	0,0094863	0,0009486	$(i_AppRate/100)*C25*d_DRP*d_MouthGrass*i_AbsorpOralnuse*d_MAF$	

Entry into treated crops				
Dermal	0,4699925	0,0469993	$(d_TcEntryCh*0.25*d_DFR*d_MAF)/1000*MAX(i_AbsorpProduct,i_Absorplnuse)$	
Hand to mouth			$(i_AppRate/100)*d_MAF*d_Turf*d_SalExt*d_AreaHM*d_ByFreqHM*d_ByExpDur*i_AbsorpOrallnuse$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Object to mouth			$(i_AppRate/100)*d_DRP*d_MouthGrass*i_AbsorpOrallnuse*d_MAF$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Adult				
Spray drift	1,5933000	0,0265550	$((C15*i_Absorplnuse*(1-d_ClothAF)t)+C17)*d_ConcAS$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	0,2070000	0,0034500	$d_AirCon*d_BreathRAD*d_BwAdult$	
Surface deposits (dermal)	0,3396340	0,0056606	$(i_AppRate/100)*C24*d_Turf*d_ByTCAd*d_ByExpDur*MAX(i_AbsorpProduct,i_Absorplnuse)*d_MAF*IF(i_AppEquip="Vehicle-mounted-Drift Reduction",0.5,1)$	
Entry into treated crops (dermal)	1,5666417	0,0261107	$(d_TcEntryAd*0.25*d_DFR*d_MAF)/1000*MAX(i_AbsorpProduct,i_Absorplnuse)$	

Recreational exposure for Malvin WG

Croptype		Golf course, turf or other sports lawns		This sheet is only to be used for treatment of grassland used for recreational purposes	
Application method		Upward spraying			
Application equipment		Vehicle-mounted			<i>i_AppEquip</i>
Formulation type		Wettable granules, soluble granules			<i>i_FormVal</i>
Application rate of the product		0,75	kg a.s./ha		<i>i_AppRate</i>
Dermal absorption of product		1,00%			<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution		10,00%			<i>i_AbsorpInuse</i>
Oral absorption		100,00%			<i>i_AbsorpOrallnuse</i>
Dislodgeable foliar residue (<i>i_AppRate</i> * <i>i_DFR</i>)		2,25	µg a.s./cm ²		<i>d_DFR</i>
Exposure duration dermal		2	hours		<i>d_ReExpDur</i>
Light clothing adjustment factor Adult resident		18,0%			<i>d_ClothAF</i>
Drift percentage on surface		100,00%			
Turf transferable residues percentage		5,00%			<i>d_Turf</i>
Transfer coeff. of surface deposits-adult		7300	cm ² /hour		<i>d_ReTCAd</i>
Transfer coeff. of surface deposits-child (1-3 year old)		2600	cm ² /hour		<i>d_ReTCCh</i>
Saliva extraction percentage		50,00%			<i>d_SalExt</i>
Surface area of hands mouthed		20	cm ²		<i>d_AreaHM</i>
Frequency of hand to mouth activity		9,5	events/hour		<i>d_ReFreqHM</i>
Ingestion rate for mouthing of grass per day		25	cm ²		<i>d_MouthGrass</i>

2. Details

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
1-3 year old child				
Surface deposits				
Dermal	0,7241366	0,0724137	$(i_AppRate/100)*C13*d_Turf*d_ReTCCh*d_ReExpDur*MAX(i_AbsorpProduct,i_AbsorpInuse)*d_MAF$	
Hand to mouth	0,2143166	0,0214317	$(i_AppRate/100)*C13*d_Turf*d_SalExt*d_AreaHM*d_ReFreqHM*d_ReExpDur*i_AbsorpOrallnuse*d_MAF$	
Object to mouth	0,1127982	0,0112798	$(i_AppRate/100)*C13*d_DRP*d_MouthGrass*i_AbsorpOrallnuse*d_MAF$	
Total systemic exposure	1,0512514	0,1051251		
% of RVNAS				
Adult				
Surface deposits (dermal)	2,0331528	0,0338859	$(i_AppRate/100)*C13*d_Turf*d_ReTCAd*d_ReExpDur*MAX(i_AbsorpProduct,i_AbsorpInuse)*d_MAF$	
% of RVNAS				

<i>d_AirConVol</i>	Concentration in air of moderately volatile substances	0,015 mg/m ³
<i>d_AirConNonVol</i>	Concentration in air of low volatile substances	0,001 mg/m ³
<i>d_AreaHM</i>	Surface area of hands mouthed	20 cm ²
<i>d_AreaTreated</i>	Area treated (defined by crop type)	10 ha
<i>d_BreathRAd</i>	Breathing rate adult residents	0,23 m ³ /day/kg
<i>d_BreathRCh</i>	Breathing rate child (1-3 year old) residents	1,07 m ³ /day/kg
<i>d_BwAdult</i>	Adult body weight	60 kg
<i>d_BwChild</i>	Child body weight (1 to < 3 year olds)	10 kg
<i>d_ByBreathRAd</i>	Breathing rate adult bystander	0,04 m ³ /hours/kg
<i>d_ByBreathRCh</i>	Breathing rate child (1-3 year old) bystander	0,19 m ³ /hours/kg
<i>d_ByExpDur</i>	Exposure duration intense activity breathing rates	2 hours
<i>d_ByFreqHM</i>	Frequency of hand to mouth activity	20 events/hour
<i>d_ByTCAd</i>	Transfer coeff. of surface deposits-adult	14500 cm ² /hour
<i>d_ByTCCh</i>	Transfer coeff. of surface deposits-child (1-3 year old)	5200 cm ² /hour
<i>d_ClothAF</i>	Light clothing adjustment factor resident and bystanders	18,0%
<i>d_ConcAs</i>	Concentration of active substance (in-use dilution for liquid applications)	1,5 g a.s./l
<i>d_DFR</i>	Dislodgeable foliar residue (i_AppRate*i_DFR)	2,25 µg a.s./cm ²
<i>d_DRP</i>	Dislodgeable residues percentage transferability for object to mouth	20,0%
<i>d_HalfLifeAS</i>	Half-life of active substance (DT50)	30 days
<i>d_InhalTcAut</i>	Inhalation transfer coefficient for automated applications	NA ha/hr*10 ^{^(-3)}
<i>d_InhalTcCut</i>	Inhalation transfer coefficient for cutting ornamentals	NA ha/hr*10 ^{^(-3)}
<i>d_InhalTcSort</i>	Inhalation transfer coefficient for sorting / bundling ornamentals	NA ha/hr*10 ^{^(-3)}
<i>d_MAF</i>	Multiple application factor	3,71
<i>d_MouthGrass</i>	Ingestion rate for mouthing of grass per day	25 cm ² grass/day
<i>d_ReExpDur</i>	Exposure duration resident dermal	2 hours
<i>d_ReExpDurInhal</i>	Exposure duration resident inhalation	24 hours
<i>d_ExpDurTreatCrop</i>	Exposure duration for resident and bystander entry into treated crops	0,25 hours
<i>d_ReFreqHM</i>	Frequency of hand to mouth activity	9,5 events/hour
<i>d_ReTCAd</i>	Transfer coeff. of surface deposits-adult	7300 cm ² /hour
<i>d_ReTCCh</i>	Transfer coeff. of surface deposits-child (1-3 year old)	2600 cm ² /hour
<i>d_SalExt</i>	Saliva extraction percentage	50,0%
<i>d_TcEntryAd</i>	Transfer coefficient for entry into treated crops 75th percentile - adult	7500 cm ² /h
<i>d_TcEntryCh</i>	Transfer coefficient for entry into treated crops 75th percentile - child	2250 cm ² /h
<i>d_TcEntryMeanAd</i>	Transfer coefficient for entry into treated crops mean - adult	5980 cm ² /h
<i>d_TcEntryMeanCh</i>	Transfer coefficient for entry into treated crops mean - child	1794 cm ² /h
<i>d_Turf</i>	Turf transferable residues percentage	5,0%
<i>d_PctExtrapolation</i>	For exposure value 75 percentiles above this amount linear extrapolation is performed	1,5 kg
<i>d_head75ProtectionFactor</i>	Coefficient to estimate head protection factor 75 th Percentile	1,79422
<i>d_head95ProtectionFactor</i>	Coefficient to estimate head protection factor 95 Percentile	1,24705

sys_KeyOperator Variables for operator exposure lookup key
sys_OperatorModel Operator model

i_IndoorOutdoor&i_FormVal&i_AppMeth&i_AppEquip&

1

RPE reduction factor	
key_MixRPE, ay_MixRPE	
None	1
FP1, P1 and similar	0,25
FP2, P2 and similar	0,1

PPE reduction factor	
key_MixPPEBody, ay_MixPPEBody	
Potential exposure	1
Work wear - arms, body and legs covered	0,1
Certified protective coverall	0,05

PPE reduction factor	
key_MixPPEHead, ay_MixPPEHead	
None	1
Hood	0,5
Hood and visor	0,05
FP1, P1 and similar	0,8
FP2, P2 and similar	0,8

Application: Gloves PPE reduction factor (depending on formulation type)		
key_AppPPEHands, ay_AppPPEHands		
Wettable powder, soluble powder	Chemical resistant gloves	0,05
Granules, fine granules	Chemical resistant gloves	0,05
Wettable granules, soluble granules	Chemical resistant gloves	0,05
Soluble concentrates, emulsifiable concentrate, etc.	Chemical resistant gloves	0,1
Wettable powder, soluble powder	None	1
Granules, fine granules	None	1
Wettable granules, soluble granules	None	1
Soluble concentrates, emulsifiable concentrate, etc.	None	1

Crop dependent exposure parameters										
key_CropType, ay_CropType	Transfer coefficients	Transfer coefficients	2250	Transfer coefficients	Area Treated					
Crop type	Arm, body and legs covered	Total potential exposure	Activity	hours per day	Body parts involved	Hands, arm, body and legs covered	Type of crop for Resident Bystander	Vehicle Mounted Applications		
Bare soil	NA	NA	NA	NA	NA	NA	Field crops		50	
Low berries and other small fruits		3000	5800 Reaching, picking		8 Hand and forearm		750 Field crops		50	
Brassica vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops		50	
Bulb vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops		50	
Cane fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Field crops		10	
Cereals		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops		50	
Citrus fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops		10	
Fruiting vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops		50	
Grapes		10100	30000 Hand harvesting		8 Hand and body	no TC available for this assessment	Grapes		10	
Grassland and lawns		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops		50	
Golf course, turf or other sports lawns		2500	5800 Maintenance		8 Hand and body		580 Field crops		50	
Hops		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Hops		10	
Leaf vegetables and fresh herbs		2500	5800 Reaching, picking		8 Hand and body		580 Field crops		50	
Legume vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops		50	
Oilfruits		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops		10	
Oilseeds		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops		50	
Ornamentals		5000	14000 Cutting, sorting, bundling, carrying		8 Hand and body		1400 Field crops		10	
Pome fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops		10	
Root and tuber vegetables		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops		50	
Stone fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops		10	
Tree nuts		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops		10	

Resident Spray Drift					
These values are the 75th Percentiles for Residents (assuming average breathing rates for inhalation exposures)					
key_ResidSpray, ay_ResidSpray	Adults Dermal	Children Dermal	Adults Inhalation	Children Inhalation	
Downward spraying2-3		0,47	0,327	0,0001	0,00022
Downward spraying5		0,24	0,22	0,00009	0,00017
Downward spraying10		0,20	0,18	0,00009	0,00013
Upward spraying2-3	NA	NA	NA	NA	
Upward spraying5		5,63	1,689	0,0021	0,00164
Upward spraying10		5,63	1,689	0,0021	0,00164

Bystander Spray Drift					
These values are the 95th Percentiles for Bystanders (assuming high breathing rates for inhalation exposures)					
key_BySpray, ay_BySpray	Adults Dermal	Children Dermal	Adults Inhalation	Children Inhalation	
Downward spraying2-3		1,21	0,74	0,0005	0,0011
Downward spraying5		0,57	0,48	0,00048	0,0008
Downward spraying10		0,48	0,39	0,00051	0,00076
Upward spraying2-3	NA	NA	NA	NA	
Upward spraying5		12,9	3,87	0,0044	0,0035
Upward spraying10		12,9	3,87	0,0044	0,0035

Mean Spray Drift					
These values are the mean values (assuming average breathing rates for inhalation exposures)					
key_AvgSpray, ay_AvgSpray	Adults Dermal	Children Dermal	Adults Inhalation	Children Inhalation	
Downward spraying2-3		0,22	0,18	0,0001	0,0002
Downward spraying5		0,12	0,12	0,0001	0,0001
Downward spraying10		0,11	0,1	0,0001	0,0001
Upward spraying2-3	NA	NA	NA	NA	
Upward spraying5		3,68	1,11	0,0017	0,0013
Upward spraying10		3,68	1,11	0,0017	0,0013

Resident and bystander Surface Deposits Drift percentage				
Ground sediments in % of the application rate calculated on the basis of percentile values (drift data acc. Rautmann)				
key_ByCropType, ay_ByCropType	Bystander surface deposit (90th Percentile)	Resident surface deposit (77th Percentile)	mean	
Field cropsnot relevant2-3		0,085	0,056	0,041
Field cropsnot relevant5		0,035	0,023	0,018
Field cropsnot relevant10		0,019	0,013	0,010
Fruit cropsnot relevant2-3		0,292	0,240	0,190
Fruit cropsnot relevant5		0,199	0,158	0,117
Fruit cropsnot relevant10		0,118	0,090	0,061
Fruit cropsearly (without leaves)2-3		0,292	0,240	0,190
Fruit cropsearly (without leaves)5		0,199	0,158	0,117

Fruit cropsearly (without leaves)10	0,118	0,090	0,061
Fruit cropslate (dense foliage)2-3	0,157	0,110	0,070
Fruit cropslate (dense foliage)5	0,084	0,060	0,037
Fruit cropslate (dense foliage)10	0,036	0,027	0,016
Grapesnot relevant2-3	0,080	0,069	0,053
Grapesnot relevant5	0,036	0,031	0,023
Grapesnot relevant10	0,012	0,010	0,008
Hopsnot relevant2-3	0,193	0,159	0,100
Hopsnot relevant5	0,116	0,086	0,059
Hopsnot relevant10	0,058	0,037	0,029

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Mixing Method	Outdoor/Indoor	Formulation type	Application method	Application equipment	Type of exposure	Mixing & Loading 75th percentile	Mixing & Loading 95th percentile	Mixing & Loading Comments	Mixing & Loading Model	Application 75th percentile	Application 95th percentile	Application Comments	Application Model
IndoorGranules, fine granulesApplication of granulesManualBody	Indoor	Granules, fine granules	Application of granules	Manual	Body			Value for application is for combination of mixing&loading and application	PHED	68,8708	253,4438	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
IndoorGranules, fine granulesApplication of granulesManualHands	Indoor	Granules, fine granules	Application of granules	Manual	Hands			Value for application is for combination of mixing&loading and application	PHED	26,5320	94,3636	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
IndoorGranules, fine granulesApplication of granulesManualInhalation	Indoor	Granules, fine granules	Application of granules	Manual	Inhalation			Value for application is for combination of mixing&loading and application	PHED	0,4677	1,5251	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesBroadcast application of granulesVehicle-mountedBody	Outdoor	Granules, fine granules	Broadcast application of granules	Vehicle-mounted	Body	0,0162	0,0427	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0,0047	0,0151	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesBroadcast application of granulesVehicle-mountedHands	Outdoor	Granules, fine granules	Broadcast application of granules	Vehicle-mounted	Hands	0,0015	0,0069	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0,0004	0,0018	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesBroadcast application of granulesVehicle-mountedInhalation	Outdoor	Granules, fine granules	Broadcast application of granules	Vehicle-mounted	Inhalation	0,0208	0,0784		PHED	0,0012	0,0045		PHED
OutdoorGranules, fine granulesIn furrow application of granulesVehicle-mountedBody	Outdoor	Granules, fine granules	In furrow application of granules	Vehicle-mounted	Body	0,0162	0,0427	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0,0047	0,0151	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesIn furrow application of granulesVehicle-mountedHands	Outdoor	Granules, fine granules	In furrow application of granules	Vehicle-mounted	Hands	0,0015	0,0069	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0,0004	0,0018	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesIn furrow application of granulesVehicle-mountedInhalation	Outdoor	Granules, fine granules	In furrow application of granules	Vehicle-mounted	Inhalation	0,0208	0,0784		PHED	0,0012	0,0045		PHED
OutdoorGranules, fine granulesManual application of granulesManualBody	Outdoor	Granules, fine granules	Manual application of granules	Manual	Body			Value for application is for combination of mixing&loading and application	PHED	68,8708	253,4438	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesManual application of granulesManualHands	Outdoor	Granules, fine granules	Manual application of granules	Manual	Hands			Value for application is for combination of mixing&loading and application	PHED	26,5320	94,3636	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesManual application of granulesManualInhalation	Outdoor	Granules, fine granules	Manual application of granules	Manual	Inhalation			Value for application is for combination of mixing&loading and application	PHED	0,4677	1,5251	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED

