

## **Instructions for using the workbook**

The information required for the exposure assessment needs to be entered in the worksheet "**Data entry**".

In the following worksheets formulas calculate the exposure values automatically

Worksheet "**Operator Outdoor Spray AOEM**" is to be for outdoor spray applications. PPE options can be selected in this worksheet

Worksheet "**Operator Granules**" is for granular applications. Currently the calculator does not allow operator exposure for indoor applications. PPE options can be selected in this worksheet

Worksheets "**Resident exposure**" and "**Bystander exposure**" are only relevant for outdoor applications

Worksheet "**Recreational Exposure**" is only applicable for golf course, turf, other sports lawns or amenity turf/grassland areas where members of the public are likely to have access

The combined results of the exposure assessment are presented in worksheet "**Summary**"

This calculator should be used in conjunction with the **Guidance on the Assessment of Exposure for Operators, Workers, Residents and Bystanders in Risk Assessment for Plant Protection Products**

Latest version: 23 Oct 2014 - Version produced to support guidance document published 23/10/2014

Note: Some drop-down menus depend on others. To avoid errors, please fill-in from top to bottom

<b>Substance name</b>	flonicamid
<b>Product name</b>	Teppeki
<b>Reference value non acutely toxic active substance (RVNAS)</b>	0,025 mg/kg bw/day
<b>Reference value acutely toxic active substance (RVAAS)</b>	0,025 mg/kg bw/day
<b>Crop type</b>	Stone fruit
<b>Substance properties</b>	
Formulation type	Wettable granules, soluble granules
Minimum volume water for application (liquids)	500 L/ha
Maximum application rate of active substance	0,07 kg a.s. /ha
50% Dissipation Time DT50	30 days
Initial Dislodgeable Foliar Residue	3 µg/cm2 of foliage/kg a.s. applied/ha
Dermal absorption of product	7,46%
Dermal absorption of in-use dilution	13,00%
Oral absorption of active substance	80,00%
Inhalation absorption of active substance	100,00%
Vapour pressure of active substance	low volatile substances having a vapour pressure of <math><5 \cdot 10^{-3}</math>Pa
<b>Scenario</b>	
Indoor or Outdoor application	Outdoor
Application method	Upward spraying
Application equipment	Vehicle-mounted
Buffer strip	5 m
Number of applications	3
Interval between multiple applications	21 days
Season (upward spraying orchards only)	late (dense foliage)

### Exposure assessment

Substance	flonicamid	Formulation = Wettable granules, soluble granules	Application rate-0,07 kg a.s. /ha	Spray dilution = 0,14 g a.s./l	Vapour pressure = low volatile substances having a vapour pressure of <math>5 \cdot 10^{-3}</math>Pa
Scenario	Stone fruit late (dense foliage) / Outdoor / Upward spraying / Vehicle-mounted			Buffer = 5	Number applications = 3, Application interval = 21 days
Percentage Absorption	Dermal for product = 7,46	Dermal for in use dilution = 13	Oral = 80	Inhalation = 100	
RVNAS	0,025 mg/kg bw/day		RVAAS	0,025 mg/kg bw/day	
DFR	3 µg a.s./cm <sup>2</sup> per kg a.s./ha		DT50	30 days	

<b>Operator Model</b>					
Mixing, loading and application AOEM					
Potential exposure	Longer term systemic exposure mg/kg bw/day		0,0230	% of RVNAS	91,90%
	Acute systemic exposure mg/kg bw/day		0,1288	% of RVAAS	515,28%
Mixing and Loading	Gloves = No		Clothing = Work wear - arms, body and legs covered	RPE = None	Soluble bags = No
Application	Gloves = No		Clothing = Work wear - arms, body and legs covered	RPE = None	Closed cabin = No
Exposure (including PPE options above)	Longer term systemic exposure mg/kg bw/day		0,0086	% of RVNAS	34,42%
	Acute systemic exposure mg/kg bw/day		0,0332	% of RVAAS	132,88%

<b>Worker - Searching, reaching, picking</b>	Potential exposure mg/kg bw/day		0,1633	% of RVNAS	653,40%
	Working clothing mg/kg bw/day		0,0327	% of RVNAS	130,68%
	Working clothing and gloves mg/kg bw/day		0,0163	% of RVNAS	65,34%

<b>Resident - child</b>	Spray drift (75th percentile) mg/kg bw/day		0,0025	% of RVNAS	10,17%
	Vapour (75th percentile) mg/kg bw/day		0,0011	% of RVNAS	4,28%
	Surface deposits (75th percentile) mg/kg bw/day		0,0004	% of RVNAS	1,53%
	Entry into treated crops (75th percentile) mg/kg bw/day		0,0031	% of RVNAS	12,25%
	All pathways (mean) mg/kg bw/day		0,0054	% of RVNAS	21,62%
<b>Resident - adult</b>	Spray drift (75th percentile) mg/kg bw/day		0,0014	% of RVNAS	5,62%
	Vapour (75th percentile) mg/kg bw/day		0,0002	% of RVNAS	0,92%
	Surface deposits (75th percentile) mg/kg bw/day		0,0001	% of RVNAS	0,53%
	Entry into treated crops (75th percentile) mg/kg bw/day		0,0017	% of RVNAS	6,81%
	All pathways (mean) mg/kg bw/day		0,0026	% of RVNAS	10,35%

<b>Bystander - child</b>	Spray drift (95th percentile) mg/kg bw/day		0,0058	% of RVAAS	23,30%
	Vapour (95th percentile) mg/kg bw/day		0,0011	% of RVAAS	4,28%
	Surface deposits (95th percentile) mg/kg bw/day		0,0010	% of RVAAS	4,11%

**Exposure assessment**

	Entry into treated crops (95th percentile) mg/kg bw/day	0,0031	% of RVAAS	12,25%
<b>Bystander - adult</b>	Spray drift (95th percentile) mg/kg bw/day	0,0032	% of RVAAS	12,88%
	Vapour (95th percentile) mg/kg bw/day	0,0002	% of RVAAS	0,92%
	Surface deposits (95th percentile) mg/kg bw/day	0,0004	% of RVAAS	1,48%
	Entry into treated crops (95th percentile) mg/kg bw/day	0,0017	% of RVAAS	6,81%
<b>Recreational Exposure</b>		Child % of RVNAS		Adult % of RVNAS

Latest version: 23 Oct 2014 - Version produced to support guidance document published 23/10/2014

**Operator exposure for Teppeki outdoor spray applications**

Application rate of active substance	0,07 kg a.s./ha	<i>i_AppRate</i>
Assumed area treated	10 ha/day	<i>d_AreaTreated</i>
Amount of active substance applied	0,7 kg a.s./day	<i>i_AmountAS</i>
Dermal absorption of the product	7,46%	<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	13,00%	<i>i_AbsorInuse</i>
Formulation type	Wettable granules, soluble granules	
Indoor or Outdoor application	Outdoor	
Application method	Upward spraying	
Application equipment	Vehicle-mounted	
Season	late (dense foliage)	

	Exposure values	µg exposure/day mixed and loaded		Reference	Comment
		75 <sup>th</sup> centile	95 <sup>th</sup> centile		
Mixing and loading	Hands	996	4736	AOEM	
	Body	961	14489	AOEM	
	Head	5	1162	AOEM	
	Protected hands (gloves)	13	22	AOEM	
	Protected body (workwear or protective garment and sturdy footwear)	14	44	AOEM	
	Protected head (hood and face shield)	0	66	AOEM	
	Inhalation	34	257	AOEM	
	<b>Protective Equipment</b>	Select for inclusion		Penetration factor	Inhalation Protection factor
	Gloves	No			
	Clothing	Work wear - arms, body and legs covered		Incl. in AOEM model	
Head and respiratory PPE	None		1	1	
Water soluble bag	No		1		

	Exposure values	µg exposure/day applied		Reference	Comment
		75 <sup>th</sup> centile	95 <sup>th</sup> centile		
Application	Hands	1839	4366	AOEM	No data available for a drift reduction scenario
	Body	6168	35991	AOEM	
	Head	811	4975	AOEM	
	Protected hands (gloves)	25	644	AOEM	
	Protected body (workwear or protective garment and sturdy footwear)	80	157	AOEM	
	Inhalation	52	58	AOEM	
	<b>Protective Equipment</b>	Select for inclusion		Penetration factor	Inhalation Protection factor
	Gloves	No			
	Clothing	Work wear - arms, body and legs covered		Incl. in AOEM model	
	Head and respiratory PPE	None		1	1
Closed cab	No		vehicle mounted upward spraying only		

**1. Total**

	Without RPE/PPE	With RPE/PPE
Longer term		
Total systemic exposure from mixing, loading and application (mg a.s./day)	1,3784288	0,5163327
Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	0,0229738	0,0086055
% of RVNAS	91,90%	34,42%

Acute			
Total systemic exposure from mixing, loading and application (mg a.s./day)	7,7291916	1,9931612	

Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	0,1288199	0,0332194	
% of RVAAS	515,28%	132,88%	

## 2. Longer term exposure

### 2.1 Mixing and loading

	Systemic exposure [ $\mu\text{g a.s. /day}$ ]	Systemic exposure [ $\mu\text{g a.s./kg bw/day}$ ]	Formula
<b>Without RPE/PPE</b>			
Hands	74,3140023	1,2385667	$D15^*i\_AbsorpProduct$
Body	71,7040346	1,1950672	$D16^*i\_AbsorpProduct$
Head	0,3390837	0,0056514	$D17^*i\_AbsorpProduct$
Inhalation	33,5808107	0,5596802	$D21^*i\_AbsorpInhalation$
Sum	179,9379312	2,9989655	
<b>With RPE/PPE (as selected above)</b>			
Hands	74,3140023	1,2385667	$D18^*i\_AbsorpProduct$
Body	1,0112263	0,0168538	$D19^*i\_AbsorpProduct$ or $D15^*i\_AbsorpProduct^*F24$
Head	0,3390837	0,0056514	$D20^*i\_AbsorpProduct$ or $D17^*i\_AbsorpProduct^*F25$
Inhalation	33,5808107	0,5596802	$D21^*i\_AbsorpInhalation^*G25$
Sum	109,2451230	1,8207521	
Water soluble bag	109,2451230	1,8207521	$C70^*F26$

### 2.2 Application

	Systemic exposure [ $\mu\text{g a.s. /day}$ ]	Systemic exposure [ $\mu\text{g a.s./kg bw/day}$ ]	Formula
<b>Without RPE/PPE</b>			
Hands	239,0897408	3,9848290	$D30^*i\_AbsorpInuse$
Body	801,8652486	13,3644208	$D31^*i\_AbsorpInuse$
Head	105,3783496	1,7563058	$D32^*i\_AbsorpInuse$
Inhalation	52,1575469	0,8692924	$D35^*i\_AbsorpInhalation$
Sum	1198,4908859	19,9748481	
<b>With RPE/PPE (as selected above)</b>			
Hands	239,0897408	3,9848290	$D33^*i\_AbsorpInuse$
Body	10,4619199	0,1743653	$D34^*i\_AbsorpInuse$ or $D31^*i\_AbsorpInuse^*F38$
Head	105,3783496	1,7563058	$D32^*i\_AbsorpInuse^*F39$
Inhalation	52,1575469	0,8692924	$D35^*i\_AbsorpInuse^*G39$
Sum	407,0875572	6,7847926	

## 3. Acute exposure

### 3.1 Mixing and loading

	Systemic exposure [ $\mu\text{g a.s. /day}$ ]	Systemic exposure [ $\mu\text{g a.s./kg bw/day}$ ]	Formula
<b>Without RPE/PPE</b>			
Hands	353,2876466	5,8881274	$E15^*i\_AbsorpProduct$
Body	1080,8523932	18,0142066	$E16^*i\_AbsorpProduct$
Head	86,6736064	1,4445601	$E17^*i\_AbsorpProduct$
Inhalation	257,3290373	4,2888173	$E21^*i\_AbsorpInhalation$
Sum	1778,1426835	29,6357114	
<b>With RPE/PPE (as selected above)</b>			
Hands	353,2876466	5,8881274	$E18^*i\_AbsorpProduct$
Body	3,2489039	0,0541484	$E19^*i\_AbsorpProduct$ or $E16^*i\_AbsorpProduct^*F24$
Head	86,6736064	1,4445601	$E20^*i\_AbsorpProduct$ or $E17^*i\_AbsorpProduct^*F25$
Inhalation	257,3290373	4,2888173	$E21^*i\_AbsorpInhalation^*G25$
Sum	700,5391942	11,6756532	
Water soluble bag	700,5391942	11,6756532	$C104^*F26$

### 2.2 Application

	Systemic exposure [ $\mu\text{g a.s. /day}$ ]	Systemic exposure [ $\mu\text{g a.s./kg bw/day}$ ]	Formula
<b>Without RPE/PPE</b>			
Hands	567,5464249	9,4591071	$E30^*i\_AbsorpInuse$

Body	4678,8744560	77,9812409	E31*i_Absorpnuse
Head	646,7573789	10,7792896	E32*i_Absorpnuse
Inhalation	57,8706073	0,9645101	E35*i_Absorplnhalation
Sum	5951,0488672	99,1841478	
<b>With RPE/PPE (as selected above)</b>			
Hands	567,5464249	9,4591071	E33*i_Absorpnuse
Body	20,4475553	0,3407926	E34*i_Absorpnuse or E31*i_Absorpnuse*F38
Head	646,7573789	10,7792896	E32*i_Absorpnuse*F39
Inhalation	57,8706073	0,9645101	E35*i_Absorplnhalation*G39
Sum	1292,6219665	21,5436994	



**Operator exposure for Teppeki granular applications**

Application rate of active substance	0,07 kg a.s./ha	<i>i_AppRate</i>
Assumed area treated	10 ha/day	<i>d_AreaTreated</i>
Amount of active substance applied	0,7 kg a.s./day	<i>i_AmountAS</i>
Dermal absorption of the product	7,46%	<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	13,00%	<i>i_AbsorInuse</i>
Formulation type	Wettable granules, soluble granules	
Indoor or Outdoor application	Outdoor <b>This sheet is only to be used for granular applications</b>	
Application method	Upward spraying	
Application equipment	Vehicle-mounted	

	Exposure values	mg exposure/kg a.s. mixed and loaded		Reference	Comment
		75 <sup>th</sup> centile	95 <sup>th</sup> centile		
Mixing and loading	Hands	#N/A	#N/A	#N/A	#N/A
	Body	#N/A	#N/A	#N/A	#N/A
	Inhalation	#N/A	#N/A	#N/A	#N/A
	<b>Protective Equipment</b>	Choose item		Penetration factor	
	Gloves	Chemical resistant gloves			Protection for granules exposure is based on measured values
	Body PPE	Certified protective coverall			
	RPE	None		1	

	Exposure values	mg exposure/kg a.s. applied		Reference	Comment
		75 <sup>th</sup> centile	95 <sup>th</sup> centile		
Application	Hands	#N/A	#N/A	#N/A	#N/A
	Body	#N/A	#N/A	#N/A	#N/A
	Inhalation	#N/A	#N/A	#N/A	#N/A
	<b>Protective Equipment</b>	Choose item		Penetration factor	
	Gloves	Chemical resistant gloves			Protection for granules exposure is based on measured values
	Body PPE	Certified protective coverall			
	RPE	FP1, P1 and similar		0,25	

1. Total			
	Without RPE/PPE	With RPE/PPE	
<b>Longer term</b>			
Total systemic exposure from mixing, loading and application (mg a.s./day)	#N/A	#N/A	
Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	#N/A	#N/A	
% of RVNAS	#N/A	#N/A	
<b>Acute</b>			

Total systemic exposure from mixing, loading and application (mg a.s./day)	#N/A	#N/A	
--	------	------	--

Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	#N/A	#N/A
% of RVAAS	#N/A	#N/A

## 2. Longer term exposure

### 2.1 Mixing and loading

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
<b>Without RPE/PPE</b>			
Hands	#N/A	#N/A	$D14*100*_i\_AmoutAS*_i\_AbsorpProduct$
Body	#N/A	#N/A	$D15*100*_i\_AmoutAS*_i\_AbsorpProduct$
Inhalation	#N/A	#N/A	$D16*_i\_AmoutAS*_i\_AbsorpInhalation$
Sum	#N/A	#N/A	
<b>With RPE/PPE (as selected above)</b>			
Hands	#N/A	#N/A	$D14*_i\_AmoutAS*_i\_AbsorpProduct$
Body	#N/A	#N/A	$D15*_i\_AmoutAS*_i\_AbsorpProduct$
Inhalation	#N/A	#N/A	$D16*_i\_AmoutAS*_i\_AbsorpInhalation*F20$
Sum	#N/A	#N/A	

### 2.2 Application

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
<b>Without RPE/PPE</b>			
Hands	#N/A	#N/A	$D25*100*_i\_AmoutAS*_i\_AbsorpInuse$
Body	#N/A	#N/A	$D26*100*_i\_AmoutAS*_i\_AbsorpInuse$
Inhalation	#N/A	#N/A	$D27*_i\_AmoutAS*_i\_AbsorpInhalation$
Sum	#N/A	#N/A	
<b>With RPE/PPE (as selected above)</b>			
Hands	#N/A	#N/A	$D25*_i\_AmoutAS*_i\_AbsorpInuse$
Body	#N/A	#N/A	$D26*_i\_AmoutAS*_i\_AbsorpInuse$
Inhalation	#N/A	#N/A	$D27*_i\_AmoutAS*_i\_AbsorpInhalation*F31$
Sum	#N/A	#N/A	

## 3. Acute exposure

### 3.1 Mixing and loading

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
<b>Without RPE/PPE</b>			
Hands	#N/A	#N/A	$E14*100*_i\_AmoutAS*_i\_AbsorpProduct$
Body	#N/A	#N/A	$E15*100*_i\_AmoutAS*_i\_AbsorpProduct$
Inhalation	#N/A	#N/A	$E16*_i\_AmoutAS*_i\_AbsorpInhalation$
Sum	#N/A	#N/A	
<b>With RPE/PPE (as selected above)</b>			
Hands	#N/A	#N/A	$E14*100*_i\_AmoutAS*_i\_AbsorpProduct$
Body	#N/A	#N/A	$E15*100*_i\_AmoutAS*_i\_AbsorpProduct$
Inhalation	#N/A	#N/A	$E16*_i\_AmoutAS*_i\_AbsorpInhalation*F20$
Sum	#N/A	#N/A	

### 3.2 Application

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
<b>Without RPE/PPE</b>			
Hands	#N/A	#N/A	$E25*100*_i\_AmoutAS*_i\_AbsorpInuse$
Body	#N/A	#N/A	$E25*100*_i\_AmoutAS*_i\_AbsorpInuse$

Inhalation	#N/A	#N/A	$E26 * i\_AmoutAS * i\_Absorpinhalation$
Sum	#N/A	#N/A	
<b>With RPE/PPE (as selected above)</b>			
Hands	#N/A	#N/A	$E25 * 100 * i\_AmoutAS * i\_Absorpinuse$
Body	#N/A	#N/A	$E26 * 100 * i\_AmoutAS * i\_Absorpinuse$
Inhalation	#N/A	#N/A	$E27 * i\_AmoutAS * i\_Absorpinhalation * F31$
Sum	#N/A	#N/A	

## Worker exposure from residues on foliage for Teppeki

Crop type	Stone fruit	
Indoor or outdoor	Outdoor	
Application method	Upward spraying	
Application equipment	Vehicle-mounted	
Worker's task	Searching, reaching, picking	
Main body parts in contact with foliage	Hand and body	
Application rate of active substance	0,07 kg a.s./ha	<i>i_AppRate</i>
Number of applications	3	<i>i_AppNo</i>
Interval between multiple applications	21 days	<i>i_AppInt</i>
Half-life of active substance	30 days	<i>d_HalfLifeAS</i>
Multiple application factor	2,0	<i>d_MAF</i>
Dermal absorption of the product	7,46%	<i>i_AbsorpProduct</i>
Dermal absorption of the in-use dilution	13,00%	<i>i_AbsorpInuse</i>
Dislodgeable foliar residue ( <i>i_AppRate</i> * <i>i_DFR</i> )	0,21 µg a.s./cm <sup>2</sup>	<i>d_DFR</i>
Working hours	8 hr	<i>d_WorkHr</i>
Dermal transfer coefficient - Total potential exposure	22500 cm <sup>2</sup> /hr	<i>d_DermTcUCV</i>
Dermal transfer coefficient - arms, body and legs covered	4500 cm <sup>2</sup> /hr	<i>d_DermTcCV1</i>
Dermal transfer coefficient - hands, arms, body and legs covered	2250 cm <sup>2</sup> /hr	<i>d_DermTcCV2</i>
Inhalation transfer coefficient for automated applications	NA ha/hr*10 <sup>-3</sup>	<i>d_InhalTcAut</i>
Inhalation transfer coefficient for cutting ornamentals	NA ha/hr*10 <sup>-3</sup>	<i>d_InhalTcCut</i>
Inhalation transfer coefficient for sorting / bundling ornamentals	NA ha/hr*10 <sup>-3</sup>	<i>d_InhalTcSort</i>

### 1. Total

	Potential exposure	Work wear - arms, body and legs covered	Working wear and gloves	Comments
Total systemic exposure (mg a.s./day)	9,8009796	1,9601959	0,9800980	
Total systemic exposure per kg body weight (mg/kg bw/day)	0,1633497	0,0326699	0,0163350	
% of RVNAS	653,40%	130,68%	65,34%	

### 2. Details

	Systemic exposure		Formula	Comments
	[mg a.s. /day]	[mg a.s./kg bw/day]		
Dermal - Potential	9,8009796	0,1633497	$d\_DermTcUCV * d\_WorkHr * i\_DFR * i\_MAF / 1000 * i\_AbsorpInuse$	
Dermal - Work wear - arms, body and legs covered	1,9601959	0,0326699	$d\_DermTcCV1 * d\_WorkHr * d\_DFR * d\_MAF / 1000 * i\_AbsorpInuse$	
Dermal - Working wear and gloves	0,9800980	0,0163350	$d\_DermTcCV2 * d\_WorkHr * d\_DFR * d\_MAF / 1000 * i\_AbsorpInuse$	
Inhalation				Na for outdoor activities

## Resident exposure for Tepeki

Croptype	Stone fruit	
Application method	Upward spraying	
Application equipment	Vehicle-mounted	<i>i_AppEquip</i>
Formulation type	Wettable granules, soluble granules	<i>i_FormVal</i>
Buffer strip	5 m	<i>i_Buffer</i>
Application rate of the product	0,07 kg a.s./ha	<i>i_AppRate</i>
Concentration of active substance (in-use dilution for liquid applications)	0,14 g a.s./l	<i>d_ConcAS</i>
Dermal absorption of product	7,46%	<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	13,00%	<i>i_AbsorpInuse</i>
Oral absorption	80,00%	<i>i_AbsorpOrallnuse</i>
Dislodgeable foliar residue ( $i\_AppRate * i\_DFR$ )	0,21 µg a.s./cm <sup>2</sup>	<i>d_DFR</i>
Vapour pressure of in-use dilution	low volatile substances having a vapour pressure of <5*10 <sup>-3</sup> Pa	<i>i_Volat</i>
Concentration in air	0,001 mg/m <sup>3</sup>	<i>d_AirCon</i>
Resident dermal spray drift exposure 75th percentile - adult	5,63 ml spray dilution/person	
Resident dermal spray drift exposure 75th percentile - child	1,689 ml spray dilution/person	
Resident inhal. spray drift exposure 75th percentile - adult	0,00210 ml spray dilution/person	
Resident inhal. spray drift exposure 75th percentile - child	0,00164 ml spray dilution/person	
Resident dermal spray drift exposure mean - adult	3,68 ml spray dilution/person	
Resident dermal spray drift exposure mean - child	1,11 ml spray dilution/person	
Resident inhal. spray drift exposure mean - adult	0,00170 ml spray dilution/person	
Resident inhal. spray drift exposure mean - child	0,00133 ml spray dilution/person	
Exposure duration dermal	2 hours	<i>d_ReExpDur</i>
Exposure duration inhalation	24 hours	<i>d_ReExpDurInhal</i>
Exposure duration entry into treated crops	0,25 hours	<i>d_ExpDurTreatCrop</i>
Light clothing adjustment factor	18,0%	<i>d_ClothAF</i>
Breathing rate adult	0,23 m <sup>3</sup> /day/kg	<i>d_BreathRAD</i>
Breathing rate child (1-3 year old)	1,07 m <sup>3</sup> /day/kg	<i>d_BreathRCh</i>
Drift percentage on surface (75th percentile)	6,04%	
Drift percentage on surface (mean)	3,73%	
Turf transferable residues percentage	5,00%	<i>d_Turf</i>
Transfer coeff. of surface deposits-adult	7300 cm <sup>2</sup> /hour	<i>d_ReTCAd</i>
Transfer coeff. of surface deposits-child (1-3 year old)	2600 cm <sup>2</sup> /hour	<i>d_ReTCCh</i>
Saliva extraction percentage	50,00%	<i>d_SalExt</i>
Surface area of hands mouthed	20 cm <sup>2</sup>	<i>d_AreaHM</i>
Frequency of hand to mouth activity	9,5 events/hour	<i>d_ReFreqHM</i>
Ingestion rate for mouthing of grass per day	25 cm <sup>2</sup>	<i>d_MouthGrass</i>
Dislodgeable residues percentage transferability for object to mouth	20,00%	<i>d_DRP</i>
Transfer coefficient for entry into treated crops (75th percentile) - adult	7500 cm <sup>2</sup> /h	<i>d_TcEntryAd</i>
Transfer coefficient for entry into treated crops (75th percentile) - child	2250 cm <sup>2</sup> /h	<i>d_TcEntryCh</i>
Transfer coefficient for entry into treated crops (mean) - adult	5980 cm <sup>2</sup> /h	<i>d_TcEntryAd</i>
Transfer coefficient for entry into treated crops (mean) - child	1794 cm <sup>2</sup> /h	<i>d_TcEntryCh</i>

1. Total

1.1 1-3 year old child

	Spray drift (75th percentile)	Vapour (75th percentile)	Surface deposits (75th percentile)	Entry into treated crops (75th percentile)	All pathways (mean)
Total systemic exposure (mg a.s./day)	0,0254367	0,0107000	0,0038285	0,0306281	0,0540507
Total systemic exposure per kg body weight (mg/kg bw/day)	0,0025437	0,0010700	0,0003828	0,0030628	0,0054051
% of RVNAS	10,17%	4,28%	1,53%	12,25%	21,62%

1.2 Adult

	Spray drift	Vapour	Surface deposits	Entry into treated crops	All pathways (mean)
Total systemic exposure (mg a.s./day)	0,0843161	0,0138000	0,0080027	0,1020935	0,1553030
Total systemic exposure per kg body weight (mg/kg bw/day)	0,0014053	0,0002300	0,0001334	0,0017016	0,0025884
% of RVNAS	5,62%	0,92%	0,53%	6,81%	10,35%

2. Resident exposure 75th Percentile

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
<b>1-3 year old child</b>				
Spray drift	0,0254367	0,0025437	$((C16 * i\_AbsorpInuse * (1 - d\_ClothAF)) + C18) * d\_ConcAS$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	0,0107000	0,0010700	$d\_AirCon * d\_BreathRCh * d\_BwChild$	
Surface deposits				
Dermal	0,0028503	0,0002850	$(i\_AppRate / 100) * C29 * d\_Turf * d\_ReTCCh * d\_ReExpDur * MAX(i\_AbsorpProduct, i\_AbsorpInuse) * d\_MAF * IF(i\_AppEquip = "Vehicle-mounted-Drift Reduction", 0.5, 1)$	
Hand to mouth	0,0006409	0,0000641	$(i\_AppRate / 100) * C29 * d\_Turf * d\_SalExt * d\_AreaHM * d\_ReFreqHM * d\_ReExpDur * i\_AbsorpOrallnuse * d\_MAF$	
Object to mouth	0,0003373	0,0000337	$(i\_AppRate / 100) * C29 * d\_DRP * d\_MouthGrass * i\_AbsorpOrallnuse * d\_MAF$	

Entry into treated crops				
Dermal	0,0306281	0,0030628	$(d\_TcEntryCh*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_Absorplnuse)$	
Hand to mouth			$(i\_AppRate/100)*d\_Turf*d\_MAF*d\_SalExt*d\_AreaHM*d\_ReFreqHM*d\_ReExpDur*i\_AbsorpOrallnuse$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Object to mouth			$(i\_AppRate/100)*d\_DRP*d\_MouthGrass*i\_AbsorpOrallnuse*d\_MAF$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
<b>Adult</b>				
Spray drift	0,0843161	0,0014053	$(C15*i\_Absorplnuse*(1-d\_ClothAF))+C17)*d\_ConcAS$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	0,0138000	0,0002300	$d\_AirCon*d\_BreathRAD*d\_BwAdult$	
Surface deposits (dermal)	0,0080027	0,0001334	$(i\_AppRate/100)*C30*d\_Turf*d\_ReTCA*d\_ReExpDur*i\_Absorplnuse$	
Entry into treated crops (dermal)	0,1020935	0,0017016	$(d\_TcEntryAd*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_Absorplnuse)$	

### 3. Summing of exposure pathways mean

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
<b>1-3 year old child</b>				
Spray drift	0,0165656	0,0016566	$((C20*i\_Absorplnuse*(1-d\_ClothAF))+C22)*d\_ConcAS$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	0,0107000	0,0010700	$d\_AirCon*d\_BreathRCh*d\_BwChild$	
Surface deposits				
Dermal	0,0017602	0,0001760	$(i\_AppRate/100)*C30*d\_Turf*d\_ReTCC*d\_ReExpDur*MAX(i\_AbsorpProduct,i\_Absorplnuse)*d\_MAF*IF(i\_AppEquip = "Vehicle-mounted-Drift Reduction",0.5,1)$	
Hand to mouth	0,0003958	0,0000396	$(i\_AppRate/100)*C30*d\_Turf*d\_SalExt*d\_AreaHM*d\_ReFreqHM*d\_ReExpDur*i\_AbsorpOrallnuse*d\_MAF$	
Object to mouth	0,0002083	0,0000208	$(i\_AppRate/100)*C30*d\_DRP*d\_MouthGrass*i\_AbsorpOrallnuse*d\_MAF$	
Entry into treated crops				
Dermal	0,0244208	0,0024421	$(d\_TcEntryMeanCh*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_Absorplnuse)$	
Hand to mouth			$(i\_AppRate/100)*1*d\_Turf*d\_MAF*d\_SalExt*d\_AreaHM*d\_ReFreqHM*d\_ReExpDur*i\_AbsorpOrallnuse$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Object to mouth			$(i\_AppRate/100)*1*d\_DRP*d\_MouthGrass*i\_AbsorpOrallnuse*d\_MAF$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
<b>Adult</b>				
Spray drift	0,0551583	0,0009193	$"(C19*i\_Absorplnuse*(1-d\_ClothAF))+C21)*d\_ConcAS"$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	0,0138000	0,0002300	$d\_AirCon*d\_BreathRAD*d\_BwAdult$	
Surface deposits (dermal)	0,0049421	0,0000824	$(i\_AppRate/100)*C30*d\_Turf*d\_ReTCA*d\_ReExpDur*MAX(i\_AbsorpProduct,i\_Absorplnuse)*d\_MAF*IF(i\_AppEquip = "Vehicle-mounted-Drift Reduction",0.5,1)$	
Entry into treated crops (dermal)	0,0814026	0,0013567	$(d\_TcEntryMeanAd*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_Absorplnuse)$	



## Bystander exposure for Teppeki

Croptype	Stone fruit	
Application method	Upward spraying	
Application equipment	Vehicle-mounted	<i>i_AppEquip</i>
Formulation type	Wettable granules, soluble granules	
Application rate of the product	0,07 kg a.s./ha	<i>i_AppRate</i>
Buffer strip	5 m	<i>i_Buffer</i>
Concentration of active substance (in-use dilution for liquid applications)	0,14 g a.s./l	<i>d_ConcAS</i>
Dermal absorption of product	7,46%	<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	13,00%	<i>i_AbsorpInuse</i>
Oral absorption	80,00%	<i>i_AbsorpOrallnuse</i>
Dislodgeable foliar residue ( $i\_AppRate \cdot i\_DFR$ )	0,21 $\mu\text{g a.s./cm}^2$	<i>d_DFR</i>
Vapour pressure of in-use dilution	low volatile substances having a vapour pressure of $<5 \cdot 10^{-3} \text{Pa}$	<i>i_Volat</i>
Concentration in air	0,001 $\text{mg/m}^3$	<i>d_AirCon</i>
Bystander dermal spray drift exposure - adult	12,9 ml spray dilution/person	
Bystander dermal spray drift exposure - child	3,87 ml spray dilution/person	
Bystander inhal. spray drift exposure - adult	0,00440 ml spray dilution/person	
Bystander inhal. spray drift exposure - child	0,00348 ml spray dilution/person	
Exposure duration	2 hours	<i>d_ByExpDur</i>
Exposure duration entry into treated crops	0,25 hours	<i>d_ExpDurTreatCrop</i>
Light clothing adjustment factor	18,0%	<i>d_ClothAF</i>
Breathing rate adult	0,23 $\text{m}^3/\text{hours/kg}$	<i>d_BreathRAD</i>
Breathing rate child (1-3 year old)	1,07 $\text{m}^3/\text{hours/kg}$	<i>d_BreathRCh</i>
Drift percentage on surface (90th percentile)	8,41%	
Turf transferable residues percentage	5,00%	<i>d_Turf</i>
Transfer coeff. of surface deposits-adult	14500 $\text{cm}^2/\text{hour}$	<i>d_ByTCAd</i>
Transfer coeff. of surface deposits-child (1-3 year old)	5200 $\text{cm}^2/\text{hour}$	<i>d_ByTCCh</i>
Saliva extraction percentage	50,00%	<i>d_SalExt</i>
Surface area of hands mouthed	20 $\text{cm}^2$	<i>d_AreaHM</i>
Frequency of hand to mouth activity	20 events/hour	<i>d_ByFreqHM</i>
Ingestion rate for mouthing of grass per day	25 $\text{cm}^2$	<i>d_MouthGrass</i>
Dislodgeable residues percentage transferability for object to mouth	20,00%	<i>d_DRP</i>
Transfer coefficient for entry into treated crops - adult	7500 $\text{cm}^2/\text{h}$	<i>d_TcEntryAd</i>
Transfer coefficient for entry into treated crops - child	2250 $\text{cm}^2/\text{h}$	<i>d_TcEntryCh</i>

### 1. Total

#### 1.1 1-3 year old child

	Spray drift	Vapour	Surface deposits	Entry into treated crops
Total systemic exposure (mg a.s./day)	0,0582435	0,0107000	0,0102857	0,0306281
Total systemic exposure per kg body weight (mg/kg bw/day)	0,0058244	0,0010700	0,0010286	0,0030628
% of RVAAS	23,30%	4,28%	4,11%	12,25%

#### 1.2 Adult

	Spray drift	Vapour	Surface deposits	Entry into treated crops
Total systemic exposure (mg a.s./day)	0,1931356	0,0138000	0,0221330	0,1020935

Total systemic exposure per kg body weight (mg/kg bw/day)	0,0032189	0,0002300	0,0003689	0,0017016
% of RVAAS	12,88%	0,92%	1,48%	6,81%

## 2. Details

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
<b>1-3 year old child</b>				
Spray drift	0,0582435	0,0058244	$((C16 * i\_AbsorpInuse * (1 - d\_ClothAF)) + C18) * d\_ConcAS$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	0,0107000	0,0010700	$d\_AirCon * d\_BreathRCh * d\_BwChild$	
Surface deposits				
Dermal	0,0079373	0,0007937	$(i\_AppRate / 100) * C24 * d\_Turf * d\_ByTCCh * d\_ByExpDur * MAX(i\_AbsorpProduct, i\_AbsorpInuse) * d\_MAF * IF(i\_AppEquip = "Vehicle-mounted-Drift Reduction", 0.5, 1)$	
Hand to mouth	0,0018787	0,0001879	$(i\_AppRate / 100) * C25 * d\_Turf * d\_SalExt * d\_AreaHM * d\_ByFreqHM * d\_ByExpDur * i\_AbsorpOralInuse * d\_MAF$	
Object to mouth	0,0004697	0,0000470	$(i\_AppRate / 100) * C25 * d\_DRP * d\_MouthGrass * i\_AbsorpOralInuse * d\_MAF$	

Entry into treated crops				
Dermal	0,0306281	0,0030628	$(d\_TcEntryCh*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_AbsorpInuse)$	
Hand to mouth			$(i\_AppRate/100)*d\_MAF*d\_Turf*d\_SalExt*d\_AreaHM*d\_ByFreqHM*d\_ByExpDur*i\_AbsorpOralInuse$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Object to mouth			$(i\_AppRate/100)*d\_DRP*d\_MouthGrass*i\_AbsorpOralInuse*d\_MAF$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
<b>Adult</b>				
Spray drift	0,1931356	0,0032189	$((C15*i\_AbsorpInuse*(1-d\_ClothAF)t)+C17)*d\_ConcAS$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	0,0138000	0,0002300	$d\_AirCon*d\_BreathRA*d\_BwAdult$	
Surface deposits (dermal)	0,0221330	0,0003689	$(i\_AppRate/100)*C24*d\_Turf*d\_ByTCA*d\_ByExpDur*MAX(i\_AbsorpProduct,i\_AbsorpInuse)*d\_MAF*IF(i\_AppEquip="Vehicle-mounted-Drift Reduction",0.5,1)$	
Entry into treated crops (dermal)	0,1020935	0,0017016	$(d\_TcEntryAd*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_AbsorpInuse)$	

## Recreational exposure for Teppeki

Cropping type		Golf course, turf or other sports lawns		This sheet is only to be used for treatment of grassland used for recreational purposes	
Application method		Upward spraying			
Application equipment		Vehicle-mounted			<i>i_AppEquip</i>
Formulation type		Wettable granules, soluble granules			<i>i_FormVal</i>
Application rate of the product		0,07	kg a.s./ha		<i>i_AppRate</i>
Dermal absorption of product		7,46%			<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution		13,00%			<i>i_Absorplnuse</i>
Oral absorption		80,00%			<i>i_AbsorpOrallnuse</i>
Dislodgeable foliar residue ( <i>i_AppRate</i> * <i>i_DFR</i> )		0,21	µg a.s./cm <sup>2</sup>		<i>d_DFR</i>
Exposure duration dermal		2	hours		<i>d_ReExpDur</i>
Light clothing adjustment factor Adult resident		18,0%			<i>d_ClothAF</i>
Drift percentage on surface		100,00%			
Turf transferable residues percentage		5,00%			<i>d_Turf</i>
Transfer coeff. of surface deposits-adult		7300	cm <sup>2</sup> /hour		<i>d_ReTCAd</i>
Transfer coeff. of surface deposits-child (1-3 year old)		2600	cm <sup>2</sup> /hour		<i>d_ReTCCh</i>
Saliva extraction percentage		50,00%			<i>d_SalExt</i>
Surface area of hands mouthed		20	cm <sup>2</sup>		<i>d_AreaHM</i>
Frequency of hand to mouth activity		9,5	events/hour		<i>d_ReFreqHM</i>
Ingestion rate for mouthing of grass per day		25	cm <sup>2</sup>		<i>d_MouthGrass</i>

### 2. Details

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
<b>1-3 year old child</b>				
Surface deposits				
Dermal	0,0471899	0,0047190	$(i\_AppRate/100)*C13*d\_Turf*d\_ReTCCh*d\_ReExpDur*MAX(i\_AbsorpProduct,i\_Absorplnuse)*d\_MAF$	
Hand to mouth	0,0106107	0,0010611	$(i\_AppRate/100)*C13*d\_Turf*d\_SalExt*d\_AreaHM*d\_ReFreqHM*d\_ReExpDur*i\_AbsorpOrallnuse*d\_MAF$	
Object to mouth	0,0055846	0,0005585	$(i\_AppRate/100)*C13*d\_DRP*d\_MouthGrass*i\_AbsorpOrallnuse*d\_MAF$	
Total systemic exposure	0,0633853	0,0063385		
% of RVNAS				
<b>Adult</b>				
Surface deposits (dermal)	0,1324947	0,0022082	$(i\_AppRate/100)*C13*d\_Turf*d\_ReTCAd*d\_ReExpDur*MAX(i\_AbsorpProduct,i\_Absorplnuse)*d\_MAF$	
% of RVNAS				

<i>d_AirConVol</i>	Concentration in air of moderately volatile substances	<b>0,015</b> mg/m <sup>3</sup>
<i>d_AirConNonVol</i>	Concentration in air of low volatile substances	<b>0,001</b> mg/m <sup>3</sup>
<i>d_AreaHM</i>	Surface area of hands mouthed	<b>20</b> cm <sup>2</sup>
<i>d_AreaTreated</i>	Area treated (defined by crop type)	<b>10</b> ha
<i>d_BreathRAd</i>	Breathing rate adult residents	<b>0,23</b> m <sup>3</sup> /day/kg
<i>d_BreathRCh</i>	Breathing rate child (1-3 year old) residents	<b>1,07</b> m <sup>3</sup> /day/kg
<i>d_BwAdult</i>	Adult body weight	<b>60</b> kg
<i>d_BwChild</i>	Child body weight (1 to < 3 year olds)	<b>10</b> kg
<i>d_ByBreathRAd</i>	Breathing rate adult bystander	<b>0,04</b> m <sup>3</sup> /hours/kg
<i>d_ByBreathRCh</i>	Breathing rate child (1-3 year old) bystander	<b>0,19</b> m <sup>3</sup> /hours/kg
<i>d_ByExpDur</i>	Exposure duration intense activity breathing rates	<b>2</b> hours
<i>d_ByFreqHM</i>	Frequency of hand to mouth activity	<b>20</b> events/hour
<i>d_ByTCAd</i>	Transfer coeff. of surface deposits-adult	<b>14500</b> cm <sup>2</sup> /hour
<i>d_ByTCCh</i>	Transfer coeff. of surface deposits-child (1-3 year old)	<b>5200</b> cm <sup>2</sup> /hour
<i>d_ClothAF</i>	Light clothing adjustment factor resident and bystanders	<b>18,0%</b>
<i>d_ConcAs</i>	Concentration of active substance (in-use dilution for liquid applications)	<b>0,14</b> g a.s./l
<i>d_DFR</i>	Dislodgeable foliar residue (i_AppRate*i_DFR)	<b>0,21</b> µg a.s./cm <sup>2</sup>
<i>d_DRP</i>	Dislodgeable residues percentage transferability for object to mouth	<b>20,0%</b>
<i>d_HalfLifeAS</i>	Half-life of active substance (DT50)	<b>30</b> days
<i>d_InhalTcAut</i>	Inhalation transfer coefficient for automated applications	<b>NA</b> ha/hr*10 <sup>^(-3)</sup>
<i>d_InhalTcCut</i>	Inhalation transfer coefficient for cutting ornamentals	<b>NA</b> ha/hr*10 <sup>^(-3)</sup>
<i>d_InhalTcSort</i>	Inhalation transfer coefficient for sorting / bundling ornamentals	<b>NA</b> ha/hr*10 <sup>^(-3)</sup>
<i>d_MAF</i>	Multiple application factor	<b>1,99</b>
<i>d_MouthGrass</i>	Ingestion rate for mouthing of grass per day	<b>25</b> cm <sup>2</sup> grass/day
<i>d_ReExpDur</i>	Exposure duration resident dermal	<b>2</b> hours
<i>d_ReExpDurInhal</i>	Exposure duration resident inhalation	<b>24</b> hours
<i>d_ExpDurTreatCrop</i>	Exposure duration for resident and bystander entry into treated crops	<b>0,25</b> hours
<i>d_ReFreqHM</i>	Frequency of hand to mouth activity	<b>9,5</b> events/hour
<i>d_ReTCAd</i>	Transfer coeff. of surface deposits-adult	<b>7300</b> cm <sup>2</sup> /hour
<i>d_ReTCCh</i>	Transfer coeff. of surface deposits-child (1-3 year old)	<b>2600</b> cm <sup>2</sup> /hour
<i>d_SalExt</i>	Saliva extraction percentage	<b>50,0%</b>
<i>d_TcEntryAd</i>	Transfer coefficient for entry into treated crops 75th percentile - adult	<b>7500</b> cm <sup>2</sup> /h
<i>d_TcEntryCh</i>	Transfer coefficient for entry into treated crops 75th percentile - child	<b>2250</b> cm <sup>2</sup> /h
<i>d_TcEntryMeanAd</i>	Transfer coefficient for entry into treated crops mean - adult	<b>5980</b> cm <sup>2</sup> /h
<i>d_TcEntryMeanCh</i>	Transfer coefficient for entry into treated crops mean - child	<b>1794</b> cm <sup>2</sup> /h
<i>d_Turf</i>	Turf transferable residues percentage	<b>5,0%</b>
<i>d_PctExtrapolation</i>	For exposure value 75 percentiles above this amount linear extrapolation is performed	<b>1,5</b> kg
<i>d_head75ProtectionFactor</i>	Coefficient to estimate head protection factor 75 th Percentile	<b>1,79422</b>
<i>d_head95ProtectionFactor</i>	Coefficient to estimate head protection factor 95 Percentile	<b>1,24705</b>

*sys\_KeyOperator* Variables for operator exposure lookup key  
*sys\_OperatorModel* Operator model

**i\_IndoorOutdoor&i\_FormVal&i\_AppMeth&i\_AppEquip&**

**1**

RPE reduction factor	
key_MixRPE, ay_MixRPE	
None	1
FP1, P1 and similar	0,25
FP2, P2 and similar	0,1

PPE reduction factor	
key_MixPPEBody, ay_MixPPEBody	
Potential exposure	1
Work wear - arms, body and legs covered	0,1
Certified protective coverall	0,05

PPE reduction factor	
key_MixPPEHead, ay_MixPPEHead	
None	1
Hood	0,5
Hood and visor	0,05
FP1, P1 and similar	0,8
FP2, P2 and similar	0,8

Application: Gloves PPE reduction factor (depending on formulation type)		
key_AppPPEHands, ay_AppPPEHands		
Wettable powder, soluble powder	Chemical resistant gloves	0,05
Granules, fine granules	Chemical resistant gloves	0,05
Wettable granules, soluble granules	Chemical resistant gloves	0,05
Soluble concentrates, emulsifiable concentrate, etc.	Chemical resistant gloves	0,1
Wettable powder, soluble powder	None	1
Granules, fine granules	None	1
Wettable granules, soluble granules	None	1
Soluble concentrates, emulsifiable concentrate, etc.	None	1

Crop dependent exposure parameters										
key_CropType, ay_CropType	Transfer coefficients	Transfer coefficients	2250	Transfer coefficients	Area Treated					
Crop type	Arm, body and legs covered	Total potential exposure	Activity	hours per day	Body parts involved	Hands, arm, body and legs covered	Type of crop for Resident Bystander	Vehicle Mounted Applications		
Bare soil	NA	NA	NA	NA	NA	NA	Field crops			50
Low berries and other small fruits		3000	5800 Reaching, picking		8 Hand and forearm		750 Field crops			50
Brassica vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops			50
Bulb vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops			50
Cane fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Field crops			10
Cereals		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops			50
Citrus fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops			10
Fruiting vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops			50
Grapes		10100	30000 Hand harvesting		8 Hand and body	no TC available for this assessment	Grapes			10
Grassland and lawns		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops			50
Golf course, turf or other sports lawns		2500	5800 Maintenance		8 Hand and body		580 Field crops			50
Hops		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Hops			50
Leaf vegetables and fresh herbs		2500	5800 Reaching, picking		8 Hand and body		580 Field crops			10
Legume vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops			50
Oilfruits		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops			10
Oilseeds		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops			50
Ornamentals		5000	14000 Cutting, sorting, bundling, carrying		8 Hand and body		1400 Field crops			10
Pome fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops			10
Root and tuber vegetables		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops			50
Stone fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops			10
Tree nuts		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops			10

Resident Spray Drift					
These values are the 75th Percentiles for Residents (assuming average breathing rates for inhalation exposures)					
key_ResidSpray, ay_ResidSpray	Adults Dermal	Children Dermal	Adults Inhalation	Children Inhalation	
Downward spraying2-3		0,47	0,327	0,0001	0,00022
Downward spraying5		0,24	0,22	0,00009	0,00017
Downward spraying10		0,20	0,18	0,00009	0,00013
Upward spraying2-3	NA	NA	NA	NA	
Upward spraying5		5,63	1,689	0,0021	0,00164
Upward spraying10		5,63	1,689	0,0021	0,00164

Bystander Spray Drift					
These values are the 95th Percentiles for Bystanders (assuming high breathing rates for inhalation exposures)					
key_BySpray, ay_BySpray	Adults Dermal	Children Dermal	Adults Inhalation	Children Inhalation	
Downward spraying2-3		1,21	0,74	0,0005	0,0011
Downward spraying5		0,57	0,48	0,00048	0,0008
Downward spraying10		0,48	0,39	0,00051	0,00076
Upward spraying2-3	NA	NA	NA	NA	
Upward spraying5		12,9	3,87	0,0044	0,0035
Upward spraying10		12,9	3,87	0,0044	0,0035

Mean Spray Drift					
These values are the mean values (assuming average breathing rates for inhalation exposures)					
key_AvgSpray, ay_AvgSpray	Adults Dermal	Children Dermal	Adults Inhalation	Children Inhalation	
Downward spraying2-3		0,22	0,18	0,0001	0,0002
Downward spraying5		0,12	0,12	0,0001	0,0001
Downward spraying10		0,11	0,1	0,0001	0,0001
Upward spraying2-3	NA	NA	NA	NA	
Upward spraying5		3,68	1,11	0,0017	0,0013
Upward spraying10		3,68	1,11	0,0017	0,0013

Resident and bystander Surface Deposits Drift percentage			
Ground sediments in % of the application rate calculated on the basis of percentile values (drift data acc. Rautmann)			
key_ByCropType, ay_ByCropType	Bystander surface deposit (90th Percentile)	Resident surface deposit (77th Percentile)	mean
Field cropsnot relevant2-3		0,085	0,056
Field cropsnot relevant5		0,035	0,023
Field cropsnot relevant10		0,019	0,013
Fruit cropsnot relevant2-3		0,292	0,240
Fruit cropsnot relevant5		0,199	0,158
Fruit cropsnot relevant10		0,118	0,090
Fruit cropsearly (without leaves)2-3		0,292	0,240
Fruit cropsearly (without leaves)5		0,199	0,158

Fruit cropsearly (without leaves)10	0,118	0,090	0,061
Fruit cropslate (dense foliage)2-3	0,157	0,110	0,070
Fruit cropslate (dense foliage)5	0,084	0,060	0,037
Fruit cropslate (dense foliage)10	0,036	0,027	0,016
Grapesnot relevant2-3	0,080	0,069	0,053
Grapesnot relevant5	0,036	0,031	0,023
Grapesnot relevant10	0,012	0,010	0,008
Hopsnot relevant2-3	0,193	0,159	0,100
Hopsnot relevant5	0,116	0,086	0,059
Hopsnot relevant10	0,058	0,037	0,029



1	2	3	4	5	6	7	8	9	10	11	12	13	14
<b>Mitch Mit-oid</b>	<b>Outdoor/Indoor</b>	<b>Formulation type</b>	<b>Application method</b>	<b>Application equipment</b>	<b>Type of exposure</b>	<b>Mixing &amp; Loading 75th percentile</b>	<b>Mixing &amp; Loading 95th percentile</b>	<b>Mixing &amp; Loading Comments</b>	<b>Mixing &amp; Loading Model</b>	<b>Application 75th percentile</b>	<b>Application 95th percentile</b>	<b>Application Comments</b>	<b>Application Model</b>
IndoorGranules, fine granulesApplication of granulesManualBody	Indoor	Granules, fine granules	Application of granules	Manual	Body			Value for application is for combination of mixing&loading and application	PHED	68,8708	253,4438	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
IndoorGranules, fine granulesApplication of granulesManualHands	Indoor	Granules, fine granules	Application of granules	Manual	Hands			Value for application is for combination of mixing&loading and application	PHED	26,5320	94,3636	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
IndoorGranules, fine granulesApplication of granulesManualInhalation	Indoor	Granules, fine granules	Application of granules	Manual	Inhalation			Value for application is for combination of mixing&loading and application	PHED	0,4677	1,5251	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesBroadcast application of granulesVehicle-mountedBody	Outdoor	Granules, fine granules	Broadcast application of granules	Vehicle-mounted	Body	0,0162	0,0427	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0,0047	0,0151	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesBroadcast application of granulesVehicle-mountedHands	Outdoor	Granules, fine granules	Broadcast application of granules	Vehicle-mounted	Hands	0,0015	0,0069	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0,0004	0,0018	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesBroadcast application of granulesVehicle-mountedInhalation	Outdoor	Granules, fine granules	Broadcast application of granules	Vehicle-mounted	Inhalation	0,0208	0,0784		PHED	0,0012	0,0045		PHED
OutdoorGranules, fine granulesIn furrow application of granulesVehicle-mountedBody	Outdoor	Granules, fine granules	In furrow application of granules	Vehicle-mounted	Body	0,0162	0,0427	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0,0047	0,0151	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesIn furrow application of granulesVehicle-mountedHands	Outdoor	Granules, fine granules	In furrow application of granules	Vehicle-mounted	Hands	0,0015	0,0069	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0,0004	0,0018	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesIn furrow application of granulesVehicle-mountedInhalation	Outdoor	Granules, fine granules	In furrow application of granules	Vehicle-mounted	Inhalation	0,0208	0,0784		PHED	0,0012	0,0045		PHED
OutdoorGranules, fine granulesManual application of granulesManualBody	Outdoor	Granules, fine granules	Manual application of granules	Manual	Body			Value for application is for combination of mixing&loading and application	PHED	68,8708	253,4438	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesManual application of granulesManualHands	Outdoor	Granules, fine granules	Manual application of granules	Manual	Hands			Value for application is for combination of mixing&loading and application	PHED	26,5320	94,3636	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesManual application of granulesManualInhalation	Outdoor	Granules, fine granules	Manual application of granules	Manual	Inhalation			Value for application is for combination of mixing&loading and application	PHED	0,4677	1,5251	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED



