

## **Instructions for using the workbook**

The information required for the exposure assessment needs to be entered in the worksheet "**Data entry**".

In the following worksheets formulas calculate the exposure values automatically

Worksheet "**Operator Outdoor Spray AOEM**" is to be for outdoor spray applications. PPE options can be selected in this worksheet

Worksheet "**Operator Granules**" is for granular applications. Currently the calculator does not allow operator exposure for indoor applications. PPE options can be selected in this worksheet

Worksheets "**Resident exposure**" and "**Bystander exposure**" are only relevant for outdoor applications

Worksheet "**Recreational Exposure**" is only applicable for golf course, turf, other sports lawns or amenity turf/grassland areas where members of the public are likely to have access

The combined results of the exposure assessment are presented in worksheet "**Summary**"

This calculator should be used in conjunction with the **Guidance on the Assessment of Exposure for Operators, Workers, Residents and Bystanders in Risk Assessment for Plant Protection Products**

Latest version: 23 Oct 2014 - Version produced to support guidance document published 23/10/2014

Note: Some drop-down menus depend on others. To avoid errors, please fill-in from top to bottom

<b>Substance name</b>	chloorthalonil
<b>Product name</b>	Allure Vloeibaar
<b>Reference value non acutely toxic active substance (RVNAS)</b>	0.009 mg/kg bw/day
<b>Reference value acutely toxic active substance (RVAAS)</b>	0.009 mg/kg bw/day
<b>Crop type</b>	Ornamentals
<b>Substance properties</b>	
Formulation type	Soluble concentrates, emulsifiable concentrate, etc.
Minimum volume water for application (liquids)	150 L/ha
Maximum application rate of active substance	0.66 kg a.s. /ha
50% Dissipation Time DT50	30 days
Initial Dislodgeable Foliar Residue	3 µg/cm <sup>2</sup> of foliage/kg a.s. applied/ha
Dermal absorption of product	0.02%
Dermal absorption of in-use dilution	0.34%
Oral absorption of active substance	30.00%
Inhalation absorption of active substance	100.00%
Vapour pressure of active substance	low volatile substances having a vapour pressure of <5*10 <sup>-3</sup> Pa
<b>Scenario</b>	
Indoor or Outdoor application	Outdoor
Application method	Downward spraying
Application equipment	Vehicle-mounted
Buffer strip	2-3 m
Number of applications	2
Interval between multiple applications	10 days
Season (upward spraying orchards only)	not relevant

### Exposure assessment

Substance	chloorthalonil	Formulation = Soluble concentrates, emulsifiable concentrate, etc.	Application rate-0,66 kg a.s. /ha	Spray dilution = 4,4 g a.s./l	Vapour pressure = low volatile substances having a vapour pressure of <math>5 \times 10^{-3}</math>Pa
Scenario	Ornamentals / Outdoor / Downward spraying / Vehicle-mounted			Buffer = 2-3	Number applications = 2, Application interval = 10 days
Percentage Absorption	Dermal for product = 0,02	Dermal for in use dilution = 0,34	Oral = 30	Inhalation = 100	
RVNAS	0,009 mg/kg bw/day		RVAAS	0,009 mg/kg bw/day	
DFR	3 µg a.s./cm <sup>2</sup> per kg a.s./ha		DT50	30 days	

<b>Operator Model</b>		Mixing, loading and application AOEM			
Potential exposure	Longer term systemic exposure mg/kg bw/day	0.0019	% of RVNAS	21.04%	
	Acute systemic exposure mg/kg bw/day	0.0052	% of RVAAS	58.20%	
Mixing and Loading	Gloves = No	Clothing = Work wear - arms, body and legs covered	RPE = None	Soluble bags = No	
Application	Gloves = No	Clothing = Work wear - arms, body and legs covered	RPE = None	Closed cabin = No	
Exposure (including PPE options above)	Longer term systemic exposure mg/kg bw/day	0.0010	% of RVNAS	11.43%	
	Acute systemic exposure mg/kg bw/day	0.0038	% of RVAAS	42.05%	

<b>Worker - Cutting, sorting, bundling, carrying</b>	Potential exposure mg/kg bw/day	0.0225	% of RVNAS	250.45%
	Working clothing mg/kg bw/day	0.0081	% of RVNAS	89.45%
	Working clothing and gloves mg/kg bw/day	0.0023	% of RVNAS	25.04%

<b>Resident - child</b>	Spray drift (75th percentile) mg/kg bw/day	0.0005	% of RVNAS	5.53%
	Vapour (75th percentile) mg/kg bw/day	0.0011	% of RVNAS	11.89%
	Surface deposits (75th percentile) mg/kg bw/day	0.0003	% of RVNAS	3.86%
	Entry into treated crops (75th percentile) mg/kg bw/day	0.0007	% of RVNAS	7.55%
	All pathways (mean) mg/kg bw/day	0.0021	% of RVNAS	23.18%
<b>Resident - adult</b>	Spray drift (75th percentile) mg/kg bw/day	0.0001	% of RVNAS	1.15%
	Vapour (75th percentile) mg/kg bw/day	0.0002	% of RVNAS	2.56%
	Surface deposits (75th percentile) mg/kg bw/day	0.0000	% of RVNAS	0.30%
	Entry into treated crops (75th percentile) mg/kg bw/day	0.0004	% of RVNAS	4.19%
	All pathways (mean) mg/kg bw/day	0.0006	% of RVNAS	6.70%

<b>Bystander - child</b>	Spray drift (95th percentile) mg/kg bw/day	0.0014	% of RVAAS	15.56%
	Vapour (95th percentile) mg/kg bw/day	0.0011	% of RVAAS	11.89%
	Surface deposits (95th percentile) mg/kg bw/day	0.0009	% of RVAAS	10.36%

**Exposure assessment**

	Entry into treated crops (95th percentile) mg/kg bw/day	0.0007	% of RVAAS	7.55%
<b>Bystander - adult</b>	Spray drift (95th percentile) mg/kg bw/day	0.0003	% of RVAAS	3.16%
	Vapour (95th percentile) mg/kg bw/day	0.0002	% of RVAAS	2.56%
	Surface deposits (95th percentile) mg/kg bw/day	0.0001	% of RVAAS	0.92%
	Entry into treated crops (95th percentile) mg/kg bw/day	0.0004	% of RVAAS	4.19%

<b>Recreational Exposure</b>	Child % of RVNAS	Adult % of RVNAS
------------------------------	------------------	------------------

Latest version: 23 Oct 2014 - Version produced to support guidance document published 23/10/2014

**Operator exposure for Allure Vloeibaar outdoor spray applications**

Application rate of active substance	0.66 kg a.s./ha	<i>i_AppRate</i>
Assumed area treated	10 ha/day	<i>d_AreaTreated</i>
Amount of active substance applied	6.6 kg a.s./day	<i>i_AmountAS</i>
Dermal absorption of the product	0.02%	<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	0.34%	<i>i_AbsorInuse</i>
Formulation type	Soluble concentrates, emulsifiable concentrate, etc.	
Indoor or Outdoor application	Outdoor	
Application method	Downward spraying	
Application equipment	Vehicle-mounted	
Season	not relevant	

	Exposure values	µg exposure/day mixed and loaded		Reference	Comment
		75 <sup>th</sup> centile	95 <sup>th</sup> centile		
Mixing and loading	Hands	20762	77352	AOEM	
	Body	13440	124617	AOEM	
	Head	342	33158	AOEM	
	Protected hands (gloves)	118	1307	AOEM	
	Protected body (workwear or protective garment and sturdy footwear)	127	965	AOEM	
	Protected head (hood and face shield)	5	1877	AOEM	
	Inhalation	6	30	AOEM	
	<b>Protective Equipment</b>	Select for inclusion		Penetration factor	Inhalation Protection factor
	Gloves	No			
	Clothing	Work wear - arms, body and legs covered		Incl. in AOEM model	
Head and respiratory PPE	None		1	1	
Water soluble bag	No		1		

	Exposure values	µg exposure/day applied		Reference	Comment
		75 <sup>th</sup> centile	95 <sup>th</sup> centile		
Application	Hands	10694	14930	AOEM	This scenario assumes that small area equipment is used
	Body	14673	18592	AOEM	
	Head	88	1031	AOEM	
	Protected hands (gloves)	61	27	AOEM	
	Protected body (workwear or protective garment and sturdy footwear)	183	217	AOEM	
	Inhalation	14	120	AOEM	
	<b>Protective Equipment</b>	Select for inclusion		Penetration factor	
	Gloves	No			
	Clothing	Work wear - arms, body and legs covered		Incl. in AOEM model	
	Head and respiratory PPE	None		1	1
Closed cab	No		vehicle mounted upward spraying only		

1. Total			
	Without RPE/PPE	With RPE/PPE	
<b>Longer term</b>			
Total systemic exposure from mixing, loading and application (mg a.s./day)	0.1136284	0.0617022	
Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	0.0018938	0.0010284	
% of RVNAS	21.04%	11.43%	
<b>Acute</b>			
Total systemic exposure from mixing, loading and application (mg a.s./day)	0.3142533	0.2270464	

Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	0.0052376	0.0037841	
% of RVAAS	58.20%	42.05%	

## 2. Longer term exposure

### 2.1 Mixing and loading

	Systemic exposure [ $\mu\text{g a.s. /day}$ ]	Systemic exposure [ $\mu\text{g a.s./kg}$ ]	Formula
<b>Without RPE/PPE</b>			
Hands	4.1524031	0.0692067	$D15^*i\_AbsorpProduct$
Body	2.6880789	0.0448013	$D16^*i\_AbsorpProduct$
Head	0.0684863	0.0011414	$D17^*i\_AbsorpProduct$
Inhalation	6.4907708	0.1081795	$D21^*i\_AbsorpInhalation$
Sum	13.3997391	0.2233290	
<b>With RPE/PPE (as selected above)</b>			
Hands	4.1524031	0.0692067	$D18^*i\_AbsorpProduct$
Body	0.0253186	0.0004220	$D19^*i\_AbsorpProduct$ or $D15^*i\_AbsorpProduct^*F24$
Head	0.0684863	0.0011414	$D20^*i\_AbsorpProduct$ or $D17^*i\_AbsorpProduct^*F25$
Inhalation	6.4907708	0.1081795	$D21^*i\_AbsorpInhalation^*G25$
Sum	10.7369788	0.1789496	
Water soluble bag	10.7369788	0.1789496	$C70^*F26$

### 2.2 Application

	Systemic exposure [ $\mu\text{g a.s. /day}$ ]	Systemic exposure [ $\mu\text{g a.s./kg}$ ]	Formula
<b>Without RPE/PPE</b>			
Hands	36.3601312	0.6060022	$D30^*i\_Absorpinuse$
Body	49.8872455	0.8314541	$D31^*i\_Absorpinuse$
Head	0.2986862	0.0049781	$D32^*i\_Absorpinuse$
Inhalation	13.6825730	0.2280429	$D35^*i\_AbsorpInhalation$
Sum	100.2286359	1.6704773	
<b>With RPE/PPE (as selected above)</b>			
Hands	36.3601312	0.6060022	$D33^*i\_Absorpinuse$
Body	0.6238741	0.0103979	$D34^*i\_Absorpinuse$ or $D31^*i\_Absorpinuse^*F38$
Head	0.2986862	0.0049781	$D32^*i\_Absorpinuse^*F39$
Inhalation	13.6825730	0.2280429	$D35^*i\_Absorpinuse^*G39$
Sum	50.9652645	0.8494211	

## 3. Acute exposure

### 3.1 Mixing and loading

	Systemic exposure [ $\mu\text{g a.s. /day}$ ]	Systemic exposure [ $\mu\text{g a.s./kg}$ ]	Formula
<b>Without RPE/PPE</b>			
Hands	15.4704155	0.2578403	$E15^*i\_AbsorpProduct$
Body	24.9233406	0.4153890	$E16^*i\_AbsorpProduct$
Head	6.6316856	0.1105281	$E17^*i\_AbsorpProduct$
Inhalation	30.0350294	0.5005838	$E21^*i\_AbsorpInhalation$
Sum	77.0604711	1.2843412	
<b>With RPE/PPE (as selected above)</b>			
Hands	15.4704155	0.2578403	$E18^*i\_AbsorpProduct$
Body	0.1930518	0.0032175	$E19^*i\_AbsorpProduct$ or $E16^*i\_AbsorpProduct^*F24$
Head	6.6316856	0.1105281	$E20^*i\_AbsorpProduct$ or $E17^*i\_AbsorpProduct^*F25$
Inhalation	30.0350294	0.5005838	$E21^*i\_AbsorpInhalation^*G25$
Sum	52.3301824	0.8721697	
Water soluble bag	52.3301824	0.8721697	$C104^*F26$

### 2.2 Application

	Systemic exposure [ $\mu\text{g a.s. /day}$ ]	Systemic exposure [ $\mu\text{g a.s./kg}$ ]	Formula
<b>Without RPE/PPE</b>			
Hands	50.7623370	0.8460389	$E30^*i\_Absorpinuse$
Body	63.2139151	1.0535653	$E31^*i\_Absorpinuse$
Head	3.5062306	0.0584372	$E32^*i\_Absorpinuse$
Inhalation	119.7103599	1.9951727	$E35^*i\_AbsorpInhalation$
Sum	237.1928427	3.9532140	
<b>With RPE/PPE (as selected above)</b>			
Hands	50.7623370	0.8460389	$E33^*i\_Absorpinuse$
Body	0.7372624	0.0122877	$E34^*i\_Absorpinuse$ or $E31^*i\_Absorpinuse^*F38$
Head	3.5062306	0.0584372	$E32^*i\_Absorpinuse^*F39$
Inhalation	119.7103599	1.9951727	$E35^*i\_AbsorpInhalation^*G39$

Sum	174.7161899	2.9119365	
-----	-------------	-----------	--

**Operator exposure for Allure Vloeibaar granular applications**

Application rate of active substance	0.66 kg a.s./ha	<i>i_AppRate</i>
Assumed area treated	10 ha/day	<i>d_AreaTreated</i>
Amount of active substance applied	6.6 kg a.s./day	<i>i_AmountAS</i>
Dermal absorption of the product	0.02%	<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	0.34%	<i>i_AbsorInuse</i>
Formulation type	Soluble concentrates, emulsifiable concentrate, etc.	
Indoor or Outdoor application	Outdoor <b>This sheet is only to be used for granular applications</b>	
Application method	Downward spraying	
Application equipment	Vehicle-mounted	

	Exposure values	mg exposure/kg a.s. mixed and loaded		Reference	Comment
		75 <sup>th</sup> centile	95 <sup>th</sup> centile		
Mixing and loading	Hands	#N/A	#N/A	#N/A	#N/A
	Body	#N/A	#N/A	#N/A	#N/A
	Inhalation	#N/A	#N/A	#N/A	#N/A
	<b>Protective Equipment</b>	Choose item		Penetration factor	
	Gloves	Chemical resistant gloves			Protection for granules exposure is based on measured values
	Body PPE	Certified protective coverall			
	RPE	None		1	

	Exposure values	mg exposure/kg a.s. applied		Reference	Comment
		75 <sup>th</sup> centile	95 <sup>th</sup> centile		
Application	Hands	#N/A	#N/A	#N/A	#N/A
	Body	#N/A	#N/A	#N/A	#N/A
	Inhalation	#N/A	#N/A	#N/A	#N/A
	<b>Protective Equipment</b>	Choose item		Penetration factor	
	Gloves	Chemical resistant gloves			Protection for granules exposure is based on measured values
	Body PPE	Certified protective coverall			
	RPE	FP1, P1 and similar		0.25	

**1. Total**

	Without RPE/PPE	With RPE/PPE
<b>Longer term</b>		
Total systemic exposure from mixing, loading and application (mg a.s./day)	#N/A	#N/A
Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	#N/A	#N/A
% of RVNAS	#N/A	#N/A
<b>Acute</b>		
Total systemic exposure from mixing, loading and application (mg a.s./day)	#N/A	#N/A



Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	#N/A	#N/A	
% of RVAAS	#N/A	#N/A	

**2. Longer term exposure**

**2.1 Mixing and loading**

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
<b>Without RPE/PPE</b>			
Hands	#N/A	#N/A	$D14*100*i\_AmoutAS*i\_AbsorpProduct$
Body	#N/A	#N/A	$D15*100*i\_AmoutAS*i\_AbsorpProduct$
Inhalation	#N/A	#N/A	$D16*i\_AmoutAS*i\_AbsorpInhalation$
Sum	#N/A	#N/A	
<b>With RPE/PPE (as selected above)</b>			
Hands	#N/A	#N/A	$D14*i\_AmoutAS*i\_AbsorpProduct$
Body	#N/A	#N/A	$D15*i\_AmoutAS*i\_AbsorpProduct$
Inhalation	#N/A	#N/A	$D16*i\_AmoutAS*i\_AbsorpInhalation*F20$
Sum	#N/A	#N/A	

**2.2 Application**

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
<b>Without RPE/PPE</b>			
Hands	#N/A	#N/A	$D25*100*i\_AmoutAS*i\_Absorpnuse$
Body	#N/A	#N/A	$D26*100*i\_AmoutAS*i\_Absorpnuse$
Inhalation	#N/A	#N/A	$D27*i\_AmoutAS*i\_AbsorpInhalation$
Sum	#N/A	#N/A	
<b>With RPE/PPE (as selected above)</b>			
Hands	#N/A	#N/A	$D25*i\_AmoutAS*i\_Absorpnuse$
Body	#N/A	#N/A	$D26*i\_AmoutAS*i\_Absorpnuse$
Inhalation	#N/A	#N/A	$D27*i\_AmoutAS*i\_AbsorpInhalation*F31$
Sum	#N/A	#N/A	

**3. Acute exposure**

**3.1 Mixing and loading**

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
<b>Without RPE/PPE</b>			
Hands	#N/A	#N/A	$E14*100*i\_AmoutAS*i\_AbsorpProduct$
Body	#N/A	#N/A	$E15*100*i\_AmoutAS*i\_AbsorpProduct$
Inhalation	#N/A	#N/A	$E16*i\_AmoutAS*i\_AbsorpInhalation$
Sum	#N/A	#N/A	
<b>With RPE/PPE (as selected above)</b>			
Hands	#N/A	#N/A	$E14*100*i\_AmoutAS*i\_AbsorpProduct$
Body	#N/A	#N/A	$E15*100*i\_AmoutAS*i\_AbsorpProduct$
Inhalation	#N/A	#N/A	$E16*i\_AmoutAS*i\_AbsorpInhalation*F20$
Sum	#N/A	#N/A	

**3.2 Application**

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
<b>Without RPE/PPE</b>			
Hands	#N/A	#N/A	$E25*100*i\_AmoutAS*i\_Absorpnuse$
Body	#N/A	#N/A	$E25*100*i\_AmoutAS*i\_Absorpnuse$
Inhalation	#N/A	#N/A	$E26*i\_AmoutAS*i\_AbsorpInhalation$
Sum	#N/A	#N/A	
<b>With RPE/PPE (as selected above)</b>			

Hands	#N/A	#N/A	$E25*100*i\_AmoutAS*i\_Absorplnuse$
Body	#N/A	#N/A	$E26*100*i\_AmoutAS*i\_Absorplnuse$
Inhalation	#N/A	#N/A	$E27*i\_AmoutAS*i\_Absorplnhalation*F31$
Sum	#N/A	#N/A	

## Worker exposure from residues on foliage for Allure Vloeibaar

Crop type	Ornamentals	
Indoor or outdoor	Outdoor	
Application method	Downward spraying	
Application equipment	Vehicle-mounted	
Worker's task	Cutting, sorting, bundling, carrying	
Main body parts in contact with foliage	Hand and body	
Application rate of active substance	0.66 kg a.s./ha	<i>i_AppRate</i>
Number of applications	2	<i>i_AppNo</i>
Interval between multiple applications	10 days	<i>i_AppInt</i>
Half-life of active substance	30 days	<i>d_HalfLifeAS</i>
Multiple application factor	1.8	<i>d_MAF</i>
Dermal absorption of the product	0.02%	<i>i_AbsorpProduct</i>
Dermal absorption of the in-use dilution	0.34%	<i>i_Absorplnuse</i>
Dislodgeable foliar residue ( $i\_AppRate * i\_DFR$ )	1.98 $\mu\text{g a.s./cm}^2$	<i>d_DFR</i>
Working hours	8 hr	<i>d_WorkHr</i>
Dermal transfer coefficient - Total potential exposure	14000 $\text{cm}^2/\text{hr}$	<i>d_DermTcUCV</i>
Dermal transfer coefficient - arms, body and legs covered	5000 $\text{cm}^2/\text{hr}$	<i>d_DermTcCV1</i>
Dermal transfer coefficient - hands, arms, body and legs covered	1400 $\text{cm}^2/\text{hr}$	<i>d_DermTcCV2</i>
Inhalation transfer coefficient for automated applications	NA $\text{ha/hr} * 10^{(-3)}$	<i>d_InhalTcAut</i>
Inhalation transfer coefficient for cutting ornamentals	NA $\text{ha/hr} * 10^{(-3)}$	<i>d_InhalTcCut</i>
Inhalation transfer coefficient for sorting / bundling ornamentals	NA $\text{ha/hr} * 10^{(-3)}$	<i>d_InhalTcSort</i>

### 1. Total

	Potential exposure	Work wear - arms, body and legs covered	Working wear and gloves	Comments
Total systemic exposure (mg a.s./day)	1.3524215	0.4830077	0.1352421	
Total systemic exposure per kg body weight (mg/kg bw/day)	0.0225404	0.0080501	0.0022540	
% of RVNAS	250.45%	89.45%	25.04%	

### 2. Details

	Systemic exposure		Formula	Comments
	[mg a.s. /day]	[mg a.s./kg bw/day]		
Dermal - Potential	1.3524215	0.0225404	$d\_DermTcUCV * d\_WorkHr * i\_DFR * i\_MAF / 1000 * i\_Absorplnuse$	
Dermal - Work wear - arms, body and legs covered	0.4830077	0.0080501	$d\_DermTcCV1 * d\_WorkHr * d\_DFR * d\_MAF / 1000 * i\_Absorplnuse$	
Dermal - Working wear and gloves	0.1352421	0.0022540	$d\_DermTcCV2 * d\_WorkHr * d\_DFR * d\_MAF / 1000 * i\_Absorplnuse$	
Inhalation				Na for outdoor activities

Resident exposure for Allure Vloeibaar

Croptype	Ornamentals		
Application method	Downward spraying		
Application equipment	Vehicle-mounted		<i>i_AppEquip</i>
Formulation type	Soluble concentrates, emulsifiable concentrate, etc.		<i>i_FormVal</i>
Buffer strip	2-3 m		<i>i_Buffer</i>
Application rate of the product	0.66 kg a.s./ha		<i>i_AppRate</i>
Concentration of active substance (in-use dilution for liquid applications)	4.4 g a.s./l		<i>d_ConcAS</i>
Dermal absorption of product	0.02%		<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	0.34%		<i>i_AbsorpInuse</i>
Oral absorption	30.00%		<i>i_AbsorpOrallnuse</i>
Dislodgeable foliar residue ( <i>i_AppRate</i> * <i>i_DFR</i> )	1.98 µg a.s./cm <sup>2</sup>		<i>d_DFR</i>
Vapour pressure of in-use dilution	low volatile substances having a vapour pressure of <5*10 <sup>-3</sup> Pa	Pa	<i>i_Volat</i>
Concentration in air	0.001 mg/m <sup>3</sup>		<i>d_AirCon</i>
Resident dermal spray drift exposure 75th percentile - adult	0.47 ml spray dilution/person		
Resident dermal spray drift exposure 75th percentile - child	0.327 ml spray dilution/person		
Resident inhal. spray drift exposure 75th percentile - adult	0.00010 ml spray dilution/person		
Resident inhal. spray drift exposure 75th percentile - child	0.00022 ml spray dilution/person		
Resident dermal spray drift exposure mean - adult	0.22318 ml spray dilution/person		
Resident dermal spray drift exposure mean - child	0.18 ml spray dilution/person		
Resident inhal. spray drift exposure mean - adult	0.00009 ml spray dilution/person		
Resident inhal. spray drift exposure mean - child	0.00017 ml spray dilution/person		
Exposure duration dermal	2 hours		<i>d_ReExpDur</i>
Exposure duration inhalation	24 hours		<i>d_ReExpDurInhal</i>
Exposure duration entry into treated crops	0.25 hours		<i>d_ExpDurTreatCrop</i>
Light clothing adjustment factor	18.0%		<i>d_ClothAF</i>
Breathing rate adult	0.23 m <sup>3</sup> /day/kg		<i>d_BreathRAD</i>
Breathing rate child (1-3 year old)	1.07 m <sup>3</sup> /day/kg		<i>d_BreathRCh</i>
Drift percentage on surface (75th percentile)	5.60%		
Drift percentage on surface (mean)	4.10%		
Turf transferable residues percentage	5.00%		<i>d_Turf</i>
Transfer coeff. of surface deposits-adult	7300 cm <sup>2</sup> /hour		<i>d_ReTCAd</i>
Transfer coeff. of surface deposits-child (1-3 year old)	2600 cm <sup>2</sup> /hour		<i>d_ReTCCh</i>
Saliva extraction percentage	50.00%		<i>d_SalExt</i>
Surface area of hands mouthed	20 cm <sup>2</sup>		<i>d_AreaHM</i>
Frequency of hand to mouth activity	9.5 events/hour		<i>d_ReFreqHM</i>
Ingestion rate for mouthing of grass per day	25 cm <sup>2</sup>		<i>d_MouthGrass</i>
Dislodgeable residues percentage transferability for object to mouth	20.00%		<i>d_DRP</i>
Transfer coefficient for entry into treated crops (75th percentile) - adult	7500 cm <sup>2</sup> /h		<i>d_TcEntryAd</i>
Transfer coefficient for entry into treated crops (75th percentile) - child	2250 cm <sup>2</sup> /h		<i>d_TcEntryCh</i>
Transfer coefficient for entry into treated crops (mean) - adult	5980 cm <sup>2</sup> /h		<i>d_TcEntryAd</i>
Transfer coefficient for entry into treated crops (mean) - child	1794 cm <sup>2</sup> /h		<i>d_TcEntryCh</i>

1. Total

1.1 1-3 year old child

	Spray drift (75th percentile)	Vapour (75th percentile)	Surface deposits (75th percentile)	Entry into treated crops (75th percentile)	All pathways (mean)
Total systemic exposure (mg a.s./day)	0.0049794	0.0107000	0.0034699	0.0067923	0.0208643
Total systemic exposure per kg body weight (mg/kg bw/day)	0.0004979	0.0010700	0.0003470	0.0006792	0.0020864
% of RVNAS	5.53%	11.89%	3.86%	7.55%	23.18%

1.2 Adult

	Spray drift	Vapour	Surface deposits	Entry into treated crops	All pathways (mean)
Total systemic exposure (mg a.s./day)	0.0062056	0.0138000	0.0016454	0.0226410	0.0361909
Total systemic exposure per kg body weight (mg/kg bw/day)	0.0001034	0.0002300	0.0000274	0.0003773	0.0006032
% of RVNAS	1.15%	2.56%	0.30%	4.19%	6.70%

2. Resident exposure 75th Percentile

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
<b>1-3 year old child</b>				
Spray drift	0.0049794	0.0004979	$((C16 * i\_AbsorpInuse * (1 - d\_ClothAF)) + C18) * d\_ConcAS$	
Vapour	0.0107000	0.0010700	$d\_AirCon * d\_BreathRCh * d\_BwChild$	
<b>Surface deposits</b>				
Dermal	0.0005860	0.0000586	$(i\_AppRate / 100) * C29 * d\_Turf * d\_ReTCCh * d\_ReExpDur * MAX(i\_AbsorpProduct, i\_AbsorpInuse) * d\_MAF * IF(i\_AppEquip = "Vehicle-mounted-Drift Reduction", 0.5, 1)$	
Hand to mouth	0.0018894	0.0001889	$(i\_AppRate / 100) * C29 * d\_Turf * d\_SalExt * d\_AreaHM * d\_ReFreqHM * d\_ReExpDur * i\_AbsorpOrallnuse * d\_MAF$	
Object to mouth	0.0009944	0.0000994	$(i\_AppRate / 100) * C29 * d\_DRP * d\_MouthGrass * i\_AbsorpOrallnuse * d\_MAF$	

Entry into treated crops				
Dermal	0.0067923	0.0006792	$(d\_TcEntryCh*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_Absorplnuse)$	
Hand to mouth			$(i\_AppRate/100)*d\_Turf*d\_MAF*d\_SalExt*d\_AreaHM*d\_ReFreqHM*d\_ReExpDur*i\_AbsorpOrallnuse$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Object to mouth			$(i\_AppRate/100)*d\_DRP*d\_MouthGrass*i\_AbsorpOrallnuse*d\_MAF$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
<b>Adult</b>				
Spray drift	0.0062056	0.0001034	$(C15*i\_Absorplnuse*(1-d\_ClothAF))+C17)*d\_ConcAS$	
Vapour	0.0138000	0.0002300	$d\_AirCon*d\_BreathRAD*d\_BwAdult$	
Surface deposits (dermal)	0.0016454	0.0000274	$(i\_AppRate/100)*C30*d\_Turf*d\_ReTCA*d\_ReExpDur*i\_Absorplnuse$	
Entry into treated crops (dermal)	0.0226410	0.0003773	$(d\_TcEntryAd*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_Absorplnuse)$	

### 3. Summing of exposure pathways mean

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
<b>1-3 year old child</b>				
Spray drift	0.0022081	0.0002208	$((C20*i\_Absorplnuse*(1-d\_ClothAF))+C22)*d\_ConcAS$	
Vapour	0.0107000	0.0010700	$d\_AirCon*d\_BreathRCh*d\_BwChild$	
Surface deposits				
Dermal	0.0004291	0.0000429	$(i\_AppRate/100)*C30*d\_Turf*d\_ReTCA*d\_ReExpDur*MAX(i\_AbsorpProduct,i\_Absorplnuse)*d\_MAF*IF(i\_AppEquip = "Vehicle-mounted-Drift Reduction",0.5,1)$	
Hand to mouth	0.0013833	0.0001383	$(i\_AppRate/100)*C30*d\_Turf*d\_SalExt*d\_AreaHM*d\_ReFreqHM*d\_ReExpDur*i\_AbsorpOrallnuse*d\_MAF$	
Object to mouth	0.0007281	0.0000728	$(i\_AppRate/100)*C30*d\_DRP*d\_MouthGrass*i\_AbsorpOrallnuse*d\_MAF$	
Entry into treated crops				
Dermal	0.0054157	0.0005416	$(d\_TcEntryMeanCh*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_Absorplnuse)$	
Hand to mouth			$(i\_AppRate/100)*1*d\_Turf*d\_MAF*d\_SalExt*d\_AreaHM*d\_ReFreqHM*d\_ReExpDur*i\_AbsorpOrallnuse$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Object to mouth			$(i\_AppRate/100)*1*d\_DRP*d\_MouthGrass*i\_AbsorpOrallnuse*d\_MAF$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
<b>Adult</b>				
Spray drift	0.0031338	0.0000522	$"(C19*i\_Absorplnuse*(1-d\_ClothAF))+C21)*d\_ConcAS"$	
Vapour	0.0138000	0.0002300	$d\_AirCon*d\_BreathRAD*d\_BwAdult$	
Surface deposits (dermal)	0.0012047	0.0000201	$(i\_AppRate/100)*C30*d\_Turf*d\_ReTCA*d\_ReExpDur*MAX(i\_AbsorpProduct,i\_Absorplnuse)*d\_MAF*IF(i\_AppEquip = "Vehicle-mounted-Drift Reduction",0.5,1)$	
Entry into treated crops (dermal)	0.0180524	0.0003009	$(d\_TcEntryMeanAd*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_Absorplnuse)$	

## Bystander exposure for Allure Vloeibaar

Croptype	<b>Ornamentals</b>		
Application method	<b>Downward spraying</b>		
Application equipment	<b>Vehicle-mounted</b>		<i>i_AppEquip</i>
Formulation type	<b>Soluble concentrates, emulsifiable concentrate, etc.</b>		
Application rate of the product	<b>0.66</b> kg a.s./ha		<i>i_AppRate</i>
Buffer strip	<b>2-3</b> m		<i>i_Buffer</i>
Concentration of active substance (in-use dilution for liquid applications)	<b>4.4</b> g a.s./l		<i>d_ConcAS</i>
Dermal absorption of product	<b>0.02%</b>		<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	<b>0.34%</b>		<i>i_AbsorpInuse</i>
Oral absorption	<b>30.00%</b>		<i>i_AbsorpOrallnuse</i>
Dislodgeable foliar residue ( $i\_AppRate * i\_DFR$ )	<b>1.98</b> µg a.s./cm <sup>2</sup>		<i>d_DFR</i>
Vapour pressure of in-use dilution	<b>low volatile substances having a vapour pressure of &lt;5*10<sup>-3</sup>Pa</b>	Pa	<i>i_Volat</i>
Concentration in air	<b>0.001</b> mg/m <sup>3</sup>		<i>d_AirCon</i>
Bystander dermal spray drift exposure - adult	<b>1.21</b> ml spray dilution/person		
Bystander dermal spray drift exposure - child	<b>0.74</b> ml spray dilution/person		
Bystander inhal. spray drift exposure - adult	<b>0.00050</b> ml spray dilution/person		
Bystander inhal. spray drift exposure - child	<b>0.00112</b> ml spray dilution/person		
Exposure duration	<b>2</b> hours		<i>d_ByExpDur</i>
Exposure duration entry into treated crops	<b>0.25</b> hours		<i>d_ExpDurTreatCrop</i>
Light clothing adjustment factor	<b>18.0%</b>		<i>d_ClothAF</i>
Breathing rate adult	<b>0.23</b> m <sup>3</sup> /hours/kg		<i>d_BreathRAd</i>
Breathing rate child (1-3 year old)	<b>1.07</b> m <sup>3</sup> /hours/kg		<i>d_BreathRCh</i>
Drift percentage on surface (90th percentile)	<b>8.50%</b>		
Turf transferable residues percentage	<b>5.00%</b>		<i>d_Turf</i>
Transfer coeff. of surface deposits-adult	<b>14500</b> cm <sup>2</sup> /hour		<i>d_ByTCAd</i>
Transfer coeff. of surface deposits-child (1-3 year old)	<b>5200</b> cm <sup>2</sup> /hour		<i>d_ByTCCh</i>
Saliva extraction percentage	<b>50.00%</b>		<i>d_SalExt</i>
Surface area of hands mouthed	<b>20</b> cm <sup>2</sup>		<i>d_AreaHM</i>
Frequency of hand to mouth activity	<b>20</b> events/hour		<i>d_ByFreqHM</i>
Ingestion rate for mouthing of grass per day	<b>25</b> cm <sup>2</sup>		<i>d_MouthGrass</i>
Dislodgeable residues percentage transferability for object to mouth	<b>20.00%</b>		<i>d_DRP</i>
Transfer coefficient for entry into treated crops - adult	<b>7500</b> cm <sup>2</sup> /h		<i>d_TcEntryAd</i>
Transfer coefficient for entry into treated crops - child	<b>2250</b> cm <sup>2</sup> /h		<i>d_TcEntryCh</i>

### 1. Total

#### 1.1 1-3 year old child

	Spray drift	Vapour	Surface deposits	Entry into treated crops
Total systemic exposure (mg a.s./day)	<b>0.0140057</b>	<b>0.0107000</b>	<b>0.0093261</b>	<b>0.0067923</b>
Total systemic exposure per kg body weight (mg/kg bw/day)	<b>0.0014006</b>	<b>0.0010700</b>	<b>0.0009326</b>	<b>0.0006792</b>
% of RVAAS	<b>15.56%</b>	<b>11.89%</b>	<b>10.36%</b>	<b>7.55%</b>

#### 1.2 Adult

	Spray drift	Vapour	Surface deposits	Entry into treated crops
Total systemic exposure (mg a.s./day)	<b>0.0170433</b>	<b>0.0138000</b>	<b>0.0049609</b>	<b>0.0226410</b>

Total systemic exposure per kg body weight (mg/kg bw/day)	0.0002841	0.0002300	0.0000827	0.0003773
% of RVAAS	3.16%	2.56%	0.92%	4.19%

2. Details

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
<b>1-3 year old child</b>				
Spray drift	0.0140057	0.0014006	$((C16 * i\_AbsorpInuse * (1 - d\_ClothAF)) + C18) * d\_ConcAS$	
Vapour	0.0107000	0.0010700	$d\_AirCon * d\_BreathRCh * d\_BwChild$	
Surface deposits				
Dermal	0.0017791	0.0001779	$(i\_AppRate / 100) * C24 * d\_Turf * d\_ByTCCh * d\_ByExpDur * MAX(i\_AbsorpProduct, i\_AbsorpInuse) * d\_MAF * IF(i\_AppEquip = "Vehicle-mounted-Drift Reduction", 0.5, 1)$	
Hand to mouth	0.0060376	0.0006038	$(i\_AppRate / 100) * C25 * d\_Turf * d\_SalExt * d\_AreaHM * d\_ByFreqHM * d\_ByExpDur * i\_AbsorpOralInuse * d\_MAF$	
Object to mouth	0.0015094	0.0001509	$(i\_AppRate / 100) * C25 * d\_DRP * d\_MouthGrass * i\_AbsorpOralInuse * d\_MAF$	



Entry into treated crops				
Dermal	<b>0.0067923</b>	<b>0.0006792</b>	$(d\_TcEntryCh*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_AbsorpInuse)$	
Hand to mouth			$(i\_AppRate/100)*d\_MAF*d\_Turf*d\_SalExt*d\_AreaHM*d\_ByFreqHM*d\_ByExpDur*i\_AbsorpOrallnuse$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Object to mouth			$(i\_AppRate/100)*d\_DRP*d\_MouthGrass*i\_AbsorpOrallnuse*d\_MAF$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
<b>Adult</b>				
Spray drift	<b>0.0170433</b>	<b>0.0002841</b>	$((C15*i\_AbsorpInuse*(1-d\_ClothAF)t)+C17)*d\_ConcAS$	
Vapour	<b>0.0138000</b>	<b>0.0002300</b>	$d\_AirCon*d\_BreathRAd*d\_BwAdult$	
Surface deposits (dermal)	<b>0.0049609</b>	<b>0.0000827</b>	$(i\_AppRate/100)*C24*d\_Turf*d\_ByTCA*d\_ByExpDur*MAX(i\_AbsorpProduct,i\_AbsorpInuse)*d\_MAF*IF(i\_AppEquip="Vehicle-mounted-Drift Reduction",0.5,1)$	
Entry into treated crops (dermal)	<b>0.0226410</b>	<b>0.0003773</b>	$(d\_TcEntryAd*0.25*d\_DFR*d\_MAF)/1000*MAX(i\_AbsorpProduct,i\_AbsorpInuse)$	

## Recreational exposure for Allure Vloeibaar

Croptype	Golf course, turf or other sports lawns	<b>This sheet is only to be used for treatment of grassland used for recreational purposes</b>	
Application method	Downward spraying		
Application equipment	Vehicle-mounted		<i>i_AppEquip</i>
Formulation type	Soluble concentrates, emulsifiable concentrate, etc.		<i>i_FormVal</i>
Application rate of the product	0.66 kg a.s./ha		<i>i_AppRate</i>
Dermal absorption of product	0.02%		<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	0.34%		<i>i_AbsorpInuse</i>
Oral absorption	30.00%		<i>i_AbsorpOrallnuse</i>
Dislodgeable foliar residue ( $i\_AppRate \cdot i\_DFR$ )	1.98 µg a.s./cm <sup>2</sup>		<i>d_DFR</i>
Exposure duration dermal	2 hours		<i>d_ReExpDur</i>
Light clothing adjustment factor Adult resident	18.0%		<i>d_ClothAF</i>
Drift percentage on surface	100.00%		
Turf transferable residues percentage	5.00%		<i>d_Turf</i>
Transfer coeff. of surface deposits-adult	7300 cm <sup>2</sup> /hour		<i>d_ReTCAd</i>
Transfer coeff. of surface deposits-child (1-3 year old)	2600 cm <sup>2</sup> /hour		<i>d_ReTCCh</i>
Saliva extraction percentage	50.00%		<i>d_SalExt</i>
Surface area of hands mouthed	20 cm <sup>2</sup>		<i>d_AreaHM</i>
Frequency of hand to mouth activity	9.5 events/hour		<i>d_ReFreqHM</i>
Ingestion rate for mouthing of grass per day	25 cm <sup>2</sup>		<i>d_MouthGrass</i>

## 2. Details

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
<b>1-3 year old child</b>				
Surface deposits				
Dermal	0.0104652	0.0010465	$(i\_AppRate/100) \cdot C13 \cdot d\_Turf \cdot d\_ReTCCh \cdot d\_ReExpDur \cdot \text{MAX}(i\_AbsorpProduct, i\_AbsorpInuse) \cdot d\_MAF$	
Hand to mouth	0.0337395	0.0033740	$(i\_AppRate/100) \cdot C13 \cdot d\_Turf \cdot d\_SalExt \cdot d\_AreaHM \cdot d\_ReFreqHM \cdot d\_ReExpDur \cdot i\_AbsorpOrallnuse \cdot d\_MAF$	
Object to mouth	0.0177576	0.0017758	$(i\_AppRate/100) \cdot C13 \cdot d\_DRP \cdot d\_MouthGrass \cdot i\_AbsorpOrallnuse \cdot d\_MAF$	
Total systemic exposure	0.0619623	0.0061962		
% of RVNAS				
<b>Adult</b>				
Surface deposits (dermal)	0.0293830	0.0004897	$(i\_AppRate/100) \cdot C13 \cdot d\_Turf \cdot d\_ReTCAd \cdot d\_ReExpDur \cdot \text{MAX}(i\_AbsorpProduct, i\_AbsorpInuse) \cdot d\_MAF$	
% of RVNAS				

<i>d_AirConVol</i>	Concentration in air of moderately volatile substances	<b>0.015</b> mg/m <sup>3</sup>
<i>d_AirConNonVol</i>	Concentration in air of low volatile substances	<b>0.001</b> mg/m <sup>3</sup>
<i>d_AreaHM</i>	Surface area of hands mouthed	<b>20</b> cm <sup>2</sup>
<i>d_AreaTreated</i>	Area treated (defined by crop type)	<b>10</b> ha
<i>d_BreathRAd</i>	Breathing rate adult residents	<b>0.23</b> m <sup>3</sup> /day/kg
<i>d_BreathRCh</i>	Breathing rate child (1-3 year old) residents	<b>1.07</b> m <sup>3</sup> /day/kg
<i>d_BwAdult</i>	Adult body weight	<b>60</b> kg
<i>d_BwChild</i>	Child body weight (1 to < 3 year olds)	<b>10</b> kg
<i>d_ByBreathRAd</i>	Breathing rate adult bystander	<b>0.04</b> m <sup>3</sup> /hours/kg
<i>d_ByBreathRCh</i>	Breathing rate child (1-3 year old) bystander	<b>0.19</b> m <sup>3</sup> /hours/kg
<i>d_ByExpDur</i>	Exposure duration intense activity breathing rates	<b>2</b> hours
<i>d_ByFreqHM</i>	Frequency of hand to mouth activity	<b>20</b> events/hour
<i>d_ByTCAd</i>	Transfer coeff. of surface deposits-adult	<b>14500</b> cm <sup>2</sup> /hour
<i>d_ByTCCh</i>	Transfer coeff. of surface deposits-child (1-3 year old)	<b>5200</b> cm <sup>2</sup> /hour
<i>d_ClothAF</i>	Light clothing adjustment factor resident and bystanders	<b>18.0%</b>
<i>d_ConcAs</i>	Concentration of active substance (in-use dilution for liquid applications)	<b>4.4</b> g a.s./l
<i>d_DFR</i>	Dislodgeable foliar residue (i_AppRate*i_DFR)	<b>1.98</b> µg a.s./cm <sup>2</sup>
<i>d_DRP</i>	Dislodgeable residues percentage transferability for object to mouth	<b>20.0%</b>
<i>d_HalfLifeAS</i>	Half-life of active substance (DT50)	<b>30</b> days
<i>d_InhalTcAut</i>	Inhalation transfer coefficient for automated applications	<b>NA</b> ha/hr*10 <sup>^(-3)</sup>
<i>d_InhalTcCut</i>	Inhalation transfer coefficient for cutting ornamentals	<b>NA</b> ha/hr*10 <sup>^(-3)</sup>
<i>d_InhalTcSort</i>	Inhalation transfer coefficient for sorting / bundling ornamentals	<b>NA</b> ha/hr*10 <sup>^(-3)</sup>
<i>d_MAF</i>	Multiple application factor	<b>1.79</b>
<i>d_MouthGrass</i>	Ingestion rate for mouthing of grass per day	<b>25</b> cm <sup>2</sup> grass/day
<i>d_ReExpDur</i>	Exposure duration resident dermal	<b>2</b> hours
<i>d_ReExpDurInhal</i>	Exposure duration resident inhalation	<b>24</b> hours
<i>d_ExpDurTreatCrop</i>	Exposure duration for resident and bystander entry into treated crops	<b>0.25</b> hours
<i>d_ReFreqHM</i>	Frequency of hand to mouth activity	<b>9.5</b> events/hour
<i>d_ReTCAd</i>	Transfer coeff. of surface deposits-adult	<b>7300</b> cm <sup>2</sup> /hour
<i>d_ReTCCh</i>	Transfer coeff. of surface deposits-child (1-3 year old)	<b>2600</b> cm <sup>2</sup> /hour
<i>d_SalExt</i>	Saliva extraction percentage	<b>50.0%</b>
<i>d_TcEntryAd</i>	Transfer coefficient for entry into treated crops 75th percentile - adult	<b>7500</b> cm <sup>2</sup> /h
<i>d_TcEntryCh</i>	Transfer coefficient for entry into treated crops 75th percentile - child	<b>2250</b> cm <sup>2</sup> /h
<i>d_TcEntryMeanAd</i>	Transfer coefficient for entry into treated crops mean - adult	<b>5980</b> cm <sup>2</sup> /h
<i>d_TcEntryMeanCh</i>	Transfer coefficient for entry into treated crops mean - child	<b>1794</b> cm <sup>2</sup> /h
<i>d_Turf</i>	Turf transferable residues percentage	<b>5.0%</b>
<i>d_PctExtrapolation</i>	For exposure value 75 percentiles above this amount linear extrapolation is performed	<b>1.5</b> kg
<i>d_head75ProtectionFactor</i>	Coefficient to estimate head protection factor 75 th Percentile	<b>1.79422</b>
<i>d_head95ProtectionFactor</i>	Coefficient to estimate head protection factor 95 Percentile	<b>1.24705</b>

*sys\_KeyOperator* Variables for operator exposure lookup key  
*sys\_OperatorModel* Operator model

**i\_IndoorOutdoor&i\_FormVal&i\_AppMeth&i\_AppEquip&**

**1**

RPE reduction factor	
key_MixRPE, ay_MixRPE	
None	1
FP1, P1 and similar	0.25
FP2, P2 and similar	0.1

PPE reduction factor	
key_MixPPEBody, ay_MixPPEBody	
Potential exposure	1
Work wear - arms, body and legs covered	0.1
Certified protective coverall	0.05

PPE reduction factor	
key_MixPPEHead, ay_MixPPEHead	
None	1
Hood	0.5
Hood and visor	0.05
FP1, P1 and similar	0.8
FP2, P2 and similar	0.8

Application: Gloves PPE reduction factor (depending on formulation type)	
key_AppPPEHands, ay_AppPPEHands	
Wettable powder, soluble powder	0.05
Chemical resistant gloves	0.05
Granules, fine granules	0.05
Chemical resistant gloves	0.05
Wettable granules, soluble granules	0.1
Chemical resistant gloves	0.1
Soluble concentrates, emulsifiable concentrate, etc.	1
Wettable powder, soluble powder	1
Granules, fine granules	1
Wettable granules, soluble granules	1
Chemical resistant gloves	1
Soluble concentrates, emulsifiable concentrate, etc.	1

Crop dependent exposure parameters									
key_CropType, ay_CropType	Transfer coefficients	Transfer coefficients	1400	Transfer coefficients	Area Treated				
Crop type	Arm, body and legs covered	Total potential exposure	Activity	hours per day	Body parts involved	Hands, arm, body and legs covered	Type of crop for Resident Bystander	Vehicle Mounted Applications	
Bare soil	NA	NA	NA	NA	NA	NA	Field crops	50	
Low berries and other small fruits		3000	5800 Reaching, picking		8 Hand and forearm		750 Field crops	50	
Brassica vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops	50	
Bulb vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops	50	
Cane fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Field crops	10	
Cereals		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops	50	
Citrus fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops	10	
Fruiting vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops	50	
Grapes		10100	30000 Hand harvesting		8 Hand and body	no TC available for this assessment	Grapes	10	
Grassland and lawns		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops	50	
Golf course, turf or other sports lawns		2500	5800 Maintenance		8 Hand and body		580 Field crops	50	
Hops		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Hops	50	
Leaf vegetables and fresh herbs		2500	5800 Reaching, picking		8 Hand and body		580 Field crops	10	
Legume vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops	50	
Oilfruits		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops	10	
Oilseeds		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops	50	
Ornamentals		5000	14000 Cutting, sorting, bundling, carrying		8 Hand and body		1400 Field crops	10	
Pome fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops	10	
Root and tuber vegetables		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops	50	
Stone fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops	10	
Tree nuts		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops	10	

Resident Spray Drift					
These values are the 75th Percentiles for Residents (assuming average breathing rates for inhalation exposures)					
key_ResidSpray, ay_ResidSpray	Adults Dermal	Children Dermal	Adults Inhalation	Children Inhalation	
Downward spraying2-3		0.47	0.327	0.0001	0.00022
Downward spraying5		0.24	0.22	0.00009	0.00017
Downward spraying10		0.20	0.18	0.00009	0.00013
Upward spraying2-3	NA	NA	NA	NA	
Upward spraying5		5.63	1.689	0.0021	0.00164
Upward spraying10		5.63	1.689	0.0021	0.00164

Bystander Spray Drift					
These values are the 95th Percentiles for Bystanders (assuming high breathing rates for inhalation exposures)					
key_BySpray, ay_BySpray	Adults Dermal	Children Dermal	Adults Inhalation	Children Inhalation	
Downward spraying2-3		1.21	0.74	0.0005	0.0011
Downward spraying5		0.57	0.48	0.00048	0.0008
Downward spraying10		0.48	0.39	0.00051	0.00076
Upward spraying2-3	NA	NA	NA	NA	
Upward spraying5		12.9	3.87	0.0044	0.0035
Upward spraying10		12.9	3.87	0.0044	0.0035

Mean Spray Drift					
These values are the mean values (assuming average breathing rates for inhalation exposures)					
key_AvgSpray, ay_AvgSpray	Adults Dermal	Children Dermal	Adults Inhalation	Children Inhalation	
Downward spraying2-3		0.22	0.18	0.0001	0.0002
Downward spraying5		0.12	0.12	0.0001	0.0001
Downward spraying10		0.11	0.1	0.0001	0.0001
Upward spraying2-3	NA	NA	NA	NA	
Upward spraying5		3.68	1.11	0.0017	0.0013
Upward spraying10		3.68	1.11	0.0017	0.0013

Resident and bystander Surface Deposits Drift percentage			
Ground sediments in % of the application rate calculated on the basis of percentile values (drift data acc. Rautmann)			
key_ByCropType, ay_ByCropType	Bystander surface deposit (90th Percentile)	Resident surface deposit (77th Percentile)	mean
Field cropsnot relevant2-3	0.085	0.056	0.041
Field cropsnot relevant5	0.035	0.023	0.018
Field cropsnot relevant10	0.019	0.013	0.010
Fruit cropsnot relevant2-3	0.292	0.240	0.190
Fruit cropsnot relevant5	0.199	0.158	0.117
Fruit cropsnot relevant10	0.118	0.090	0.061
Fruit cropsearly (without leaves)2-3	0.292	0.240	0.190
Fruit cropsearly (without leaves)5	0.199	0.158	0.117

Fruit cropsearly (without leaves)10	0.118	0.090	0.061
Fruit cropslate (dense foliage)2-3	0.157	0.110	0.070
Fruit cropslate (dense foliage)5	0.084	0.060	0.037
Fruit cropslate (dense foliage)10	0.036	0.027	0.016
Grapesnot relevant2-3	0.080	0.069	0.053
Grapesnot relevant5	0.036	0.031	0.023
Grapesnot relevant10	0.012	0.010	0.008
Hopsnot relevant2-3	0.193	0.159	0.100
Hopsnot relevant5	0.116	0.086	0.059
Hopsnot relevant10	0.058	0.037	0.029

	1	2	3	4	5	6	7	8	9	10	11	12	13	14
<b>Match Method</b>														
IndoorGranules, fine granulesApplication of granulesManualBody	Outdoor/Indoor	Formulation type	Application method	Application equipment	Type of exposure	Mixing & Loading 75th percentile	Mixing & Loading 95th percentile	Mixing & Loading Comments	Mixing & Loading Model	Application 75th percentile	Application 95th percentile	Application Comments	Application Model	
IndoorGranules, fine granulesApplication of granulesManualHands	Indoor	Granules, fine granules	Application of granules	Manual	Body			Value for application is for combination of mixing&loading and application	PHED	68.8708	253.4438	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	
IndoorGranules, fine granulesApplication of granulesManualInhalation	Indoor	Granules, fine granules	Application of granules	Manual	Hands			Value for application is for combination of mixing&loading and application	PHED	28.5320	94.3636	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	
OutdoorGranules, fine granulesBroadcast application of granulesVehicle-mountedBody	Outdoor	Granules, fine granules	Application of granules	Manual	Inhalation			Value for application is for combination of mixing&loading and application	PHED	0.4677	1.5251	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	
OutdoorGranules, fine granulesBroadcast application of granulesVehicle-mountedHands	Outdoor	Granules, fine granules	Broadcast application of granules	Vehicle-mounted	Body	0.0162	0.0427	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0.0047	0.0151	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	
OutdoorGranules, fine granulesBroadcast application of granulesVehicle-mountedInhalation	Outdoor	Granules, fine granules	Broadcast application of granules	Vehicle-mounted	Hands	0.0015	0.0069	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0.0004	0.0018	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	
OutdoorGranules, fine granulesIn furrow application of granulesVehicle-mountedBody	Outdoor	Granules, fine granules	Broadcast application of granules	Vehicle-mounted	Inhalation	0.0208	0.0784		PHED	0.0012	0.0045		PHED	
OutdoorGranules, fine granulesIn furrow application of granulesVehicle-mountedHands	Outdoor	Granules, fine granules	In furrow application of granules	Vehicle-mounted	Body	0.0162	0.0427	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0.0047	0.0151	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	
OutdoorGranules, fine granulesIn furrow application of granulesVehicle-mountedInhalation	Outdoor	Granules, fine granules	In furrow application of granules	Vehicle-mounted	Hands	0.0015	0.0069	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0.0004	0.0018	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	
OutdoorGranules, fine granulesManual application of granulesManualBody	Outdoor	Granules, fine granules	Manual application of granules	Manual	Body			Value for application is for combination of mixing&loading and application	PHED	68.8708	253.4438	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	
OutdoorGranules, fine granulesManual application of granulesManualHands	Outdoor	Granules, fine granules	Manual application of granules	Manual	Hands			Value for application is for combination of mixing&loading and application	PHED	28.5320	94.3636	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	
OutdoorGranules, fine granulesManual application of granulesManualInhalation	Outdoor	Granules, fine granules	Manual application of granules	Manual	Inhalation			Value for application is for combination of mixing&loading and application	PHED	0.4677	1.5251	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	





