

Instructions for using the workbook

The information required for the exposure assessment needs to be entered in the worksheet "**Data entry**".

In the following worksheets formulas calculate the exposure values automatically

Worksheet "**Operator Outdoor Spray AOEM**" is to be for outdoor spray applications. PPE options can be selected in this worksheet

Worksheet "**Operator Granules**" is for granular applications. Currently the calculator does not allow operator exposure for indoor applications. PPE options can be selected in this worksheet

Worksheets "**Resident exposure**" and "**Bystander exposure**" are only relevant for outdoor applications

Worksheet "**Recreational Exposure**" is only applicable for golf course, turf, other sports lawns or amenity turf/grassland areas where members of the public are likely to have access

The combined results of the exposure assessment are presented in worksheet "**Summary**"

This calculator should be used in conjunction with the **Guidance on the Assessment of Exposure for Operators, Workers, Residents and Bystanders in Risk Assessment for Plant Protection Products**

Latest version: 23 Oct 2014 - Version produced to support guidance document published 23/10/2014

Note: Some drop-down menus depend on others. To avoid errors, please fill-in from top to bottom

Substance name	thiacloprid
Product name	Calypso
Reference value non acutely toxic active substance (RVNAS)	0,02 mg/kg bw/day
Reference value acutely toxic active substance (RVAAS)	0,02 mg/kg bw/day
Crop type	Pome fruit
Substance properties	
Formulation type	Soluble concentrates, emulsifiable concentrate, etc.
Minimum volume water for application (liquids)	500 L/ha
Maximum application rate of active substance	0,18 kg a.s. /ha
50% Dissipation Time DT50	30 days
Initial Dislodgeable Foliar Residue	3 µg/cm ² of foliage/kg a.s. applied/ha
Dermal absorption of product	1,00%
Dermal absorption of in-use dilution	10,00%
Oral absorption of active substance	95,00%
Inhalation absorption of active substance	100,00%
Vapour pressure of active substance	low volatile substances having a vapour pressure of $5 \cdot 10^{-3}$ Pa
Scenario	
Indoor or Outdoor application	Outdoor
Application method	Upward spraying
Application equipment	Manual-Hand held
Buffer strip	5 m
Number of applications	2
Interval between multiple applications	60 days
Season (upward spraying orchards only)	early (without leaves)

Exposure assessment

Substance	thiacloprid	Formulation = Soluble concentrates, emulsifiable concentrate, etc.	Application rate-0,18 kg a.s. /ha	Spray dilution = 0,36 g a.s./l	Vapour pressure = low volatile substances having a vapour pressure of $5 \cdot 10^{-3}$Pa
Scenario	Pome fruit early (without leaves) / Outdoor / Upward spraying / Manual-Hand held			Buffer = 5	Number applications = 2, Application interval = 60 days
Percentage Absorption	Dermal for product = 1	Dermal for in use dilution = 10	Oral = 95	Inhalation = 100	
RVNAS	0,02 mg/kg bw/day		RVAAS	0,02 mg/kg bw/day	
DFR	3 µg a.s./cm ² per kg a.s./ha		DT50	30 days	

Operator Model	Mixing, loading and application AOEM			
Potential exposure	Longer term systemic exposure mg/kg bw/day	0,1032	% of RVNAS	515,82%
	Acute systemic exposure mg/kg bw/day	0,3259	% of RVAAS	1629,27%
Mixing and Loading	Gloves = No	Clothing = Work wear - arms, body and legs covered	RPE = None	Soluble bags = No
Application	Gloves = No	Clothing = Work wear - arms, body and legs covered	RPE = None	Closed cabin = No
Exposure (including PPE options above)	Longer term systemic exposure mg/kg bw/day	0,0071	% of RVNAS	35,56%
	Acute systemic exposure mg/kg bw/day	0,0212	% of RVAAS	105,96%

Worker - Searching, reaching, picking	Potential exposure mg/kg bw/day	0,2025	% of RVNAS	1012,50%
	Working clothing mg/kg bw/day	0,0405	% of RVNAS	202,50%
	Working clothing and gloves mg/kg bw/day	0,0203	% of RVNAS	101,25%

Resident - child	Spray drift (75th percentile) mg/kg bw/day	#NA	% of RVNAS	#NA
	Vapour (75th percentile) mg/kg bw/day	#NA	% of RVNAS	#NA
	Surface deposits (75th percentile) mg/kg bw/day	#NA	% of RVNAS	#NA
	Entry into treated crops (75th percentile) mg/kg bw/day	#NA	% of RVNAS	#NA
	All pathways (mean) mg/kg bw/day	#NA	% of RVNAS	#NA
Resident - adult	Spray drift (75th percentile) mg/kg bw/day	#NA	% of RVNAS	#NA
	Vapour (75th percentile) mg/kg bw/day	#NA	% of RVNAS	#NA
	Surface deposits (75th percentile) mg/kg bw/day	#NA	% of RVNAS	#NA
	Entry into treated crops (75th percentile) mg/kg bw/day	#NA	% of RVNAS	#NA
	All pathways (mean) mg/kg bw/day	#NA	% of RVNAS	#NA

Bystander - child	Spray drift (95th percentile) mg/kg bw/day	#NA	% of RVAAS	#NA
	Vapour (95th percentile) mg/kg bw/day	#NA	% of RVAAS	#NA
	Surface deposits (95th percentile) mg/kg bw/day	#NA	% of RVAAS	#NA

Exposure assessment

	Entry into treated crops (95th percentile) mg/kg bw/day	#NA	% of RVAAS	#NA
Bystander - adult	Spray drift (95th percentile) mg/kg bw/day	#NA	% of RVAAS	#NA
	Vapour (95th percentile) mg/kg bw/day	#NA	% of RVAAS	#NA
	Surface deposits (95th percentile) mg/kg bw/day	#NA	% of RVAAS	#NA
	Entry into treated crops (95th percentile) mg/kg bw/day	#NA	% of RVAAS	#NA
Recreational Exposure		Child % of RVNAS	Adult % of RVNAS	

Latest version: 23 Oct 2014 - Version produced to support guidance document published 23/10/2014

Operator exposure for Calypso outdoor spray applications

Application rate of active substance	0,18 kg a.s./ha	<i>i_AppRate</i>
Assumed area treated	4 ha/day	<i>d_AreaTreated</i>
Amount of active substance applied	0,72 kg a.s./day	<i>i_AmountAS</i>
Dermal absorption of the product	1,00%	<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	10,00%	<i>i_AbsorInuse</i>
Formulation type	Soluble concentrates, emulsifiable concentrate, etc.	
Indoor or Outdoor application	Outdoor	
Application method	Upward spraying	
Application equipment	Manual-Hand held	
Season	early (without leaves)	

Mixing and loading	Exposure values	µg exposure/day mixed and loaded		Reference	Comment
		75 th centile	95 th centile		
		Hands	3772		
Body	2832	65468	AOEM		
Head	37	205	AOEM		
Protected hands (gloves)	28	143	AOEM		
Protected body (workwear or protective garment and sturdy footwear)	18	105	AOEM		
Protected head (hood and face shield)	1	12	AOEM		
Inhalation	3	28	AOEM		
Protective Equipment	Select for inclusion		Penetration factor	Inhalation Protection factor	
Gloves	No				
Clothing	Work wear - arms, body and legs covered		Incl. in AOEM model		
Head and respiratory PPE	None		1	1	
Water soluble bag	No		1		

Application	Exposure values	µg exposure/day applied		Reference	Comment
		75 th centile	95 th centile		
		Hands	2067		
Body	58382	178199	AOEM		
Head	137	738	AOEM		
Protected hands (gloves)	17	89	AOEM		
Protected body (workwear or protective garment and sturdy footwear)	1033	1938	AOEM		
Inhalation	61	150	AOEM		
Protective Equipment	Select for inclusion		Penetration factor	Inhalation Protection factor	
Gloves	No				
Clothing	Work wear - arms, body and legs covered		Incl. in AOEM model		
Head and respiratory PPE	None		1	1	
Closed cab	No		vehicle mounted upward spraying only		

1. Total	Without RPE/PPE	With RPE/PPE
Longer term		
Total systemic exposure from mixing, loading and application (mg a.s./day)	6,1897963	0,4267545
Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	0,1031633	0,0071126
% of RVNAS	515,82%	35,56%
Acute		

Total systemic exposure from mixing, loading and application (mg a.s./day)	19,551,892	1,271,504	
--	------------	-----------	--

Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	0,3258532	0,0211917	
% of RVAAS	1629,27%	105,96%	

2. Longer term exposure

2.1 Mixing and loading

	Systemic exposure [$\mu\text{g a.s. /day}$]	Systemic exposure [$\mu\text{g a.s./kg bw/day}$]	Formula
Without RPE/PPE			
Hands	37,7164037	0,6286067	D15*i_AbsorpProduct
Body	28,3168364	0,4719473	D16*i_AbsorpProduct
Head	0,3735618	0,0062260	D17*i_AbsorpProduct
Inhalation	3,3569290	0,0559488	D21*i_AbsorpInhalation
Sum	69,7637309	1,1627288	
With RPE/PPE (as selected above)			
Hands	37,7164037	0,6286067	D18*i_AbsorpProduct
Body	0,1776372	0,0029606	D19*i_AbsorpProduct or D15*i_AbsorpProduct*F24
Head	0,3735618	0,0062260	D20*i_AbsorpProduct or D17*i_AbsorpProduct*F25
Inhalation	3,3569290	0,0559488	D21*i_AbsorpInhalation*G25
Sum	41,6245317	0,6937422	
Water soluble bag	41,6245317	0,6937422	C70*F26

2.2 Application

	Systemic exposure [$\mu\text{g a.s. /day}$]	Systemic exposure [$\mu\text{g a.s./kg bw/day}$]	Formula
Without RPE/PPE			
Hands	206,7242604	3,4454043	D30*i_Absorpnuse
Body	5838,2095949	97,3034932	D31*i_Absorpnuse
Head	13,7056458	0,2284274	D32*i_Absorpnuse
Inhalation	61,3930182	1,0232170	D35*i_AbsorpInhalation
Sum	6120,0325193	102,0005420	
With RPE/PPE (as selected above)			
Hands	206,7242604	3,4454043	D33*i_Absorpnuse
Body	103,3070595	1,7217843	D34*i_Absorpnuse or D31*i_Absorpnuse*F38
Head	13,7056458	0,2284274	D32*i_Absorpnuse*F39
Inhalation	61,3930182	1,0232170	D35*i_Absorpnuse*G39
Sum	385,1299839	6,4188331	

3. Acute exposure

3.1 Mixing and loading

	Systemic exposure [$\mu\text{g a.s. /day}$]	Systemic exposure [$\mu\text{g a.s./kg bw/day}$]	Formula
Without RPE/PPE			
Hands	137,7972409	2,2966207	E15*i_AbsorpProduct
Body	654,6847683	10,9114128	E16*i_AbsorpProduct
Head	2,0488138	0,0341469	E17*i_AbsorpProduct
Inhalation	28,4912603	0,4748543	E21*i_AbsorpInhalation
Sum	823,0220832	13,7170347	
With RPE/PPE (as selected above)			
Hands	137,7972409	2,2966207	E18*i_AbsorpProduct
Body	1,0530100	0,0175502	E19*i_AbsorpProduct or E16*i_AbsorpProduct*F24
Head	2,0488138	0,0341469	E20*i_AbsorpProduct or E17*i_AbsorpProduct*F25
Inhalation	28,4912603	0,4748543	E21*i_AbsorpInhalation*G25
Sum	169,3903250	2,8231721	
Water soluble bag	169,3903250	2,8231721	C104*F26

2.2 Application

	Systemic exposure [$\mu\text{g a.s. /day}$]	Systemic exposure [$\mu\text{g a.s./kg bw/day}$]	Formula
Without RPE/PPE			
Hands	684,4417711	11,4073629	E30*i_Absorpnuse
Body	17819,8741332	296,9979022	E31*i_Absorpnuse
Head	73,8437085	1,2307285	E32*i_Absorpnuse
Inhalation	150,0075257	2,5001254	E35*i_AbsorpInhalation

Sum	18728,1671385	312,1361190	
With RPE/PPE (as selected above)			
Hands	684,4417711	11,4073629	<i>E33*i_Absorplnuse</i>
Body	193,8206296	3,2303438	<i>E34*i_Absorplnuse or E31*i_Absorplnuse*F38</i>
Head	73,8437085	1,2307285	<i>E32*i_Absorplnuse*F39</i>
Inhalation	150,0075257	2,5001254	<i>E35*i_Absorplnhalation*G39</i>
Sum	1102,1136350	18,3685606	

Operator exposure for Calypso granular applications

Application rate of active substance	0,18 kg a.s./ha	<i>i_AppRate</i>
Assumed area treated	4 ha/day	<i>d_AreaTreated</i>
Amount of active substance applied	0,72 kg a.s./day	<i>i_AmountAS</i>
Dermal absorption of the product	1,00%	<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	10,00%	<i>i_AbsorInuse</i>
Formulation type	Soluble concentrates, emulsifiable concentrate, etc.	
Indoor or Outdoor application	Outdoor This sheet is only to be used for granular applications	
Application method	Upward spraying	
Application equipment	Manual-Hand held	

	Exposure values	mg exposure/kg a.s. mixed and loaded		Reference	Comment
		75 th centile	95 th centile		
Mixing and loading	Hands	#N/A	#N/A	#N/A	#N/A
	Body	#N/A	#N/A	#N/A	#N/A
	Inhalation	#N/A	#N/A	#N/A	#N/A
	Protective Equipment	Choose item		Penetration factor	
	Gloves	Chemical resistant gloves			Protection for granules exposure is based on measured values
	Body PPE	Certified protective coverall			
	RPE	None		1	

	Exposure values	mg exposure/kg a.s. applied		Reference	Comment
		75 th centile	95 th centile		
Application	Hands	#N/A	#N/A	#N/A	#N/A
	Body	#N/A	#N/A	#N/A	#N/A
	Inhalation	#N/A	#N/A	#N/A	#N/A
	Protective Equipment	Choose item		Penetration factor	
	Gloves	Chemical resistant gloves			Protection for granules exposure is based on measured values
	Body PPE	Certified protective coverall			
	RPE	FP1, P1 and similar		0,25	

1. Total			
	Without RPE/PPE	With RPE/PPE	
Longer term			
Total systemic exposure from mixing, loading and application (mg a.s./day)	#N/A	#N/A	
Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	#N/A	#N/A	
% of RVNAS	#N/A	#N/A	
Acute			

Total systemic exposure from mixing, loading and application (mg a.s./day)	#N/A	#N/A	
--	------	------	--

Total systemic exposure from mixing, loading and application per kg body weight (mg/kg bw/day)	#N/A	#N/A
% of RVAAS	#N/A	#N/A

2. Longer term exposure

2.1 Mixing and loading

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
Without RPE/PPE			
Hands	#N/A	#N/A	$D14*100*i_AmoutAS*i_AbsorpProduct$
Body	#N/A	#N/A	$D15*100*i_AmoutAS*i_AbsorpProduct$
Inhalation	#N/A	#N/A	$D16*i_AmoutAS*i_Absorplnhalation$
Sum	#N/A	#N/A	
With RPE/PPE (as selected above)			
Hands	#N/A	#N/A	$D14*i_AmoutAS*i_AbsorpProduct$
Body	#N/A	#N/A	$D15*i_AmoutAS*i_AbsorpProduct$
Inhalation	#N/A	#N/A	$D16*i_AmoutAS*i_Absorplnhalation*F20$
Sum	#N/A	#N/A	

2.2 Application

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
Without RPE/PPE			
Hands	#N/A	#N/A	$D25*100*i_AmoutAS*i_Absorplnuse$
Body	#N/A	#N/A	$D26*100*i_AmoutAS*i_Absorplnuse$
Inhalation	#N/A	#N/A	$D27*i_AmoutAS*i_Absorplnhalation$
Sum	#N/A	#N/A	
With RPE/PPE (as selected above)			
Hands	#N/A	#N/A	$D25*i_AmoutAS*i_Absorplnuse$
Body	#N/A	#N/A	$D26*i_AmoutAS*i_Absorplnuse$
Inhalation	#N/A	#N/A	$D27*i_AmoutAS*i_Absorplnhalation*F31$
Sum	#N/A	#N/A	

3. Acute exposure

3.1 Mixing and loading

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
Without RPE/PPE			
Hands	#N/A	#N/A	$E14*100*i_AmoutAS*i_AbsorpProduct$
Body	#N/A	#N/A	$E15*100*i_AmoutAS*i_AbsorpProduct$
Inhalation	#N/A	#N/A	$E16*i_AmoutAS*i_Absorplnhalation$
Sum	#N/A	#N/A	
With RPE/PPE (as selected above)			
Hands	#N/A	#N/A	$E14*100*i_AmoutAS*i_AbsorpProduct$
Body	#N/A	#N/A	$E15*100*i_AmoutAS*i_AbsorpProduct$
Inhalation	#N/A	#N/A	$E16*i_AmoutAS*i_Absorplnhalation*F20$
Sum	#N/A	#N/A	

3.2 Application

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula
Without RPE/PPE			
Hands	#N/A	#N/A	$E25*100*i_AmoutAS*i_Absorplnuse$
Body	#N/A	#N/A	$E25*100*i_AmoutAS*i_Absorplnuse$

Inhalation	#N/A	#N/A	$E26 * i_AmountAS * i_Absorplnhalation$
Sum	#N/A	#N/A	
With RPE/PPE (as selected above)			
Hands	#N/A	#N/A	$E25 * 100 * i_AmountAS * i_Absorplnuse$
Body	#N/A	#N/A	$E26 * 100 * i_AmountAS * i_Absorplnuse$
Inhalation	#N/A	#N/A	$E27 * i_AmountAS * i_Absorplnhalation * F31$
Sum	#N/A	#N/A	

Worker exposure from residues on foliage for Calypso

Crop type	Pome fruit	
Indoor or outdoor	Outdoor	
Application method	Upward spraying	
Application equipment	Manual-Hand held	
Worker's task	Searching, reaching, picking	
Main body parts in contact with foliage	Hand and body	
Application rate of active substance	0,18 kg a.s./ha	<i>i_AppRate</i>
Number of applications	2	<i>i_AppNo</i>
Interval between multiple applications	60 days	<i>i_AppInt</i>
Half-life of active substance	30 days	<i>d_HalfLifeAS</i>
Multiple application factor	1,3	<i>d_MAF</i>
Dermal absorption of the product	1,00%	<i>i_AbsorpProduct</i>
Dermal absorption of the in-use dilution	10,00%	<i>i_Absorplnuse</i>
Dislodgeable foliar residue (<i>i_AppRate</i> * <i>i_DFR</i>)	0,54 µg a.s./cm ²	<i>d_DFR</i>
Working hours	8 hr	<i>d_WorkHr</i>
Dermal transfer coefficient - Total potential exposure	22500 cm ² /hr	<i>d_DermTcUCV</i>
Dermal transfer coefficient - arms, body and legs covered	4500 cm ² /hr	<i>d_DermTcCV1</i>
Dermal transfer coefficient - hands, arms, body and legs covered	2250 cm ² /hr	<i>d_DermTcCV2</i>
Inhalation transfer coefficient for automated applications	NA ha/hr*10 ⁻³	<i>d_InhalTcAut</i>
Inhalation transfer coefficient for cutting ornamentals	NA ha/hr*10 ⁻³	<i>d_InhalTcCut</i>
Inhalation transfer coefficient for sorting / bundling ornamentals	NA ha/hr*10 ⁻³	<i>d_InhalTcSort</i>

1. Total

	Potential exposure	Work wear - arms, body and legs covered	Working wear and gloves	Comments
Total systemic exposure (mg a.s./day)	12,1500000	2,4300000	1,2150000	
Total systemic exposure per kg body weight (mg/kg bw/day)	0,2025000	0,0405000	0,0202500	
% of RVNAS	1012,50%	202,50%	101,25%	

2. Details

	Systemic exposure		Formula	Comments
	[mg a.s. /day]	[mg a.s./kg bw/day]		
Dermal - Potential	12,1500000	0,2025000	$d_DermTcUCV * d_WorkHr * i_DFR * i_MAF / 1000 * i_Absorplnuse$	
Dermal - Work wear - arms, body and legs covered	2,4300000	0,0405000	$d_DermTcCV1 * d_WorkHr * d_DFR * d_MAF / 1000 * i_Absorplnuse$	
Dermal - Working wear and gloves	1,2150000	0,0202500	$d_DermTcCV2 * d_WorkHr * d_DFR * d_MAF / 1000 * i_Absorplnuse$	
Inhalation				Na for outdoor activities

Resident exposure for Calypso

Croptype	Pome fruit	
Application method	Upward spraying	
Application equipment	Manual-Hand held	<i>i_AppEquip</i>
Formulation type	Soluble concentrates, emulsifiable concentrate, etc.	<i>i_FormVal</i>
Buffer strip	5 m	<i>i_Buffer</i>
Application rate of the product	0,18 kg a.s./ha	<i>i_AppRate</i>
Concentration of active substance (in-use dilution for liquid applications)	0,36 g a.s./l	<i>d_ConcAS</i>
Dermal absorption of product	1,00%	<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	10,00%	<i>i_AbsorpInuse</i>
Oral absorption	100,00%	<i>i_AbsorpOrallnuse</i>
Dislodgeable foliar residue ($i_AppRate * i_DFR$)	0,54 $\mu\text{g a.s./cm}^2$	<i>d_DFR</i>
Vapour pressure of in-use dilution	low volatile substances having a vapour pressure of $<5 * 10^{-3} \text{Pa}$	<i>i_Volat</i>
Concentration in air	0,001 mg/m^3	<i>d_AirCon</i>
Resident dermal spray drift exposure 75th percentile - adult	5,63 ml spray dilution/person	
Resident dermal spray drift exposure 75th percentile - child	1,689 ml spray dilution/person	
Resident inhal. spray drift exposure 75th percentile - adult	0,00210 ml spray dilution/person	
Resident inhal. spray drift exposure 75th percentile - child	0,00164 ml spray dilution/person	
Resident dermal spray drift exposure mean - adult	3,68 ml spray dilution/person	
Resident dermal spray drift exposure mean - child	1,11 ml spray dilution/person	
Resident inhal. spray drift exposure mean - adult	0,00170 ml spray dilution/person	
Resident inhal. spray drift exposure mean - child	0,00133 ml spray dilution/person	
Exposure duration dermal	2 hours	<i>d_ReExpDur</i>
Exposure duration inhalation	24 hours	<i>d_ReExpDurInhal</i>
Exposure duration entry into treated crops	0,25 hours	<i>d_ExpDurTreatCrop</i>
Light clothing adjustment factor	18,0%	<i>d_ClothAF</i>
Breathing rate adult	0,23 $\text{m}^3/\text{day}/\text{kg}$	<i>d_BreathRAD</i>
Breathing rate child (1-3 year old)	1,07 $\text{m}^3/\text{day}/\text{kg}$	<i>d_BreathRCh</i>
Drift percentage on surface (75th percentile)	15,79%	
Drift percentage on surface (mean)	11,69%	
Turf transferable residues percentage	5,00%	<i>d_Turf</i>
Transfer coeff. of surface deposits-adult	7300 cm^2/hour	<i>d_ReTCAd</i>
Transfer coeff. of surface deposits-child (1-3 year old)	2600 cm^2/hour	<i>d_ReTCCh</i>
Saliva extraction percentage	50,00%	<i>d_SalExt</i>
Surface area of hands mouthed	20 cm^2	<i>d_AreaHM</i>
Frequency of hand to mouth activity	9,5 events/hour	<i>d_ReFreqHM</i>
Ingestion rate for mouthing of grass per day	25 cm^2	<i>d_MouthGrass</i>
Dislodgeable residues percentage transferability for object to mouth	20,00%	<i>d_DRP</i>
Transfer coefficient for entry into treated crops (75th percentile) - adult	7500 cm^2/h	<i>d_TcEntryAd</i>
Transfer coefficient for entry into treated crops (75th percentile) - child	2250 cm^2/h	<i>d_TcEntryCh</i>
Transfer coefficient for entry into treated crops (mean) - adult	5980 cm^2/h	<i>d_TcEntryAd</i>
Transfer coefficient for entry into treated crops (mean) - child	1794 cm^2/h	<i>d_TcEntryCh</i>

1. Total

1.1 1-3 year old child

	Spray drift (75th percentile)	Vapour (75th percentile)	Surface deposits (75th percentile)	Entry into treated crops (75th percentile)	All pathways (mean)
Total systemic exposure (mg a.s./day)	0,0504509	0,0107000	0,0141311	0,0379688	0,0842028
Total systemic exposure per kg body weight (mg/kg bw/day)	0,0050451	0,0010700	0,0014131	0,0037969	0,0084203
% of RVNAS	25,23%	5,35%	7,07%	18,98%	42,10%

1.2 Adult

	Spray drift	Vapour	Surface deposits	Entry into treated crops	All pathways (mean)
Total systemic exposure (mg a.s./day)	0,1669536	0,0138000	0,0259351	0,1265625	0,2431589
Total systemic exposure per kg body weight (mg/kg bw/day)	0,0027826	0,0002300	0,0004323	0,0021094	0,0040526
% of RVNAS	13,91%	1,15%	2,16%	10,55%	20,26%

2. Resident exposure 75th Percentile

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
1-3 year old child				
Spray drift	0,0504509	0,0050451	$((C16 * i_AbsorpInuse * (1 - d_ClothAF)) + C18) * d_ConcAS$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	0,0107000	0,0010700	$d_AirCon * d_BreathRCh * d_BwChild$	
Surface deposits				
Dermal	0,0092372	0,0009237	$(i_AppRate/100) * C29 * d_Turf * d_ReTCCh * d_ReExpDur * MAX(i_AbsorpProduct, i_AbsorpInuse) * d_MAF * IF(i_AppEquip = "Vehicle-mounted-Drift Reduction", 0.5, 1)$	
Hand to mouth	0,0032064	0,0003206	$(i_AppRate/100) * C29 * d_Turf * d_SalExt * d_AreaHM * d_ReFreqHM * d_ReExpDur * i_AbsorpOrallnuse * d_MAF$	
Object to mouth	0,0016876	0,0001688	$(i_AppRate/100) * C29 * d_DRP * d_MouthGrass * i_AbsorpOrallnuse * d_MAF$	

Entry into treated crops				
Dermal	0,0379688	0,0037969	$(d_TcEntryCh*0.25*d_DFR*d_MAF)/1000*MAX(i_AbsorpProduct,i_Absorplnuse)$	
Hand to mouth			$(i_AppRate/100)*d_Turf*d_MAF*d_SalExt*d_AreaHM*d_ReFreqHM*d_ReExpDur*i_AbsorpOrallnuse$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Object to mouth			$(i_AppRate/100)*d_DRP*d_MouthGrass*i_AbsorpOrallnuse*d_MAF$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Adult				
Spray drift	0,1669536	0,0027826	$(C15*i_Absorplnuse*(1-d_ClothAF))+C17)*d_ConcAS$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	0,0138000	0,0002300	$d_AirCon*d_BreathRAD*d_BwAdult$	
Surface deposits (dermal)	0,0259351	0,0004323	$(i_AppRate/100)*C30*d_Turf*d_ReTCAd*d_ReExpDur*i_Absorplnuse$	
Entry into treated crops (dermal)	0,1265625	0,0021094	$(d_TcEntryAd*0.25*d_DFR*d_MAF)/1000*MAX(i_AbsorpProduct,i_Absorplnuse)$	

3. Summing of exposure pathways mean

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
1-3 year old child				
Spray drift	0,0327672	0,0032767	$((C20*i_Absorplnuse*(1-d_ClothAF))+C22)*d_ConcAS$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	0,0107000	0,0010700	$d_AirCon*d_BreathRCh*d_BwChild$	
Surface deposits				
Dermal	0,0068387	0,0006839	$(i_AppRate/100)*C30*d_Turf*d_ReTCCh*d_ReExpDur*MAX(i_AbsorpProduct,i_Absorplnuse)*d_MAF*IF(i_AppEquip = "Vehicle-mounted-Drift Reduction",0.5,1)$	
Hand to mouth	0,0023738	0,0002374	$(i_AppRate/100)*C30*d_Turf*d_SalExt*d_AreaHM*d_ReFreqHM*d_ReExpDur*i_AbsorpOrallnuse*d_MAF$	
Object to mouth	0,0012494	0,0001249	$(i_AppRate/100)*C30*d_DRP*d_MouthGrass*i_AbsorpOrallnuse*d_MAF$	
Entry into treated crops				
Dermal	0,0302738	0,0030274	$(d_TcEntryMeanCh*0.25*d_DFR*d_MAF)/1000*MAX(i_AbsorpProduct,i_Absorplnuse)$	
Hand to mouth			$(i_AppRate/100)*1*d_Turf*d_MAF*d_SalExt*d_AreaHM*d_ReFreqHM*d_ReExpDur*i_AbsorpOrallnuse$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Object to mouth			$(i_AppRate/100)*1*d_DRP*d_MouthGrass*i_AbsorpOrallnuse*d_MAF$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Adult				
Spray drift	0,1092456	0,0018208	$"(C19*i_Absorplnuse*(1-d_ClothAF))+C21)*d_ConcAS"$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	0,0138000	0,0002300	$d_AirCon*d_BreathRAD*d_BwAdult$	
Surface deposits (dermal)	0,0192008	0,0003200	$(i_AppRate/100)*C30*d_Turf*d_ReTCAd*d_ReExpDur*MAX(i_AbsorpProduct,i_Absorplnuse)*d_MAF*IF(i_AppEquip = "Vehicle-mounted-Drift Reduction",0.5,1)$	
Entry into treated crops (dermal)	0,1009125	0,0016819	$(d_TcEntryMeanAd*0.25*d_DFR*d_MAF)/1000*MAX(i_AbsorpProduct,i_Absorplnuse)$	

Bystander exposure for Calypso

Croptype	Pome fruit	
Application method	Upward spraying	
Application equipment	Manual-Hand held	<i>i_AppEquip</i>
Formulation type	Soluble concentrates, emulsifiable concentrate, etc.	
Application rate of the product	0,18 kg a.s./ha	<i>i_AppRate</i>
Buffer strip	5 m	<i>i_Buffer</i>
Concentration of active substance (in-use dilution for liquid applications)	0,36 g a.s./l	<i>d_ConcAS</i>
Dermal absorption of product	1,00%	<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	10,00%	<i>i_Absorpnuse</i>
Oral absorption	100,00%	<i>i_AbsorpOrallnuse</i>
Dislodgeable foliar residue ($i_AppRate \cdot i_DFR$)	0,54 $\mu\text{g a.s./cm}^2$	<i>d_DFR</i>
Vapour pressure of in-use dilution	low volatile substances having a vapour pressure of $<5 \cdot 10^{-3}$ Pa	<i>i_Volat</i>
Concentration in air	0,001 mg/m^3	<i>d_AirCon</i>
Bystander dermal spray drift exposure - adult	12,9 ml spray dilution/person	
Bystander dermal spray drift exposure - child	3,87 ml spray dilution/person	
Bystander inhal. spray drift exposure - adult	0,00440 ml spray dilution/person	
Bystander inhal. spray drift exposure - child	0,00348 ml spray dilution/person	
Exposure duration	2 hours	<i>d_ByExpDur</i>
Exposure duration entry into treated crops	0,25 hours	<i>d_ExpDurTreatCrop</i>
Light clothing adjustment factor	18,0%	<i>d_ClothAF</i>
Breathing rate adult	0,23 $\text{m}^3/\text{hours/kg}$	<i>d_BreathRAD</i>
Breathing rate child (1-3 year old)	1,07 $\text{m}^3/\text{hours/kg}$	<i>d_BreathRCh</i>
Drift percentage on surface (90th percentile)	19,89%	
Turf transferable residues percentage	5,00%	<i>d_Turf</i>
Transfer coeff. of surface deposits-adult	14500 cm^2/hour	<i>d_ByTCAd</i>
Transfer coeff. of surface deposits-child (1-3 year old)	5200 cm^2/hour	<i>d_ByTCCh</i>
Saliva extraction percentage	50,00%	<i>d_SalExt</i>
Surface area of hands mouthed	20 cm^2	<i>d_AreaHM</i>
Frequency of hand to mouth activity	20 events/hour	<i>d_ByFreqHM</i>
Ingestion rate for mouthing of grass per day	25 cm^2	<i>d_MouthGrass</i>
Dislodgeable residues percentage transferability for object to mouth	20,00%	<i>d_DRP</i>
Transfer coefficient for entry into treated crops - adult	7500 cm^2/h	<i>d_TcEntryAd</i>
Transfer coefficient for entry into treated crops - child	2250 cm^2/h	<i>d_TcEntryCh</i>

1. Total

1.1 1-3 year old child

	Spray drift	Vapour	Surface deposits	Entry into treated crops
Total systemic exposure (mg a.s./day)	0,1154964	0,0107000	0,0339000	0,0379688
Total systemic exposure per kg body weight (mg/kg bw/day)	0,0115496	0,0010700	0,0033900	0,0037969
% of RVAAS	57,75%	5,35%	16,95%	18,98%

1.2 Adult

	Spray drift	Vapour	Surface deposits	Entry into treated crops
Total systemic exposure (mg a.s./day)	0,3823920	0,0138000	0,0648911	0,1265625

Total systemic exposure per kg body weight (mg/kg bw/day)	0,0063732	0,0002300	0,0010815	0,0021094
% of RVAAS	31,87%	1,15%	5,41%	10,55%

2. Details

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
1-3 year old child				
Spray drift	0,1154964	0,0115496	$((C16 * i_AbsorpInuse * (1 - d_ClothAF)) + C18) * d_ConcAS$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	0,0107000	0,0010700	$d_AirCon * d_BreathRCh * d_BwChild$	
Surface deposits				
Dermal	0,0232713	0,0023271	$(i_AppRate / 100) * C24 * d_Turf * d_ByTCCh * d_ByExpDur * MAX(i_AbsorpProduct, i_AbsorpInuse) * d_MAF * IF(i_AppEquip = "Vehicle-mounted-Drift Reduction", 0.5, 1)$	
Hand to mouth	0,0085030	0,0008503	$(i_AppRate / 100) * C25 * d_Turf * d_SalExt * d_AreaHM * d_ByFreqHM * d_ByExpDur * i_AbsorpOrallnuse * d_MAF$	
Object to mouth	0,0021257	0,0002126	$(i_AppRate / 100) * C25 * d_DRP * d_MouthGrass * i_AbsorpOrallnuse * d_MAF$	

Entry into treated crops				
Dermal	0,0379688	0,0037969	$(d_TcEntryCh*0.25*d_DFR*d_MAF)/1000*MAX(i_AbsorpProduct,i_AbsorpInuse)$	
Hand to mouth			$(i_AppRate/100)*d_MAF*d_Turf*d_SalExt*d_AreaHM*d_ByFreqHM*d_ByExpDur*i_AbsorpOrallnuse$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Object to mouth			$(i_AppRate/100)*d_DRP*d_MouthGrass*i_AbsorpOrallnuse*d_MAF$	Considered only for application on grassland and lawns and for application on golf course, turf or other sports lawns.
Adult				
Spray drift	0,3823920	0,0063732	$((C15*i_AbsorpInuse*(1-d_ClothAF)t)+C17)*d_ConcAS$	the only available values are for the 8 m distance downwind from the middle of the tree trunk, which are assumed to represent 5 m distance from the edge of orchard; the same value is used for 5 and 10 m.
Vapour	0,0138000	0,0002300	$d_AirCon*d_BreathRA*d_BwAdult$	
Surface deposits (dermal)	0,0648911	0,0010815	$(i_AppRate/100)*C24*d_Turf*d_ByTCA*d_ByExpDur*MAX(i_AbsorpProduct,i_AbsorpInuse)*d_MAF*IF(i_AppEquip="Vehicle-mounted-Drift Reduction",0.5,1)$	
Entry into treated crops (dermal)	0,1265625	0,0021094	$(d_TcEntryAd*0.25*d_DFR*d_MAF)/1000*MAX(i_AbsorpProduct,i_AbsorpInuse)$	

Recreational exposure for Calypso

Golf course, turf or other sports lawns		This sheet is only to be used for treatment of grassland used for recreational purposes	
Croptype			
Application method	Upward spraying		
Application equipment	Manual-Hand held		<i>i_AppEquip</i>
Formulation type	Soluble concentrates, emulsifiable concentrate, etc.		<i>i_FormVal</i>
Application rate of the product	0,18 kg a.s./ha		<i>i_AppRate</i>
Dermal absorption of product	1,00%		<i>i_AbsorpProduct</i>
Dermal absorption of in-use dilution	10,00%		<i>i_Absorplnuse</i>
Oral absorption	100,00%		<i>i_AbsorpOrallnuse</i>
Dislodgeable foliar residue ($i_AppRate \cdot i_DFR$)	0,54 $\mu\text{g a.s./cm}^2$		<i>d_DFR</i>
Exposure duration dermal	2 hours		<i>d_ReExpDur</i>
Light clothing adjustment factor Adult resident	18,0%		<i>d_ClothAF</i>
Drift percentage on surface	100,00%		
Turf transferable residues percentage	5,00%		<i>d_Turf</i>
Transfer coeff. of surface deposits-adult	7300 cm^2/hour		<i>d_ReTCAd</i>
Transfer coeff. of surface deposits-child (1-3 year old)	2600 cm^2/hour		<i>d_ReTCCh</i>
Saliva extraction percentage	50,00%		<i>d_SalExt</i>
Surface area of hands mouthed	20 cm^2		<i>d_AreaHM</i>
Frequency of hand to mouth activity	9,5 events/hour		<i>d_ReFreqHM</i>
Ingestion rate for mouthing of grass per day	25 cm^2		<i>d_MouthGrass</i>

2. Details

	Systemic exposure [mg a.s. /day]	Systemic exposure [mg a.s./kg bw/day]	Formula	Comments
1-3 year old child				
Surface deposits				
Dermal	0,0585000	0,0058500	$(i_AppRate/100) \cdot C13 \cdot d_Turf \cdot d_ReTCCh \cdot d_ReExpDur \cdot \text{MAX}(i_AbsorpProduct, i_Absorplnuse) \cdot d_MAF$	
Hand to mouth	0,0203063	0,0020306	$(i_AppRate/100) \cdot C13 \cdot d_Turf \cdot d_SalExt \cdot d_AreaHM \cdot d_ReFreqHM \cdot d_ReExpDur \cdot i_AbsorpOrallnuse \cdot d_MAF$	
Object to mouth	0,0106875	0,0010688	$(i_AppRate/100) \cdot C13 \cdot d_DRP \cdot d_MouthGrass \cdot i_AbsorpOrallnuse \cdot d_MAF$	
Total systemic exposure	0,0894938	0,0089494		
% of RVNAS				
Adult				
Surface deposits (dermal)	0,1642500	0,0027375	$(i_AppRate/100) \cdot C13 \cdot d_Turf \cdot d_ReTCAd \cdot d_ReExpDur \cdot \text{MAX}(i_AbsorpProduct, i_Absorplnuse) \cdot d_MAF$	
% of RVNAS				

<i>d_AirConVol</i>	Concentration in air of moderately volatile substances	0,015 mg/m ³
<i>d_AirConNonVol</i>	Concentration in air of low volatile substances	0,001 mg/m ³
<i>d_AreaHM</i>	Surface area of hands mouthed	20 cm ²
<i>d_AreaTreated</i>	Area treated (defined by crop type)	4 ha
<i>d_BreathRAd</i>	Breathing rate adult residents	0,23 m ³ /day/kg
<i>d_BreathRCh</i>	Breathing rate child (1-3 year old) residents	1,07 m ³ /day/kg
<i>d_BwAdult</i>	Adult body weight	60 kg
<i>d_BwChild</i>	Child body weight (1 to < 3 year olds)	10 kg
<i>d_ByBreathRAd</i>	Breathing rate adult bystander	0,04 m ³ /hours/kg
<i>d_ByBreathRCh</i>	Breathing rate child (1-3 year old) bystander	0,19 m ³ /hours/kg
<i>d_ByExpDur</i>	Exposure duration intense activity breathing rates	2 hours
<i>d_ByFreqHM</i>	Frequency of hand to mouth activity	20 events/hour
<i>d_ByTCAd</i>	Transfer coeff. of surface deposits-adult	14500 cm ² /hour
<i>d_ByTCCh</i>	Transfer coeff. of surface deposits-child (1-3 year old)	5200 cm ² /hour
<i>d_ClothAF</i>	Light clothing adjustment factor resident and bystanders	18,0%
<i>d_ConcAs</i>	Concentration of active substance (in-use dilution for liquid applications)	0,36 g a.s./l
<i>d_DFR</i>	Dislodgeable foliar residue (i_AppRate*i_DFR)	0,54 µg a.s./cm ²
<i>d_DRP</i>	Dislodgeable residues percentage transferability for object to mouth	20,0%
<i>d_HalfLifeAS</i>	Half-life of active substance (DT50)	30 days
<i>d_InhalTcAut</i>	Inhalation transfer coefficient for automated applications	NA ha/hr*10 ^{^(-3)}
<i>d_InhalTcCut</i>	Inhalation transfer coefficient for cutting ornamentals	NA ha/hr*10 ^{^(-3)}
<i>d_InhalTcSort</i>	Inhalation transfer coefficient for sorting / bundling ornamentals	NA ha/hr*10 ^{^(-3)}
<i>d_MAF</i>	Multiple application factor	1,25
<i>d_MouthGrass</i>	Ingestion rate for mouthing of grass per day	25 cm ² grass/day
<i>d_ReExpDur</i>	Exposure duration resident dermal	2 hours
<i>d_ReExpDurInhal</i>	Exposure duration resident inhalation	24 hours
<i>d_ExpDurTreatCrop</i>	Exposure duration for resident and bystander entry into treated crops	0,25 hours
<i>d_ReFreqHM</i>	Frequency of hand to mouth activity	9,5 events/hour
<i>d_ReTCAd</i>	Transfer coeff. of surface deposits-adult	7300 cm ² /hour
<i>d_ReTCCh</i>	Transfer coeff. of surface deposits-child (1-3 year old)	2600 cm ² /hour
<i>d_SalExt</i>	Saliva extraction percentage	50,0%
<i>d_TcEntryAd</i>	Transfer coefficient for entry into treated crops 75th percentile - adult	7500 cm ² /h
<i>d_TcEntryCh</i>	Transfer coefficient for entry into treated crops 75th percentile - child	2250 cm ² /h
<i>d_TcEntryMeanAd</i>	Transfer coefficient for entry into treated crops mean - adult	5980 cm ² /h
<i>d_TcEntryMeanCh</i>	Transfer coefficient for entry into treated crops mean - child	1794 cm ² /h
<i>d_Turf</i>	Turf transferable residues percentage	5,0%
<i>d_PctExtrapolation</i>	For exposure value 75 percentiles above this amount linear extrapolation is performed	1,5 kg
<i>d_head75ProtectionFactor</i>	Coefficient to estimate head protection factor 75 th Percentile	1,79422
<i>d_head95ProtectionFactor</i>	Coefficient to estimate head protection factor 95 Percentile	1,24705

sys_KeyOperator Variables for operator exposure lookup key
sys_OperatorModel Operator model

i_IndoorOutdoor&i_FormVal&i_AppMeth&i_AppEquip&

1

RPE reduction factor	
key_MixRPE, ay_MixRPE	
None	1
FP1, P1 and similar	0,25
FP2, P2 and similar	0,1

PPE reduction factor	
key_MixPPEBody, ay_MixPPEBody	
Potential exposure	1
Work wear - arms, body and legs covered	0,1
Certified protective coverall	0,05

PPE reduction factor	
key_MixPPEHead, ay_MixPPEHead	
None	1
Hood	0,5
Hood and visor	0,05
FP1, P1 and similar	0,8
FP2, P2 and similar	0,8

Application: Gloves PPE reduction factor (depending on formulation type)		
key_AppPPEHands, ay_AppPPEHands		
Wettable powder, soluble powder	Chemical resistant gloves	0,05
Granules, fine granules	Chemical resistant gloves	0,05
Wettable granules, soluble granules	Chemical resistant gloves	0,05
Soluble concentrates, emulsifiable concentrate, etc.	Chemical resistant gloves	0,1
Wettable powder, soluble powder	None	1
Granules, fine granules	None	1
Wettable granules, soluble granules	None	1
Soluble concentrates, emulsifiable concentrate, etc.	None	1

Crop dependent exposure parameters										
key_CropType, ay_CropType	Transfer coefficients	Transfer coefficients	2250	Transfer coefficients	Area Treated					
Crop type	Arm, body and legs covered	Total potential exposure	Activity	hours per day	Body parts involved	Hands, arm, body and legs covered	Type of crop for Resident Bystander	Vehicle Mounted Applications		
Bare soil	NA	NA	NA	NA	NA	NA	Field crops			50
Low berries and other small fruits		3000	5800 Reaching, picking		8 Hand and forearm		750 Field crops			50
Brassica vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops			50
Bulb vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops			50
Cane fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Field crops			10
Cereals		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops			50
Citrus fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops			10
Fruiting vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops			50
Grapes		10100	30000 Hand harvesting		8 Hand and body	no TC available for this assessment	Grapes			10
Grassland and lawns		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops			50
Golf course, turf or other sports lawns		2500	5800 Maintenance		8 Hand and body		580 Field crops			50
Hops		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Hops			50
Leaf vegetables and fresh herbs		2500	5800 Reaching, picking		8 Hand and body		580 Field crops			10
Legume vegetables		2500	5800 Reaching, picking		8 Hand and body		580 Field crops			50
Oilfruits		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops			10
Oilseeds		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops			50
Ornamentals		5000	14000 Cutting, sorting, bundling, carrying		8 Hand and body		1400 Field crops			10
Pome fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops			10
Root and tuber vegetables		1400	12500 Inspection, irrigation		2 Hand and body	no TC available for this assessment	Field crops			50
Stone fruit		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops			10
Tree nuts		4500	22500 Searching, reaching, picking		8 Hand and body		2250 Fruit crops			10

Resident Spray Drift					
These values are the 75th Percentiles for Residents (assuming average breathing rates for inhalation exposures)					
key_ResidSpray, ay_ResidSpray	Adults Dermal	Children Dermal	Adults Inhalation	Children Inhalation	
Downward spraying2-3		0,47	0,327	0,0001	0,00022
Downward spraying5		0,24	0,22	0,00009	0,00017
Downward spraying10		0,20	0,18	0,00009	0,00013
Upward spraying2-3	NA	NA	NA	NA	
Upward spraying5		5,63	1,689	0,0021	0,00164
Upward spraying10		5,63	1,689	0,0021	0,00164

Bystander Spray Drift					
These values are the 95th Percentiles for Bystanders (assuming high breathing rates for inhalation exposures)					
key_BySpray, ay_BySpray	Adults Dermal	Children Dermal	Adults Inhalation	Children Inhalation	
Downward spraying2-3		1,21	0,74	0,0005	0,0011
Downward spraying5		0,57	0,48	0,00048	0,0008
Downward spraying10		0,48	0,39	0,00051	0,00076
Upward spraying2-3	NA	NA	NA	NA	
Upward spraying5		12,9	3,87	0,0044	0,0035
Upward spraying10		12,9	3,87	0,0044	0,0035

Mean Spray Drift					
These values are the mean values (assuming average breathing rates for inhalation exposures)					
key_AvgSpray, ay_AvgSpray	Adults Dermal	Children Dermal	Adults Inhalation	Children Inhalation	
Downward spraying2-3		0,22	0,18	0,0001	0,0002
Downward spraying5		0,12	0,12	0,0001	0,0001
Downward spraying10		0,11	0,1	0,0001	0,0001
Upward spraying2-3	NA	NA	NA	NA	
Upward spraying5		3,68	1,11	0,0017	0,0013
Upward spraying10		3,68	1,11	0,0017	0,0013

Resident and bystander Surface Deposits Drift percentage				
Ground sediments in % of the application rate calculated on the basis of percentile values (drift data acc. Rautmann)				
key_ByCropType, ay_ByCropType	Bystander surface deposit (90th Percentile)	Resident surface deposit (77th Percentile)	mean	
Field cropsnot relevant2-3		0,085	0,056	0,041
Field cropsnot relevant5		0,035	0,023	0,018
Field cropsnot relevant10		0,019	0,013	0,010
Fruit cropsnot relevant2-3		0,292	0,240	0,190
Fruit cropsnot relevant5		0,199	0,158	0,117
Fruit cropsnot relevant10		0,118	0,090	0,061
Fruit cropsearly (without leaves)2-3		0,292	0,240	0,190
Fruit cropsearly (without leaves)5		0,199	0,158	0,117

Fruit cropsearly (without leaves)10	0,118	0,090	0,061
Fruit cropslate (dense foliage)2-3	0,157	0,110	0,070
Fruit cropslate (dense foliage)5	0,084	0,060	0,037
Fruit cropslate (dense foliage)10	0,036	0,027	0,016
Grapesnot relevant2-3	0,080	0,069	0,053
Grapesnot relevant5	0,036	0,031	0,023
Grapesnot relevant10	0,012	0,010	0,008
Hopsnot relevant2-3	0,193	0,159	0,100
Hopsnot relevant5	0,116	0,086	0,059
Hopsnot relevant10	0,058	0,037	0,029

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Mixing Method	Outdoor/Indoor	Formulation type	Application method	Application equipment	Type of exposure	Mixing & Loading 75th percentile	Mixing & Loading 95th percentile	Mixing & Loading Comments	Mixing & Loading Model	Application 75th percentile	Application 95th percentile	Application Comments	Application Model
IndoorGranules, fine granulesApplication of granulesManualBody	Indoor	Granules, fine granules	Application of granules	Manual	Body			Value for application is for combination of mixing&loading and application	PHED	68,8708	253,4438	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
IndoorGranules, fine granulesApplication of granulesManualHands	Indoor	Granules, fine granules	Application of granules	Manual	Hands			Value for application is for combination of mixing&loading and application	PHED	26,5320	94,3636	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
IndoorGranules, fine granulesApplication of granulesManualInhalation	Indoor	Granules, fine granules	Application of granules	Manual	Inhalation			Value for application is for combination of mixing&loading and application	PHED	0,4677	1,5251	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesBroadcast application of granulesVehicle-mountedBody	Outdoor	Granules, fine granules	Broadcast application of granules	Vehicle-mounted	Body	0,0162	0,0427	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0,0047	0,0151	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesBroadcast application of granulesVehicle-mountedHands	Outdoor	Granules, fine granules	Broadcast application of granules	Vehicle-mounted	Hands	0,0015	0,0069	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0,0004	0,0018	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesBroadcast application of granulesVehicle-mountedInhalation	Outdoor	Granules, fine granules	Broadcast application of granules	Vehicle-mounted	Inhalation	0,0208	0,0784		PHED	0,0012	0,0045		PHED
OutdoorGranules, fine granulesIn furrow application of granulesVehicle-mountedBody	Outdoor	Granules, fine granules	In furrow application of granules	Vehicle-mounted	Body	0,0162	0,0427	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0,0047	0,0151	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesIn furrow application of granulesVehicle-mountedHands	Outdoor	Granules, fine granules	In furrow application of granules	Vehicle-mounted	Hands	0,0015	0,0069	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED	0,0004	0,0018	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesIn furrow application of granulesVehicle-mountedInhalation	Outdoor	Granules, fine granules	In furrow application of granules	Vehicle-mounted	Inhalation	0,0208	0,0784		PHED	0,0012	0,0045		PHED
OutdoorGranules, fine granulesManual application of granulesManualBody	Outdoor	Granules, fine granules	Manual application of granules	Manual	Body			Value for application is for combination of mixing&loading and application	PHED	68,8708	253,4438	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesManual application of granulesManualHands	Outdoor	Granules, fine granules	Manual application of granules	Manual	Hands			Value for application is for combination of mixing&loading and application	PHED	26,5320	94,3636	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED
OutdoorGranules, fine granulesManual application of granulesManualInhalation	Outdoor	Granules, fine granules	Manual application of granules	Manual	Inhalation			Value for application is for combination of mixing&loading and application	PHED	0,4677	1,5251	Exposure value originally included use of PPE, calculated potential exposure is 100 times higher assuming a 'worst case' reduction factor of 1% for gloves/coverall	PHED

